



**Efficacy of Perineural versus Systemic
Dexamethasone to prolong analgesia after
Supraclavicular Nerve Block in patients
undergoing Wrist Arthroscopy**

Thesis

*Submitted for Partial Fulfilment of Master
Degree in Anaesthesia*

By

Ahmed Ali Nadeem Hussein Refaat

M.B.B.Ch., Faculty of Medicine, Ain Shams University

Supervised by

Prof. Dr. Alaa Eid Mohammed

Professor of Anaesthesia, ICU & Pain Management

Faculty of Medicine- Ain Shams University

Prof. Dr. Ayman Ahmed Abdellatif

Assistant Professor of Anaesthesia, ICU & Pain Management

Faculty of Medicine- Ain Shams University

Prof. Dr. Hoda Shokri Abd ElSamie

Assistant Professor of Anaesthesia, ICU & Pain Management

Faculty of Medicine- Ain Shams University

*Faculty of Medicine
Ain Shams University
2018*

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

سبحانك لا علم لنا
إلا ما علمتنا إنك أنت
العليم العظيم

صدق الله العظيم

سورة البقرة الآية: ٣٢

Acknowledgment

*First and foremost, I feel always indebted to **ALLAH**, the Most Kind and Most Merciful.*

*I'd like to express my respectful thanks and profound gratitude to **Prof. Dr. Alaa Eid Mohammed**, Professor of Anaesthesia, ICU & Pain Management Faculty of Medicine- Ain Shams University for his keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.*

*I am also delighted to express my deepest gratitude and thanks to **Prof. Dr. Ayman Ahmed Abdellatif**, Assistant Professor of Anaesthesia, ICU & Pain Management Faculty of Medicine- Ain Shams University, for his kind care, continuous supervision, valuable instructions, constant help and great assistance throughout this work.*

*I am deeply thankful to **Prof. Dr. Hoda Shokri Abd ElSamie**, Assistant Professor of Anaesthesia, ICU & Pain Management Faculty of Medicine- Ain Shams University, for her great help, active participation and guidance.*

Ahmed Ali Nadeem Hussein Refaat

List of Contents

Title	Page No.
List of Tables	5
List of Figures	6
List of Abbreviations	8
Introduction	1
Aim of the Work.....	11
Review of Literature	
▪ Anatomy of the Brachial Plexus	12
▪ Bupivacaine	29
▪ Dexamethasone.....	38
Patients and Methods	46
Results	56
Discussion	69
Conclusion.....	74
Summary	75
References	77
Arabic Summary	—

List of Tables

Table No.	Title	Page No.
Table (1):	Pharmacokinetics of dexamethasone	41
Table (2):	Comparison between groups according to demographic data & peri-operative times.....	56
Table (3):	Comparison between groups according to mean blood pressure.	60
Table (4):	Comparison between groups according to heart rate.....	62
Table (5):	Comparison between groups according to oxygen saturation.....	64
Table (6):	Comparison between groups according to numeric pain rating scale.	66
Table (7):	Comparison between groups according to total no. of rescue analgesic doses (Pethidine).....	68

List of Figures

Fig. No.	Title	Page No.
Figure (1):	Brachial plexus with terminal branches labeled.....	13
Figure (2):	Typical cervical vertebra	14
Figure (3):	Basic relationships of the brachial plexus to the axillary artery, which is a continuation of the subclavian artery.....	16
Figure (4):	Distribution of the musculocutaneous nerve	18
Figure (5):	Distribution of the ulnar nerve	19
Figure (6):	Distribution of the median nerve.....	20
Figure (7):	Distribution of axillary and radial nerves	21
Figure (8):	Anatomical relations of the brachial plexus	26
Figure (9):	Chemical structure of bupivacaine Hcl molecule	29
Figure (10):	Probe position.....	50
Figure (11):	Approach.	50
Figure (12):	Injection.....	51
Figure (13):	Numeric Rating Scale for pain (NRS pain).....	54
Figure (14):	Bar chart between groups according to age.....	57
Figure (15):	Bar chart between groups according to sex.	58
Figure (16):	Bar chart between groups according to ASA.....	58
Figure (17):	Bar chart between groups according to duration of surgery.	58
Figure (18):	Bar chart between groups according to time for first rescue analgesic.	59
Figure (19):	Comparison between groups according to mean blood pressure.....	61

List of Figures Cont...

Fig. No.	Title	Page No.
Figure (20):	Comparison between groups according to heart rate.	63
Figure (21):	Comparison between groups according to oxygen saturation.....	65
Figure (22):	Comparison between groups according to numeric pain rating scale.....	67
Figure (23):	Comparison between groups according to total no. of rescue analgesic doses.	68

List of Abbreviations

Abb.	Full term
CNS	Central nervous system
COX I	Cyclooxygenase I
COX II	Cyclooxygenase II
DNA	Deoxyribo Nucleic Acid
DST	Dexamethasone suppression test
HPA axis	Hypothalamus pituitary adrenal axis
HR	Heart rate
HS	Highly significant
IL-1.....	Interleukin 1
IL-6.....	Interleukin 6
IM.....	Intramuscular
IV.....	Intravenous
K+	Potassium
LAs	Local anaesthetics
MBP	Mean blood pressure
Na+.....	Sodium
NS.....	Non significant
S	Significant
TNF	Tumor necrosis factor

INTRODUCTION

Brachial plexus block is a popular and widely employed regional nerve block technique for peri-operative anaesthesia and analgesia for surgery of the upper extremity. Supraclavicular approach is the easiest and most consistent method for surgery below the shoulder joint. Regional nerve block minimizes the stress response to surgery and allows using minimal anaesthetic drugs which is always beneficial for the patients with various cardio-respiratory co-morbidities (*Shrestha et al, 2007*).

Ultrasound guidance is a reliable and safe technique in peripheral nerve blocks. It also plays a crucial and an increasing role in medicine nowadays due its low cost, the absence of ionizing radiation and its high temporal resolution (*Jerrold et al, 2010*).

Local anaesthetics alone for supraclavicular brachial plexus block provide good operative conditions, but they have short duration of postoperative analgesia. Therefore, various adjuvants such as opioids, clonidine, neostigmine and midazolam were added to local anaesthetics in brachial plexus block to achieve quick, dense and prolonged block (*Gowala et al, 2009*).

Bupivacaine is a widely used local anaesthetic which is related chemically and pharmacologically to the amide local

anaesthetics available. Using 0.5% bupivacaine in supraclavicular brachial plexus block, the onset time may be up to 20 min with surgical anaesthesia taking up to 40 min. The duration of block ranges from 4-12h with residual anaesthesia still being present at 24h (*Berde, 2015*).

Perineural dexamethasone was first explored clinically more than 12 years ago, followed by countless clinical trials. Recently, a meta-analysis concluded that perineural dexamethasone, compared with placebo, prolonged the duration of analgesia by >8h, when combined with long-acting local anaesthetics, suggesting that patients could benefit from a pain-free postoperative night. The mechanism of action for this prolongation of block is not fully understood, but suggested possibilities include decreased nociceptive C-fiber activity via a direct effect on glucocorticoid receptors (*Johansson et al, 1995*), a direct effect on inhibitory potassium channels, a local vasoconstrictive effect or systemic anti-inflammatory effect.

However, perineural route of dexamethasone administration remains off label. An alternative choice of intravenous dexamethasone has likewise been explored, which at moderate doses offers the potential for a systemic anti-inflammatory effect (*De Olivera et al, 2011*).

AIM OF THE WORK

The aim of this work is to study the effect of dexamethasone as an adjuvant to bupivacaine either intravenous or perineural in ultrasound guided supraclavicular brachial plexus block in prolongation of postoperative analgesia in patients undergoing wrist arthroscopy.

Chapter 1

ANATOMY OF THE BRACHIAL PLEXUS

Brachial plexus is a complex network of nerves supplying the whole upper limb, including its motor and sensory supply, arising from the neck and passing through the axilla to the upper limb. It is composed of 5 roots, 3 trunks, 6 divisions, 3 cords, and terminal branches (**Figure 1**) (*Andres and Sala, 2001*).

1. Roots:

The ventral rami of spinal nerves from C5 to T1 are referred to as the roots of the brachial plexus. The typical spinal nerve root results from the union of the ventral nerve rootlets originating in the anterior horn cells of the spinal cord and the dorsal nerve rootlets (*Andres and Sala, 2001*).




Figure (1): Brachial plexus with terminal branches labeled: MC is musculocutaneous nerve, AXI is axillary nerve, RAD is radial nerve, MED is median nerve, and ULN is ulnar nerve (*Andres and Sala, 2001*).

The roots then get exit through the transverse processes of the cervical vertebrae just posterior to the vertebral artery, which runs in a cephalic direction through the transverse foraminae. Each transverse process consists of a posterior and anterior tubercle, which meet laterally to form the costotransverse bar (**Figure 2**). The transverse foramen lies medial to the costotransverse bar and between the posterior and anterior tubercles. The spinal nerves that form the brachial plexus run in an inferior and anterior direction within the sulci which is formed by these structures (*Gloss et al, 2006*).




Figure (2): Typical cervical vertebra (*Gloss et al., 2006*).

2. Trunks:

Shortly after emerging from the intervertebral foraminae, the 5 roots (C5-T1) unite to form 3 trunks. One on top of the other, as they traverse the triangular interscalene space formed between the anterior and the middle scalene muscles. This space becomes wider in the anteroposterior plane as the muscles approach their insertion on the first rib. Although the roots of the plexus are long, the trunks are as short (approximately 1 cm) as they are wide, which shortly give rise to a total of six divisions (three anterior and three posterior), as they reach the clavicle (*Franco and Vieira, 2008*).

The area of the trunks corresponds to the point where the brachial plexus is confined to its smallest surface area, 3 nerve structures, closely related to each other, carrying the entire sensory, motor and sympathetic innervation of the upper

extremity, with the exception of a small area in the axilla and upper middle arm, which is innervated by the intercosto-brachial nerve, a branch of the 2nd intercostal nerve. This great reduction in surface area allows the plexus to negotiate the narrow passage between the clavicle and the first rib at the apex of the axilla (*Singhal et al, 2007*).

The ventral rami of C5 and C6 unite to form the upper (superior) trunk, from which the suprascapular nerve and the nerve to the subclavius arise. The suprascapular nerve carries sensory fibers to the shoulder joint and provides motor innervation to the supraspinatus and infraspinatus muscles. The ventral ramus of C7 continues as the middle trunk. The ventral rami of C8 and T1 unite to form the lower (inferior) trunk (**Figure 3**) (*Fazan et al, 2001*).