MYCORHIZAE AND RHIZOBACTERIA FOR MITIGATING SALT STRESS IN SOME WOODY TREES

By

HANAA SHERIF ELSAYED AHMED

B. Sc. Agric. Sci. (Microbiology), Fac. Agric., Ain Shams Univ., 2001

THESIS

Submitted in Partial Fulfillment of the Requirements for the Degree of

MASTER OF SCIENCE

In

Agricultural Sciences (Agricultural Microbiology)

Department of Agricultural Microbiology
Faculty of Agriculture
Cairo University
EGYPT

2018

APPROVAL SHEET

MYCORHIZAE AND RHIZOBACTERIA FOR MITIGATING SALT STRESS IN SOME WOODY TREES

M. Sc. Thesis
In
Agric. Sci. (Agricultural Microbiology)

By

HANAA SHERIF ELSAYED AHMED

B. Sc. Agric. Sci. (Microbiology), Fac. Agric., Ain Shams Univ., 2001

APPROVAL COMMITTEE

Dr. DIAA EL DIN MOHAMED SWEILAM	
Head Researcher of Soil Microbiology, SWERI, ARC, Giza.	
Dr. NADIA FAHMY AMIN EMAM	
Professor of Agricultural Microbiology, Fac. Agric., Cairo University.	
Dr. ESSAM MOHAMED ABDEL-MOATY DARWISH	
Associate Professor of Plant Physiology, Fac. Agric., Cairo Universit	y
Dr. MOHAMED ABDEL-ALIM ALI	
Professor of Agricultural Microbiology, Fac. Agric., Cairo University.	

Date: 18 / 7 / 2018

SUPERVISION SHEET

MYCORHIZAE AND RHIZOBACTERIA FOR MITIGATING SALT STRESS IN SOME WOODY TREES

M. Sc. Thesis
In
Agric. Sci. (Agricultural Microbiology)

By

HANAA SHERIF ELSAYED AHMED

B. Sc. Agric. Sci. (Microbiology), Fac. Agric., Ain Shams Univ., 2001

SUPERVISION COMMITTEE

Dr. MOHAMED ABDEL-ALIM ALI Professor of Agricultural Microbiology, Fac. Agric., Cairo University

Dr. ESSAM MOHAMED ABDEL-MOATY DARWISH Associate Professor of Plant Physiology, Fac. Agric., Cairo University

Dr. ESSAM MOSTAFA ABD EL-KADER Researcher - Horticulture Research Institute, ARC, Giza

Name of Candidate: Hanaa Sherif Elsayed Ahmed Degree: M. Sc.

Title of Thesis: Mycorhyzae and Rhizobacteria for Mitigating Salt Stress in

Some Woody Trees.

Supervisors: Dr. Mohamed Abdel Alim Ali

Dr. Essam Mohamed Abdel Moaty Darwish

Dr. Essam Mostafa Abdel Kader

Department: Agricultural Microbiology. **Approval**: 18 / 7/2018

ABSTRACT

This study included the following points: The first one concerning isolation and identification of plant growth promoting rhizobacteria (PGPR). Twelve bacterial isolates were isolated from rhizosphere of taxodium seedlings. Data revealed that six isolates were cocci, three isolates were short rods, two isolates were long rods in chains and one was long rods. All isolates were Gram positive except one isolates was Gram negative bacteria. All isolates tolerate salinity up to 12000 ppm. IAA production in there isolates was ranged from 3.85 to 7.59 mg/L. Polysaccharides ranged from 0.1609 to 12.226 mg/L. The highest two producers were characterized as: G+ cocci; 7.165 mg/L IAA; 12.226 mg/L polysaccharides; followed by G- short rods, 7.592 mg/L IAA, 8.364 mg/L polysaccharides. The two isolates were identified using Biolog as Kocuria varians and Enterobacter cloacae, respectively. The second point two PGPR isolates were used with or without mycorrhiza to alleviate salt stress on seedlings of three woody tree species (Taxodium deisticum, Khaya senegalensis and Balanites aegyptiaca). Salinity levels were 0.0, 2000, 4000 and 8000 ppm. Data revealed that all Balanites aegyptiaca seedlings exposed to 8000 ppm of saline were dead while 25% were able to stilled alive at 4000 ppm when infected with mycorrhizae. Survival percentage of Khaya senegalensis and Taxodium deisticum seedlings were 69.45% and 54.76%, respectively under 8000 ppm salinity when inoculated with mycorrhizae. All growth parameters (i.e), plant height; number of leaves, fresh and dry weight was decreased by increasing salinity level. Chlorophyll content, sodium, potassium, chloride, nitrogen and phosphorus were also investigated.

Key words: salinity, PGPR, mycorrhiza, *Taxodium deisticum, Khaya senegalensis* Balanites aegyptiaca.

DEDICATION

This work is dedicated to whom my heartfelt thanks; to my mother, my father, my brothers, my sisters and my friends for their patience, help and for all the support they lovely offered during my postgraduation studies.

ACKNOWLEDGEMENT

Praise and thanks be to ALLAH, for assisting and directing me to the right way.

The authoress wishes to express her sincere thanks, deepest gratitude and appreciation to **Dr.Mohamed A. Ali,** Professor of Agricultural Microbiology, Faculty of Agricultur, Cairo University for suggesting the problem, his supervision, kind assistance, continuous guidance and encouragement throughout study. Without his assistance, none of this work would have been possible. My deep thanks to **Dr. Essam Mohamed Abdel Moaty,** Associate Professor of plant Physiology Faculty of Agriculture, Cairo University for his encouragement, support, sincere advices and helpful supervision. I also would like to express my thanks, deepest gratitude to **Dr. Essam Mostafa Abdel Kader,** researcher at Horticulture Research Institute. ARC, Giza.

CONTENTS

INTRODUCTION
REVIEW OF LITERATURE
1. Salinity
2. PGPR and Mycorrhizae
3. Woody trees
MATERIALS AND METHODS
RESULTS
1. Plant growth promoting rhizobacteria (PGPR)
2. Effect of inoculation with PGPR on seedlings growth and
salinity tolerance
a. Balanites aegyptiaca
(1) Plant survival percentage
b. Khaya senegalensis
(1) Survival percentage
(2) Plant height, number of leaves, stem diameter and root length
(3) Shoot fresh weight, root fresh weight, shoot dry
weight and root dry weight
(4) Moisture content, IAA and proline
(5) Chlorophyll a, b and carotenoid
(6) N, P and K
(7) Na and Cl
c. Taxodium distichum
(1) Survival percentage
(2) Plant height, number of leaves, stem diameter and root length
(3) Shoot fresh weight, root fresh weight, shoot dry weight and root dry weight
(4) Moisture content, IAA and proline
(5) Chlorophyll a, b and carotenoid
(6) N, P and K

(7) Na and Cl	
DISCUSSION	,
PGPR as a tool to alleviate salt stress	
1. Effect of inoculation with PGPR on growth characters	
2. Chemical composition	
3. Growth parameters	
4. Chemical composition parameters	
SUMMARY	
REFERENCES	
ARABIC SUMMARY	