

بِسْمِ اللّٰهِ الرَّحْمٰنِ الرَّحِیْمِ





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغييرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بعض الوثائق الأصلية تالفة





بالرسالة صفحات لم ترد بالأصل



**MODIFIED PUBO-VAGINAL SLING
IN COMPLICATED FEMALE
URINARY INCONTINENCE**

THESIS

*Submitted to the faculty of Medicine Menoufya
University*

In partial fulfillment of the requirements of the degree of

**DOCTOR OF UROLOGICAL
SURGERY**

By

Alaa El-Din Mahmoud Shaaban

*MBBCh, Alex
MCH Ur Alex*

*Faculty of Medicine
Menoufya University*

1994

B

11717

SUPERVISORS

Prof. Dr. Gamal M. Ghoniem
MD., F.A.C.S.

Associate Professor of Urology and Gynecology
Chief, Section of Neurourology and Female Urology
Residency Program Director
Tulane University School of Medicine
New Orleans, Louisiana, USA

Prof. Dr. Mohamed Ali El-Sayed
Professor and Chairman
Department of Urology
Faculty of Medicine
Menoufya University

Prof. Dr. Nagi Ibrahim Eid
Professor of Urology
Department of Urology
Faculty of Medicine
Alexandria University

ACKNOWLEDGMENT

Praise to be done to ALLAH most merciful and most compassionate

I would like to express my profound gratitude and deep appreciation to Professor Gamal M. Ghoniem Associate Professor of Urology and Gynecology Chief, Section of Neurourology and Female Urology Tulane University School of Medicine, for suggesting the research protocol, support, meticulous guidance, and generous cooperation since the early stages of this work until now.

I am deeply grateful to Prof. Mohamed Ali El-Sayed Professor and Chairman, Department of Urology Faculty of Medicine, Menoufya University, for his support, advice , helpful instructions cooperation and guidance.

I am very grateful to Prof. Nagi Ibrahim Eid Professor of Urology, Department of Urology, Faculty of Medicine Alexandria University, for his help and encouragement that made the accomplishment of this work possible.

It is a pleasure to express my regard to Prof. Shoukri M. Nassef Professor of urology Faculty of Medicine, Menoufya University, for his support and guidance during the early stages of this work.

I am very grateful to Debbie Dubuc and Mike W. Aertker from the Urology Department, Tulane Regional Primate Center, for their technical and moral support, friendship and encouragement that made the accomplishment of this work and others possible.

Last but not least, I would like to thank Prof. Pamela J. Moore Prof. of Anatomy, Vice Dean, Tulane University, for her help and valuable guidance.

LIST OF TABLES

No.	Title	Page
1	Structures potentially involved in continence	32
2	Micturition reflexes	58
3	Lower urinary tract causes of established incontinence	69
4	Results of anterior colporaphy	102
5	The postoperative bladder drainage periods	140
6	A comparison between the patients who underwent MPVS as the only procedure, and the patients who underwent MPVS in conjunction with pelvic surgery	144
7	A comparison between the pre and postoperative complications in MPVS group and MPVS + pelvic surgery group	145
8	The patient complaints	158
9	Physical examination and investigations	158
10	Operative data	159
11	Analysis of DI	159
12	Number of strips used at different level of stretch	162
13	Major contribution in sling procedures	174
14	Sling materials	178
15	Different follow-up periods	197
16	Comparison between organic and inorganic slings	201
17	Detrusor instability among 80 patients	206
18	The effect of preoperative DI on the presence of postoperative DI	208

LIST OF FIGURES

No.	Title	Page
1	The pelvic side wall	5
2	Space of Retzus	7
3	The Pubovesical muscle	12
4	The Pubourethral ligament	14
5	Pelvic outlet	19
6	The striated urogenital sphincter	26
7	The component and location of the internal and external urogenital sphincter	27
8	Cerebral-brain stem circuit (LOOP I)	36
9	The brain stem sacral loop (LOOP II)	38
10	Vesical-sacral sphincter loop (LOOP III)	40
11	Cerebral-sacral loop (LOOP IV)	41
12	The skeletal muscle motor end plate	47
13	Storage-favoring reflexes (SDIR) + (SSCR)	49
14	Storage-favoring reflexes (PDIR)	50
15	Urethosphincteric guarding reflex (USGR)	52
16	Perineobulber detrusor facilitative reflex	53
17	Detrusodetrusor facilitative reflex (DDFR)	54
18	Detrusourethral inhibitory reflex (DUIR)	56
19	Schematic drawing of the Base Plate A: Bladder neck close during storage phase B: Bladder neck opening during voiding	63
20	Type 0 SUI	72
21	Type I SUI	73
22	Type IIA SUI	74
23	Type IIB SUI	75
24	Type III SUI	77

25	Anterior colporaphy A: Kelly plication, B: Kelly-Kennedy modification	101
26	A: Burch colposuspension (Cooper's ligament), B: Marshall-Marchetti-Krantz procedure (symphysis pubis), C: Richardson paravaginal approach (ATFP)	106
27	Three variations of NUP	106
28	A diagram for the AS 800 (artificial urinary sphincter)	109
29	Distribution of patients at the different hospitals	112
30	Age groups of the patients	
31	Weight groups of the patients (Pound)	114
32	The 7.5 X 2.5 cm fascial flap	118
33	Rectus fascia sling with the number 1 polypropylene suture at each end	119
34	An index finger develops the periurethral avascular space. The fingertip is directed against the inferior posterolateral pubic ramus, penetrating the urogenital diaphragm at its attachment	121
35	A: The vaginal finger positioned retropubically makes contact with the tip of the carrier needle. This contact is maintained as the needle is advanced and the finger withdrawn. B: The ends of the sutures are threaded through the aperture in the emerging needle carrier.	122
36	Passage of the sutures to the suprapubic area	123
37	Positioning and fixation of the sling	124
38	A diagram of the tissue bath used for measuring tensile strength of rectus fascia/vaginal wall	128
39	Complaint of the patients	133
40	Different previous anti-incontinence operations (total = 86)	134
41	Different presentation for pelvic floor weakness	135
42	Pre-operative urethral closure pressure (cm H ₂ O)	137
43	Different types of reconstructive pelvic surgeries (48 procedures)	139

44	The outcome of MPVS procedure	147
45	Age of the patients in the different outcome groups (mean + SD, p value)	148
46	Weight of the patients in the different outcome groups (mean + SD, p value)	149
47	Previous anti-incontinence surgeries in the different outcome groups (mean + SD, p value)	150
48	Amount of blood loss in the different outcome groups (mean + SD, p value)	151
49	Duration of bladder drainage following MPVS in the different outcome groups (mean + SD, p value)	152
51	Number of previous pregnancies in the different outcome groups (mean + SD, p value)	155
52	Number of previous deliveries in the different outcome groups (mean + SD, p value)	156
53	Number of pads used by the patients in the different outcome groups (mean + SD, p value)	157
54	A comparison between the weight of the different strips/their length (*RF=rectus fascia versus **VW=vaginal wall (mean and SD))	161
55	Absolute acute tension (in gram) rectus fascia versus vaginal wall at different levels of stretch	164
56	Absolute relaxed tension (in gram) rectus fascia versus vaginal wall	165
57	Acute tension (gram/100 mg tissue) rectus fascia versus vaginal wall	167
58	Relaxed tension (gram/100 mg tissue) rectus fascia versus vaginal wall	168
59	The time needed by the strips to relax (rectus fascia versus vaginal wall)	170
60	A comparison between the compliance of rectus fascia and vaginal wall strips at various stretch levels	171

ABBREVIATIONS AND SYMBOLS

ATFP	Arcus Tendineus Fasciae Pelvis
ATLA	Arcus Tendineus Levatory Ani
AUS	Artificial urinary sphincter
BUN	blood urea nitrogen
cm	centimeter
CMG	cystometrogram
CIC	clean intermittent catheterization
DI	detrusor instability
DSIR	detrusor sphincter inhibitory reflex
EMG	electromyography
FB	foreign body
FUDS	fluorourodynamic study
GET	general endotracheal
GSI	genuine stress incontinence
GSM	genitourinary spheroidal membrane
Hz	Hertz (frequency unite)
IC	interstitial cystitis
ISD	internal sphincteric dysfunction
mA	milliampere
mg	milligram
ml	milliliter
ml/min	milliliter/minute
mm	millimeter
MMK	Marshall-Marchetti-Krantz
MPVS	modified pubo-vaginal sling
m sec	millisecond
NUP	needle urethropexy

PC	personal computer
PMRC	Pontine micturition reflex center
PPA	phenylpropanolamine
PTR	pressure transmission ratio
PUV	posterior urethrovesical
PVM	pubovesical muscle
PVR	post voiding residual
RF	rectus fascia
RPV	retropubic urethropexy
SD	standard deviation
sec	second
SMRS	spinal micturition reflex center
SP	substance P
SUI	stress urinary incontinence
T	tension
U	urethral
UI	urge incontinence
UPP	urethral pressure profile
UTI	urinary tract infection
UVJ	urethrovesical junction
VVF	vesicovaginal fistula
VW	vaginal wall
#	number
Δt	delta t = change in time
Δv	delta v = change in volume

CONTENTS

No.	Chapter	Page
I.	INTRODUCTION	1-109
II.	AIM OF THE WORK	110
III.	PATIENT	111-114
IV.	METHODS	115-129
V.	RESULTS	131-171
VI.	DISCUSSION	173-215
VII.	SUMMARY AND CONCLUSION	217-218
VIII.	REFERENCES	220-256

PROTOCOL

ARABIC SUMMERY