

بسم الله الرحمن الرحيم









شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة يعيدا عن الغيار












بالرسالة صفحات لم ترد بالأصل





LEAST COST NUTRITIONALLY ADEQUATE DIET FOR AVERAGE EGYPTIAN FAMILY UNDER PREVAILING ECONOMIC CONDITIONS

BY

KHALED RAMADAN IBRAHIM MOHAMED

B.Sc., In Food Science and Nutrition Faculty of Home Economics Helwan University .1989

Thesis

Submitted in Partial Fulfillment of the Requirements for the Degree of M.Sc. (Nutrition)

SUPERVISOR

PROF. DR. AIDA MOHAMED EL-ASFAHANI

Professor of Nutrition and Food Science Faculty of Home Economics, Al-Menoufia University.

PROF. DR. AMIN ISMAIL ABDOU

Professor of Agricultural Economics National Research Center, Giza

FOOD SCIENCE AND NUTRITION DEPARTMENT FACULTY
OF HOME ECONOMICS, AL-MONOUFIA UNIVERSITY

1998

APPROVAL SHEET

LEAST COST NUTRITIONALLY ADEQUATE DIET FOR AVERAGE EGYPTIAN FAMILY UNDER PREVAILING ECONOMIC CONDITIONS

BY KHALED RAMADAN IBRAHIM MOHAMMED

B.Sc. Helwan University (1989)

APPROVED BY:

Aida El Arfahani Am Ami

Man Source

Committee in charge

Date:

25/1/1998

AL-MENOUFIA UNIVERSITY

1998



ACKNOWLEDGEMENT

I would like to express my deepest gratitude to Prof. Dr. Aida Mohamed El-Asfahani, Professor of Nutrition and Food Science Faculty of Home Economics, Al-Menoufia University, Shebin El-Kom, Egypt, for her articulate selection of the topic research coupled with her exceptional devotion, utmost effort she has shown throughout the thesis. Her constructive guidance together with her valued instructions and profound help were all of extreme importance in the verification of this study.

Moreover, I am truly indebted to Prof. Dr. Amin Ismail Abdou professor of agricultural Economics, National Research Center, Giza, Egypt, for his continuous assistance and encouragement. His instructions and remarks have been of great help. He spared no effort in bringing this work to the best possible results.

Khaled Ramadan

LIST OF ABBREVIATIONS

ALISH: Average Low-Income Strata Household.

CAPMAS: Central Agency for Public Mobilization and

Statistics.

FAO: Food and Agriculture Organization.

FBS: Food Balance Sheet.

FNB: Food and Nutrition Board.

HBS: Household Budget Survey.

LCAD : Least Cost Acceptable Diet.

RDA: Recommended Dietary Allowances.

UNU: United Nations University.

WHO: World Health Organization.

TABLE OF CONTENTS

	page
INTRODUCTION	1
REVIEW OF LITERATURE	3
1- Per capita consumption of major food groups.	3
2- Nutritional deficiencies.	7
3- Factors affecting food consumption levels and role of income.	11
4- Confrontation of the Nutritional problem in Egypt; applied	14
and suggested methods and programs.	
a) Target groups intervention.	14
b) Food subsidies.	16
MATERIAL AND METHODS.	17
RESULTS AND DISCUSSION.	20
I- General food consumption condition.	20
I.1. General nutritional status in Egypt compared with other	20
countries.	
I.2. Correlation between food per capita consumption and	21
income level.	
II- Target group characteristics.	27
II.1. Major characteristics of selected average low income	27
strata household (ALISH)	
II.2. The ALISH nutritional requirements.	27
II.3. ALISH food consumption patterns and nutrients intakes	29
II.4. Comparison between ALISH and other income brackets	36
food consumption levels.	
II.5. ALISH daily intake of nutrients compared to RDA.	36

	-V1-
II.6. ALISH food expenditure.	38
III- Diet planning.	
III.1. Basic concepts.	44
III.2. Elementary diet model for calories and protein	44
satisfaction.	
III.3. Suggested LCAD models for ALISH in urban regions.	45
III.4. Suggested LCAD models for ALISH in rural regions.	61
III.5. Summary: basic concepts and remarks.	69
III.6. Suggested LCAD models compared to actual food	78
intake of ALISH.	
III.6.1. Composition of LCAD models:	78
a)Comparison with HBS (1990/1991).	78
b)Comparison with FBS (1990/1991).	81
III.7. Nutritional value of ALISH LCAD models compared	85
with those of HBS and FBS.	
III.8. Costs.	88
SUMMARY AND CONCLUSION.	89
REFERENCES.	93
APPENDICES.	
ARABIC SUMMARY.	

LIST OF TABLES

			Page
Table 1	:	Per capita daily supply of calories, protein and share of	22
		vegetable sources in Egypt compared to other countries	
		(Average: 1961-1963, 1986-1988).	
Table 2	:	Per capita share of some major foods and total annual	26
		expenditure correlation coefficient, (1990/1991).	
Table 3	:	The ALISH sex and age composition in urban and rural	28
		Egypt, (1990/1991).	
Table 4	:	Recommended daily dietary allowances (RDA) for the	30
		ALISH in urban and rural areas, (1990/1991)	
Table 5	:	Average of ALISH food consumption and nutrients	32
		intakes, (1990/1991).	
Table 6		Food groups per capita daily consumption for ALISH and	37 .
		other brackets, (1990/1991).	
Table 7	:	Mean nutrients intakes of ALISH in (1990/1991)	39
		compared with RDA.	
Table 8	:	Average annual total expenditure and food share for the	43
		ALISH in urban and rural regions, (1990/1991).	
Table 9	:	The average minimum cost of suggested diet model for	46
		the ALISH.	
Table 10	:	Suggested LCAD model 1 for the ALISH in the urban	48
		regions, of the first half 1997.	
Table 11	:	Suggested LCAD model 2 for the ALISH in the urban	
		regions, of the first half 1997.	
Table 12	:	Suggested LCAD model 3 for the ALISH in the urban	50
		regions, of the first half 1997.	

Γable 13	:	Suggested LCAD model 4 for the ALISH in the urban	52
		regions, of the second half 1997.	
Table 14	:	Suggested LCAD model 5 for the ALISH in the urban	55
		regions, of the second half 1997.	
Table 15	:	Suggested LCAD model 6 for the ALISH in the urban	57
		regions, of the second half 1997.	
Table 16	:	Suggested LCAD model 7 for the ALISH in the rural	59
		regions, of the first half 1997.	
Table 17	:	Suggested LCAD model 8 for the ALISH in the rural	62
		regions, of the first half 1997.	
Table 18	:	Suggested LCAD model 9 for the ALISH in the rural	64
		regions, of the first half 1997.	
Table 19	:	Suggested LCAD model 10 for the ALISH in the rural	66
		regions, of the second half 1997.	
Table 20	:	Suggested LCAD model 11 for the ALISH in the rural	70
		regions, of the second half 1997.	
Table 21	:	Suggested LCAD model 12 for the ALISH in the rural	72
		regions, of the second half 1997.	
Table 22	:	Levels of consumption for the ALISH as reported by HBS	74
		(1990/1991) compared with those of suggested LCAD	
		models for the ALISH of (1997) in the urban regions.	80
Table 23	:	Levels of consumption for the ALISH as reported by HBS	
		(1990/1991) compared with those of suggested LCAD	
		models for the ALISH of (1997) in the rural regions.	82
Table 24	:	Daily per capita of food items reported by FBS of	
		(1990/1991) compared with those of suggested LCAD	
		models of (1997) in urban and rural regions.	84

Table 25 :	Per capita major nutrients intake and carbohydrate foods	87
	contribution in suggested LCAD models compared to	
	HBS and FBS food intake, (1990/1991).	
Table 26 :	Annual cost value of ALISH suggested LCAD models	88

and percentage share of total annual expenditure compared to actual patterns HBS of (1990/1991).

LIST OF FIGURES

		pag
Figure 1 :	Vegetable share as percentage of total caloric supply in Egypt	23
	and other countries (Average: 1961-1963, 1986-1988).	ė
Figure 2 :	Vegetable share as percentage of total protein supply in	24
	Egypt and other countries (Average: 1961-1963, 1986-1988).	
Figure 3 :	Mean nutrients intakes as percentage of RDA for the ALISH	4]
	in urban and rural areas (1990-1991).	
Figure 4 :	Average nutrients content of suggested LCAD models (1-6)	76
	as percentages of RDA for the ALISH in the urban regions.	
Figure 5 :	Average nutrients content of suggested LCAD models (7-12)	73
	as percentages of RDA for the ALISH in the rural regions.	