



Faculty of Science
Dept. of Entomology

Efficiency of venom components of Egyptian and Carniolan honeybee in Egypt at different storage conditions

A Thesis

Submitted to the Department of Entomology,
Faculty of Science, Ain Shams University
For the award of the Ph.D. Degree of Science in Entomology

By

Hind Abd El-Ghany Zidan Amer

M. Sc.

Supervisors

Prof. Dr. Zahia Kamel Mostafa

Emeritus Professor of Insect
Physiology- Entomology Department
Faculty of Science
Ain Shams University

Prof. Dr. EL-Said Ibrahim Haggag

Emeritus Chief Researcher of Honeybee
Research Department
Plant Protection Research Institute
Agricultural Research Center

Prof. Dr. Mahmoud A. Ibrahim

Emeritus Researcher Professor of
Biochemistry
Molecular Biology Department
National Research Centre

Dr. Abir Abdel-Hamid Elfiky

Director of ANDI Center of excellence
Antivenom research Vacsera, Egypt.

Dr. Doaa Abdel-Khalek Darwish

Associate Researcher Professor of Biochemistry
Molecular Biology Department
National Research Centre

Department of Entomology

**Faculty of Science
Ain Shams University
Cair – Egypt**

2018

﴿قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا﴾

إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ

سورة البقرة آية (٣٢)

Biography

Name: Hind Abd El- Ghany Zidan Amer

Degree Awarded: M.Sc. in Entmology- Faculty of Science-
Ain Shams University.

Occupation: Assistant Researcher- Honeybee Research Depart.
Plant Protection Research Institute- Dokki- A.R.C

Date of Registration for Ph.D.: 2014

Acknowledgment

First of all I do thank **ALLAH** for the unlimited help. I wish to express my deepest appreciation and respect to **Prof. Dr. Zahia Kamel Mostafa** Professor of Entomology, Ain Shams University, for suggesting the problem, the critical review of the manuscript and her kind supervision.

My sincere thanks go to **Prof. Dr. Mahmoud A. Ibrahim**, Professor of Biochemistry Molecular Biology Department National Research Centre, for his excellent ideas, energetic guidance, offering all facilities to me and close supervision.

I am also particularly indebted to **Prof. Dr. EL-Said Ibrahim Haggag**, Professor of Honeybee Research Department Plant Protection Research Institute Agricultural Research Center for generous assistance.

I am extremely grateful to **Dr. Doaa Abdel-Khalek Darwish**, Associate Professor of Biochemistry Molecular Biology Department National Research Centre for her advice and guidance from very early stage of this work as well as providing me extraordinary experiences throughout the work.

Deep and sincere gratitude to **Dr. Abir Abdel-Hamid Elfiky** Director of ANDI Center of excellence Antivenom research Vacsera. For her encouragement and advises as a member of my supervisory committee.

Finally I am grateful to the head and staff members of the Entomology Department, Faculty of Science, Ain Shams and my friends who have helped me, contributed with me on way during the advancement of the different stages of this work.

CONTENTS		Page
I.	INTRODUCTION	1
II.	LETERATURE REVIEW	5
III.	MATERIAL AND METHODS	18
	Experimental Insects	18
	Venom Collection	18
	1. Manual extracted method	18
	2. Electrical stimulated method	19
	i. Spectrophotometer studies	20
	- Protein determination	20
	1. Hyaluronidase assay	21
	2. Phospholipase A2 (PLA2) assay	22
	3. Haemolytic activity assay of melittin	23
	4. Protease assay	24
	ii. Electrophoresis studies	25
	1. Native polyacrylamide gel electrophoresis (Native-PAGE)	25
	2. Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE)	28
	3. Isoelectrofocusing (IEF) on polyacrylamide gel	33
	4. Hyaluronidase activity on SDS– PAGE.	37
	iii. Toxicity studies	38
	- Determination of LD50	38
IV.	RESULTS	40
	i. Spectrophotometer studies	40
	■ Venom manually and electrically extracted before storage	40
	1. Hyaluronidase activity	40
	2. PLA2 activity	41
	3. Haemolytic activity of melittin	41

	4. Protease activity	42
	■ Venoms electrically extracted during storage for 24 months	45
	1. Hyaluronidase activity	45
	2. PLA2 activity	50
	3. Haemolytic activity of melittin	55
	ii. Electrophoresis studies	61
	■ Venom manually and electrically extracted before storage	61
	1. Native- PAGE	61
	2. SDS- PAGE	65
	3. Isoelectrofocusing on polyacrylamide gel	70
	4. Hyaluronidase activity on SDS– PAGE	75
	■ Venoms electrically extracted after storage for 24 months	77
	1. Native- PAGE	77
	- Egyptian honeybee subspecies	77
	- Carniolan honeybee subspecies	77
	2. SDS- PAGE	81
	- Egyptian honeybee subspecies	81
	- Carniolan honeybee subspecies	81
	3. Isoelectrofocusing on polyacrylamide gel	85
	- Egyptian honeybee subspecies	85
	- Carniolan honeybee subspecies	85
	4. Hyaluronidase activity on SDS– PAGE	89
	iii. Toxicity studies	91
	■ Venoms manually and electrically extracted before storage	91
	■ Venoms electrically extracted after storage for 12 and 24 months	93

V.	DISCUSSION	99
	i. Spectrophotometer studies	99
	■ Venoms manually and electrically extracted before storage	99
	■ Venoms electrically extracted during storage for 24 months	99
	ii. Electrophoresis studied	102
	■ Venoms manually and electrically extracted before storage	102
	■ Venoms electrically extracted after storage for 24 months	109
	■ Hyaluronidase activity on SDS – PAGE	110
	- Venoms manually and electrically extracted before storage	110
	- Venoms electrically extracted after storage for 24 months	110
	iii. Toxicity studies	111
	■ Venoms manually and electrically extracted before storage	111
	■ Venoms electrically extracted after storage for 12 and 24 months	113
VI.	SUMMARY	114
	REFERENCES	120
	ARABIC SUMMARY	1

List of Tables

No.	Title	Page
1	Preparation of SDS-PAGE.....	32
2	Preparation of the gel for isoelectrofocusing (IEF)	36
3	The hyaluronidase, PLA2, melittin and protease activities of the two honeybee subspecies venoms extracted using manual (A) and electric (B) methods.....	43
4	The mean values of hyaluronidase activity (U/mg) \pm S.E. for the two subspecies of the foraging workers venom extracted by electrical stimulated method during storage for 24 months at the different conditions.....	47
5	The mean values of PLA2 activity (U/mg) \pm S.E. for the two subspecies of the foraging workers venom extracted by electrical stimulated method during storage for 24 months at the different conditions	52
6	The mean values of lysis (%) of melittin \pm S.E. for the two subspecies of the foraging workers venom extracted by electrical stimulated method during storage for 24 months at the different conditions	57
7	The relative mobility (Rm) and its corresponding relative concentration (%) values of the non-denatured protein bands of the foraging workers venom extracted using two different methods for two <i>Apis mellifer</i> subspecies.....	64
8	Estimated similarity index (S.I) for the non-denatured proteins of the foraging workers venom extracted using two different methods between the two <i>Apis mellifera</i> subspecies.	64
9	The molecular weight (MW) and its corresponding relative concentration (%) values of the denatured protein bands of the foraging workers venom extracted using two different methods for the two <i>Apis mellifera</i> subspecies	68
10	Estimated similarity index (S.I) for the denatured proteins of the foraging workers venom extracted using two different methods between two <i>Apis mellifera</i> subspecies	69
11	The Isoelectric point (pI) and its corresponding relative concentration (%) values of the IEF protein bands of the foraging workers venom extracted using two different methods for two <i>Apis mellifera</i> subspecies.....	73
12	Estimated similarity index (S.I) for the IEF proteins of the foraging workers venom extracted using two different methods between two <i>Apis mellifera</i> subspecies	74

No.	Title	Page
13	The relative mobility (R _m) and its corresponding relative concentration (%) values of the non- denatured protein bands of the Egyptian and Carniolan foraging workers venom electrically stimulated after storage for 24 months at four different conditions	80
14	The molecular weight (MW) and its corresponding relative concentration (%) values of the denatured protein bands of the Egyptian and Carniolan foraging workers venom electrically stimulated after storage for 24 months at four different conditions	84
15	Isoelectric point (pI) and its corresponding relative concentration (%) values of the denatured protein bands of the Egyptian and Carniolan foraging workers venom electrically stimulated after storage for 24 months at four different conditions	88
16	The mean values of LD ₅₀ (mg/kg) ± S.E. for the foraging workers venom extracted using two different methods for two <i>Apis mellifera</i> subspecies	92
17	The mean values of LD ₅₀ (mg/kg) ± S.E. for the two subspecies of the foraging w foraging workers venom collected using electrical stimulated method during ke during storage for 24 months at four different conditions	95

List of Figers

No.	Title	Page
1	Manually extracted of venom.....	19
2	Bee venom collector device	20
3	Standard curve of the protein concentration using bovine serum albumin as a standard protein, according to the method of Bradford (1976)	21
4	Standard curve of the melittin according to the method of Owen and Pfaff (1995).....	24
5	Effects of manual (A) and electrical extracted (B) methods on hyaluronidase, PLA2, melittin and protease activities of venoms for the two honeybee subspecies	44
6	Effects of the different storage conditions on hyaluronidase activity of Egyptian and Carniolan bee venoms during the storage periods	48
7	The reduction (%) in the level of hyaluronidase activity for the two honeybee subspecies venoms in the different storage conditions during the storage periods	49
8	Effects of the different storage conditions on PLA2 activity of Egyptian and Carniolan bee venoms during the storage periods	53
9	The reduction (%) in the level of PLA2 activity for the two honeybee subspecies venoms in the different storage conditions during the storage periods	54
10	Effects of the different storage conditions on haemolytic activity of melittin of Egyptian and Carniolan bee venoms during the storage periods	58
11	The reduction (%) in the level of haemolytic activity of melittin for the two honeybee subspecies venoms in the different storage conditions during the storage periods	59
12	The reduction (%) of the hyaluronidase, PLA2 and milittin activities in the storage venoms for the two honeybee subspecies after 24 months in the different storage conditions.	60
13	Native- PAGE of the non- denatured proteins of the foraging workers venom extracted using two different methods for two <i>Apis mellifera</i> subspecies	63
14	SDS-PAGE of the denatured proteins of the foraging workers venom extracted using two different methods for two <i>Apis mellifera</i> subspecies	67