

---

---

# **THE ROLE OF CARDIAC MAGNETIC RESONANCE IMAGING IN THE DIAGNOSIS OF CARDIOMYOPATHY IN PEDIATRICS**

**Thesis**

**Submitted for partial fulfillment of**

**M.D Degree in Radiodiagnosis**

**BY**

**Sara Mahmoud Kamel Ali**

**M.B.B.Ch,M.Sc**

*Supervised by*

**Prof. Hassan Ali Hassan El Kiki**

*Professor of Radiodiagnosis*

*Faculty of Medicine*

*Cairo University*

**Prof. Zeinab Salah El Din Mohamed Selim**

*Professor of Pediatric*

*Faculty of Medicine*

*Cairo University*

**Dr. Rania Zakaria Hassan**

*Lecturer of Radiodiagnosis*

*Faculty of Medicine*

*Cairo University*

*Faculty of Medicine*

*Cairo University*

**2015**

**"قالوا سبحانك لا علم لنا إلا ما علمتنا  
إنك أنت العظيم الحكيم"**

سورة البقرة-الآية 32

# *Acknowledgement*

First and foremost, I would like to express my deepest gratitude and thankfulness to Allah for giving me the will and strength to fulfill this work and then for my family for their continuous encouragement and great helps.

My sincere gratitude and unlimited thanks are addressed to my mentor Dr. Hassan Elkiki, professor of radiodiagnosis, Cairo University, for his continuous advice, encouragement and sincere remarks. It has been an extreme pleasure for me to proceed with this work under his supervision.

I would also like to thank Dr.Zeinab Salah Eldin for her support, encouragement, cooperation and input what made this research possible.

I am also grateful to Dr.Rania Zakaria, lecturer of radiodiagnosis, Cairo University. I am extremely thankful and indebted to her for sharing expertise, and sincere and valuable guidance and encouragement extended to me.

Last but not least, my deep gratitude and thanks to all my dear professors and colleagues of the radiology department for their help and cooperation.

# *Dedication*

To all my family members for  
their inspiration, indispensable  
support and continuous  
encouragement

# TABLE OF CONTENTS

List of figures.....	i
List of tables.....	x
List of abbreviations.....	xi
Abstract.....	xvi
Introduction.....	xvii
Aim of the work.....	xix
<b>Review of literature.....</b>	<b>1</b>
Diagnosis of cardiomyopathy.....	2
Technique of cardiac MRI .....	16
Cardiac MRI.....	36
Cardiac MRI in cardiomyopathy.....	69
<b>Patients and methods.....</b>	<b>100</b>
<b>Results.....</b>	<b>107</b>
<b>Case presentation.....</b>	<b>119</b>
<b>Discussion.....</b>	<b>146</b>
<b>Summary and conclusion.....</b>	<b>152</b>
<b>Recommendations.....</b>	<b>154</b>
<b>References.....</b>	<b>155</b>

الملخص العربي

## LIST OF FIGURES

No.		Page
<b>Figure 1</b>	Four-chamber bidimensional echocardiographic view of a 16 year-old asymptomatic male patient with hypertrophic cardiomyopathy with asymmetric septal hypertrophy. The septum and the posterior wall measure 21 mm and 9.5 mm respectively ( <b>Alday and Moreyra; 2012</b> ).	<b>9</b>
<b>Figure 2</b>	CMR set-up for paediatric general anaesthetic cases. View of the MR scanner room showing the anesthetic machine (A) and monitoring equipment (B). Ventilation tubing and leads from both pieces of equipment pass through a small opening in the wall (C) into the control room, so that the anesthetist can control breath-holding and monitor the patient from within the control room ( <b>Ntsinjana et al; 2011</b> ).	<b>18</b>
<b>Figure 3</b>	A 1.5T magnet used in cardiac imaging ( <b>Norton et al; 2013</b> ).	<b>19</b>
<b>Figure 4</b>	A typical RF coil used in cardiac MRI. This is placed on the patient's chest during the study ( <b>Norton et al;2013</b> ).	<b>19</b>
<b>Figure 5</b>	Representative black blood images of the heart in the short-axis (A), vertical long axis (B), 4-chamber (C) and left ventricular outflow tract (D) views are shown in this figure. Notice effective nulling of blood independent of the orientation of the acquisition ( <b>Krishnamurthy et al; 2014</b> ).	<b>22</b>
<b>Figure 6</b>	Example of Cardiac Magnetic Resonance Tagging .The tags are seen as dark lines. The deformation of the tag lines can be seen at end-systole ( <b>Petersen et al; 2011</b> ).	<b>24</b>
<b>Figure 7</b>	Phase-contrast images of a normal subject. A and B, Transverse aortic (A) and pulmonary artery (B) phase-contrast images ( <b>Ginat et al; 2011</b> ).	<b>25</b>
<b>Figure 8</b>	Gadolinium enhancement patterns in ischemic versus nonischemic cardiomyopathy. Top row indicates carton of enhancement pattern, and bottom row shows patient examples from short-axis delayed enhancement MRI. A, Subendocardial enhancement or transmural enhancement is associated with infarction from ischemic disease.	<b>28</b>

	<b>B</b> , Nonischemic cardiomyopathy with midwall stripe pattern. This pattern is common in idiopathic dilated cardiomyopathy. <b>C</b> , Nonischemic cardiomyopathy with epicardial enhancement pattern .This patient had viral myocarditis. ( <b>Hunold et al; 2005</b> ).	
<b>Figure 9</b>	Cardiac amyloidosis (a, b) Delayed contrast-enhanced short axis cardiac MR images show diffuse subendocardial enhancement (arrows) that is not limited to a vascular distribution, along with the absence of blood pool enhancement, findings that are characteristic of cardiac amyloidosis ( <b>Cummings et al; 2009</b> ).	<b>30</b>
<b>Figure 10</b>	Histiocytoid cardiomyopathy in a 7-month-old girl who presented with recurrent ventricular tachyarrhythmias. Delayed contrast-enhanced short-axis single-shot true FISP images obtained from the base ( <b>a</b> ) toward the apex ( <b>c</b> ) show multiple enhancing nodules (arrow) involving the subendocardium of the septal, inferior, and lateral walls. Transcatheter myocardial biopsy revealed histiocytoid myocytes consistent with histiocytoid cardiomyopathy.The patient subsequently underwent cardiac transplantation ( <b>Cummings et al; 2009</b> ).	<b>32</b>
<b>Figure 11</b>	Representative Segmentation and Pattern of Contrast HE over 17-Segments of the LAD, RCA, and LCX Arteries .A single basal, mid, apical, and 2-chamber contrast-enhanced views of 3 subjects (left panels) following an acute myocardial infarction with occlusion to the left anterior descending (LAD) artery, right coronary artery (RCA), and left circumflex (LCX) artery. The dotted yellow lines delineate and separate the 17-segments over the entire left ventricle. Bulls-eye diagrams (right) reveal the pattern of infarction for each example. CMR cardiac magnetic resonance; HE hyperenhancement ( <b>Ortiz-Pérez et al;2008</b> ).	<b>34</b>
<b>Figure 12</b>	Chart illustrates the differential diagnosis of delayed contrast enhancement at cardiac MR imaging by location ( <b>Cummings et al; 2009</b> ).	<b>35</b>
<b>Figure 13</b>	Schematic shows orientation of major body planes with respect to patient and their corresponding appearance on bright blood imaging sequences ( <b>Ginat et al;2011</b> ).	<b>37</b>
<b>Figure 14</b>	Schematic shows orientation of major cardiac planes with respect to heart and their corresponding appearance on bright blood sequences ( <b>Ginat et al; 2011</b> ).	<b>38</b>
<b>Figure 15</b>	Normal cardiac MRI anatomy shown in healthy subject. A–E, Short-axis (A), horizontal long-axis (B), two-chamber (C), right	<b>40</b>

	ventricular outflow tract (D), and left ventricular outflow tract (E) views ( <b>Ginat et al; 2011</b> ).	
<b>Figure 16</b>	Normal cardiac MRI anatomy shown in healthy subject. Axial fast imaging employing steady-state acquisition image shows prominent crista terminalis ( <i>arrow</i> ).This finding is normal variant not to be confused with tumor or thrombus ( <b>Ginat et al; 2011</b> ).	<b>43</b>
<b>Figure 17</b>	Transverse images in normal cardiac MRI ( <b>Bogaert et al; 2005</b> ).	<b>44</b>
<b>Figure 18</b>	Coronal images in normal cardiac MRI. Lm left main stem coronary artery.mv mitral valve.sv sinus of valsalva ;Other abbreviations as in the previous figures ( <b>Bogaert et al;2005</b> ).	<b>45</b>
<b>Figure 19</b>	Sagittal images in normal cardiac MRI .lad left anterior descending coronary artery;lcx left circumflex artery ;pavg posterior (or left) atrioventricular groove; puv pulmonary valve; sv sinus of valsalva;other abbreviations as in <b>Figure 16</b> ( <b>Bogaert et al;2005</b> ).	<b>46</b>
<b>Figure 20</b>	Short-axis images in normal cardiac MRI ,aivg anterior interventricular groove ;d diaphragm; lvap LVapex; pivg posterior interventricular groove ;other abbreviations as in preceeding figures ( <b>Bogaert et al;2005</b> ).	<b>47</b>
<b>Figure 21</b>	Horizontal long –axis images in normal cardiac MRI.fo fossa ovalis; pda posterior descending artery. Other abbreviations as in preceeding figures ( <b>Bogaert et al;2005</b> ).	<b>48</b>
<b>Figure 22</b>	Vertical long-axis images in normal cardiac MRI .Abbreviations as in preceeding figures ( <b>Bogaert et al;2005</b> ).	<b>49</b>
<b>Figure 23</b>	Left ventricular outflow tract images. Abbreviations as in preceeding figures ( <b>Bogaert et al;2005</b> ).	<b>50</b>
<b>Figure 24</b>	Right ventricular outflow tract image. Abbreviations as preceeding figures ( <b>Bogaert et al;2005</b> ).	<b>50</b>
<b>Figure 25</b>	Display, on a circumferential polar plot, of the 17 myocardial segments and the recommended nomenclature for tomographic imaging of the heart ( <b>Cerqueira et al; 2002</b> ).	<b>51</b>
<b>Figure 26</b>	Assignment of the 17 myocardial segments to the territories of the left anterior descending (LAD), right coronary artery (RCA), and the left circumflex coronary artery (LCX) ( <b>Cerqueira et al; 2002</b> ).	<b>52</b>
<b>Figure 27</b>	MR tissue tagging in short-axis orientation for strain analysis of the myocardial motion ( <b>Westenberg; 2011</b> ).	<b>55</b>
<b>Figure 28</b>	LV time-volume relation (c) determined by planimetry from a	<b>58</b>



	multislice cine short-axis dataset (b), planned perpendicular to the long-axis of the LV (a). From the time-volume relation, early peak filling rate (EPFR) is determined from the steepest gradient in the volume curve in the early filling phase. Atrial filling fraction (AFF) is determined in the atrial filling phase ( <b>Westenberg; 2011</b> ).	
<b>Figure 29</b>	Left atrial size is determined by biplane Simpson's rule on the atrial areas determined in four-chamber (a) and two-chamber (b) view ( <b>Westenberg; 2011</b> ).	<b>59</b>
<b>Figure 30</b>	At the moment of end-systole, an acquisition plane is positioned at the mitral valve (a). Through-plane one-directional velocity-encoded MRI (magnitude and velocity image in b) results in a time-flow rate graph (c), which is used for wave form analysis. Early (E) and atrial (A) peak filling rate can be determined, as well as the deceleration time (DT) of the E-peak ( <b>Westenberg; 2011</b> ).	<b>60</b>
<b>Figure 31</b>	Three-dimensional three-directional velocity-encoded MRI at the aorta and mitral valve (a) results in velocity images reconstructed at the ascending aorta (b) and the mitral valve (d). From the velocity images, time-flow rate graphs (c and e) are obtained from which the isovolumic relaxation time (IVRT) can be determined ( <b>Westenberg; 2011</b> ).	<b>61</b>
<b>Figure 32</b>	In-plane one-directional velocity-encoded MRI in four chamber orientation (magnitude image in (a) and velocity image in (b) Velocity sampling at the pulmonary vein (arrow) results in a time velocity graph (c), from which peak systolic velocity (S), peak anterograde diastolic velocity (D), and peak atrial reversal velocity (Ar) can be determined ( <b>Westenberg; 2011</b> ).	<b>62</b>
<b>Figure 33</b>	The time-resolved velocity vector field (presented with color coding) on a four-chamber view is obtained from three-dimensional three-directional velocity-encoded MRI. From this inflow velocity vector field, the flow propagation velocity can be determined ( <b>Westenberg; 2011</b> ).	<b>64</b>
<b>Figure 34</b>	<b>Left ventricular volumes and mass.</b> Lines indicate median and 95% prediction bands ( <b>Buechel et al;2009</b> ).	<b>65</b>
<b>Figure 35</b>	<b>Right ventricular volumes and mass.</b> Lines indicate median and 95% prediction bands ( <b>Buechel et al;2009</b> ).	<b>67</b>
<b>Figure 36</b>	Cine balanced steady-state free precession (SSFP) images are routinely used for ventricular volumetry, due to their high blood-	<b>71</b>

	to-myocardial contrast, and SNR. For LV and RV volume quantification, a stack of short axis slices (~8 mm slice thickness) covering both the ventricles from the base of the heart to the apex is obtained (A); (B) shows representative images from the stack at three different locations, and at systole and diastole. Contours demarcating the LV and RV cavity are drawn at end-systole and end-diastole for all the slices. The areas are summed to obtain the end-diastolic volume (EDV), end-systolic volume (ESV). From EDV, and ESV, the stroke volume ( $SV = EDV - ESV$ ), and ejection fraction ( $EF = SV/EDV \times 100$ ) are calculated ( <b>Krishnamurthy et al;2014</b> ).	
<b>Figure 37</b>	Contour tracing in the endsystolic phase ( <b>Buechel et al; 2009</b> ).	<b>71</b>
<b>Figure 38</b>	A. Long axis view of a cardiac magnetic resonance image of a 5 year-old asymptomatic boy with severe hypertrophic cardiomyopathy. <b>B:</b> Magnetic resonance imaging of a short axis projection of the heart of a 3 year-old boy with severe heart failure. (IVS: interventricular septum, LV: left ventricle, PW: posterior wall, RV: right ventricle) ( <b>Alday and Moreyra; 2012</b> ).	<b>78</b>
<b>Figure 39</b>	<b>A</b> .Cardiac magnetic resonance image of the heart of a patient with positive delayed gadolinium enhancement indicating a diffuse pattern of fibrosis (arrowheads). <b>B:</b> Long axis view of a cardiac magnetic resonance image of the heart of a 9 year-old boy with a localized delayed gadolinium enhancement image in the interventricular septum (arrow) ( <b>Alday and Moreyra; 2012</b> ).	<b>79</b>
<b>Figure 40</b>	<b><u>Differential diagnosis of LVH.</u></b> Diastolic SSFP cines (above) and LGE-sequences (below) in the 4-chamber view of several types of hypertrophy are depicted. From left to right: hypertensive cardiomyopathy, hypertrophic cardiomyopathy, Fabry's disease, cardiac amyloid infiltration and cardiac sarcoidosis. Different gadolinium myocardial enhancement patterns (arrows) are shown corresponding to each condition: very subtle focal intramyocardial fibrosis in the basal septum of a hypertensive cardiomyopathy patient, confluent intramyocardial fibrosis in a case of hypertrophic cardiomyopathy, subendocardial fibrosis in the lateral wall in a case of Fabry's disease, global diffuse subendocardial fibrosis in amyloid cardiomyopathy and subendocardial/intramyocardial/subepicardial fibrosis in a case of sarcoidosis ( <b>Maceira and Mohiaddin ; 2012</b> ).	<b>80</b>
<b>Figure 41</b>	ARVD. Right-ventricular horizontal long-axis DE-MRI	<b>83</b>

	demonstrates diffuse, subendocardial enhancement of thinned RV wall, consistent with fibrous replacement (arrows) ( <b>Rajiah, and Flamm; 2009</b> ).	
<b>Figure 42</b>	Coronal bright blood cine (to the left) and axial bright blood cine (to the right) images from cardiac MRI showing a markedly dilated left ventricle and left atrium. The left ventricle shows prominent trabeculation particularly along the apex. A small pericardial effusion and a small pleural effusion are also noted ( <b>Fox et al; 2006</b> ).	<b>84</b>
<b>Figure 43</b>	Nonischemic dilated cardiomyopathy. Four-chamber DE-MRI demonstrates a dilated LV with linear mid myocardial scarring in the basal septum (arrows). The coronary arteries were normal in this patient ( <b>Rajiah, and Flamm; 2009</b> ).	<b>87</b>
<b>Figure 44</b>	Assessment of increased ventricular coupling using real-time cine CMR in constrictive pericarditis. Short-axis cine CMR at onset of inspiration (a) and onset of expiration (b). Images are acquired at early ventricular filling. Septal inversion occurs at inspiration (a), with increased right-sided septal motion at expiration (b), leading to an abnormal respiratory septal shift. The horizontal dashed lines indicate the position of the left hemidiaphragm ( <b>Bogaert and Francone; 2009</b> )	<b>90</b>
<b>Figure 45</b>	Acute myocarditis. Three-chamber long-axis DE-MRI demonstrates patchy, mid myocardial-to-epicardial enhancement in the basal and mid inferolateral and inferior walls (arrows) ( <b>Rajiah, and Flamm; 2009</b> ).	<b>93</b>
<b>Figure 46</b>	Examples of cardiomyopathies. A. 4-chamber, balanced-SSFP view in hypertrophic cardiomyopathy. Note the marked thickening of the septum with compression of the RV cavity. B. 4-chamber, balanced-SSFP view in left ventricular non-compaction. Note the arrowheads show areas of thin compacted myocardium. C. 4-chamber, late gadolinium enhancement (LGE) image in idiopathic dilated cardiomyopathy. Note no LGE. D. Short-axis, LGE image in a patient with critical aortic stenosis, restrictive cardiomyopathy secondary to global, sub-endocardial fibrosis ( <b>Ntsinjana et al;2011</b> ).	<b>94</b>
<b>Figure 47</b>	Cardiac sarcoidosis. Mid ventricular short-axis DE-MRI demonstrates patchy, mid myocardial scarring in the inferoseptum (arrow) in a patient with known systemic	<b>97</b>

	sarcoidosis ( <b>Rajiah, and Flamm; 2009</b> ).	
<b>Figure 48</b>	This poster shows easy and efficient planning of the main cardiac views using the Philips Intera Cardiac Package( <b>www.Philips.com</b> ).	<b>104</b>
<b>Figure 49</b>	Sex distribution of the study sample.	<b>107</b>
<b>Figure 50</b>	Pie chart representing the types of cardiomyopathy in the study sample.	<b>109</b>
<b>Figure 51</b>	Correlation between LV EDV and BSA among the study sample, LVEDV: left ventricular end diastolic volume, BSA: body surface area.	<b>115</b>
<b>Figure 52</b>	Correlation between LV ESV and BSA among the study sample, LVESV: left ventricular end systolic volume, BSA: body surface area.	<b>116</b>
<b>Figure 53</b>	Correlation between RV EDV and BSA among the study sample, RVEDV: right ventricular end diastolic volume, BSA: body surface area.	<b>117</b>
<b>Figure 54</b>	Correlation between RV ESV and BSA among the study sample, RVESV: right ventricular end systolic volume, BSA: body surface area.	<b>118</b>
<b>Figure 55</b>	Four chamber SSFP view showing markedly dilated right atrium (red circle to the left), and markedly dilated left atrium (yellow circle to the right) with moderate mitral regurge (MR).	<b>119</b>
<b>Figure 56</b>	Four chamber view SSFP showing average size of both ventricle.	<b>120</b>
<b>Figure 57</b>	Inversion Recovery (IR) Short axis view showing subendocardial enhancement at the inferoseptal wall of the left ventricle (red arrow).	<b>120</b>
<b>Figure 58</b>	Cardiac MRI four chamber view showing enlarged both atria with normal size of both ventricles, (RA) right atrium, (LA) left atrium.	<b>123</b>
<b>Figure 59</b>	Cardiac MRI (to the left) four chamber view sowing the congested hepatic veins (arrow head), (to the right) IR (inversion recovery) delayed post contrast short axis view, 10 minutes after contrast administration showing no enhancement.	<b>124</b>
<b>Figure 60</b>	Automated wall motion analysis of the left ventricle using the Philips work station, showing global hypokinesia with patchy supero-septal areas of marked hypokinesia.	<b>124</b>

<b>Figure 61</b>	Cardiac MRI four chamber view showing markedly dilated both atria with normal size of both ventricles. Moderate pericardial effusion is also noted (arrow head). (RA) right atrium, (LA) left atrium.	<b>127</b>
<b>Figure 62</b>	Short axis view during early diastole (to the right) and late diastole (to the left) showing the flattening of the interventricular septum during early diastole (arrow).	<b>128</b>
<b>Figure 63</b>	Axial T2-TSE cardiac MRI showing the pericardial thickness of 2.3mm. Mean normal pericardial thickness is $1.2 \pm 0.8$ mm and does not exceed 2.5 mm. Pericardial thickness $\geq 3$ mm is considered thickened.	<b>128</b>
<b>Figure 64</b>	Automated wall motion analysis of the left ventricle using the Philips work station, showing dyskinesia at the septal wall of the left ventricle (interventricular septum) (blue, arrowed at the upper right image) with mild hypokinesia of the rest of the left ventricular walls.	<b>129</b>
<b>Figure 65</b>	Four chamber view SSFP during systole (to the left) and diastole (to the right) showing the dilated left ventricle.	<b>132</b>
<b>Figure 66</b>	Four chamber view showing normal origin of the right coronary artery (yellow arrow) and the left coronary artery (red arrow) from the aorta.	<b>133</b>
<b>Figure 67</b>	IR (inversion recovery) delayed post contrast short axis view, 10 minutes after contrast administration showing no enhancement.	<b>133</b>
<b>Figure 68</b>	Automated wall motion analysis of the left ventricle using the Philips work station, showing global left ventricular hypokinesia.	<b>134</b>
<b>Figure 69</b>	Four chamber view SSFP showing cardiomegaly, with left ventricular enlargement.	<b>136</b>
<b>Figure 70</b>	Four chamber view SSFP showing normal origin of the left coronary artery (yellow arrow) and right coronary artery from the aorta (red arrow).	<b>137</b>
<b>Figure 71</b>	Automated wall motion analysis of the left ventricle using the Philips work station, showing global hypokinesia with patchy supero-septal areas of marked hypokinesia.	<b>137</b>
<b>Figure 72</b>	IR (inversion recovery) delayed post contrast short axis view, 10 minutes after contrast administration showing no enhancement.	<b>138</b>
<b>Figure 73</b>	Four chamber view showing the ratio of the non compacted to compacted myocardium at the lateral wall of the left ventricle	<b>140</b>

	(arrow head) <b>2.5</b> and at the apex (red arrow) <b>5.6</b> (Both are more than 2.3)	
<b>Figure 74</b>	Four chamber view showing the deep intertrabecular recesses of the non compacted layer.	<b>141</b>
<b>Figure 75</b>	Four chamber view during systole (to the left) and during diastole (to the right) showing almost complete obliteration of the left ventricular cavity during the systole with marked asymmetrical myocardial thickening and mild mitral regurge (arrow to the left).	<b>143</b>
<b>Figure 76</b>	Vertical long axis view during systole (to the left) and during diastole (to the right) showing the same previous findings.	<b>144</b>
<b>Figure 77</b>	Inversion Recovery (IR) Short axis view showing patchy areas of myocardial enhancement not confined to a known vascular distribution (arrows).	<b>144</b>

## LIST OF TABLES

No.		Page
<b>Table 1</b>	European Society of Cardiology classification of primary cardiomyopathies ( <b>modified from Elliot, et al., 2008</b> ).	<b>3</b>
<b>Table 2</b>	World Health Organization classification of cardiomyopathies ( <b>Maron et al;2006</b> ).	<b>4</b>
<b>Table 3</b>	AHA classification model for primary cardiomyopathies ( <b>Maron et al; 2006</b> ).	<b>5</b>
<b>Table 4</b>	Optimal Planes for Imaging Cardiac Structures and Chambers ( <b>Ginat et al;2011</b> ).	<b>39</b>
<b>Table 5</b>	Classification of diastolic dysfunction ( <b>Westenberg; 2011</b> ).	<b>57</b>
<b>Table 6</b>	Sex distribution of the study population	<b>107</b>
<b>Table 7</b>	Demographic data of the study population	<b>108</b>
<b>Table 8</b>	Types of cardiomyopathy in the study population	<b>109</b>
<b>Table 9</b>	Cardiac MRI findings within the study population	<b>110</b>
<b>Table 10</b>	Relative atrial volume ratio (RAR) in cases referred to have constrictive pericarditis versus restrictive cardiomyopathy.	<b>111</b>
<b>Table 11</b>	Left ventricular volumes measured by MRI in the study population.	<b>112</b>
<b>Table 12</b>	Right ventricular volumes measured by MRI in the study population.	<b>113</b>
<b>Table 13</b>	CMR findings in Dilated cardiomyopathy patients.	<b>114</b>