سامية محمد مصطفى



شبكة المعلومات الحامعية

بسم الله الرحمن الرحيم



-Caro-

سامية محمد مصطفي



شبكة العلومات الحامعية



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





سامية محمد مصطفى

شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسو

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة يعيدا عن الغيار



سامية محمد مصطفي



شبكة المعلومات الجامعية



المسلمة عين شعور المسلمة عين شعور المسلمة عين شعور المسلمة عين شعور المسلمة ا

سامية محمد مصطفى

شبكة المعلومات الحامعية



بالرسالة صفحات لم ترد بالأصل



POSTURAL CONTROL TRAINING: EFFECT ON BALANCE AND LOCOMOTION IN HEMIPARETIC CHILDREN

A Thesis

Submitted to Physical Therapy Department for Growth and Developmental Disorders in Children and its Surgery in the Partial Fulfillment of the Requirements for the Doctoral Degree in Physical Therapy

By

Heba Mohamed Youssr Mohamed El Basatiny
M.Sc. Physical Therapy

Faculty of Physical Therapy
Cairo University
2004

5

VC V-0

A. ElTohan Supervisors

Prof. Amira Mohamed El-Tohamy

Professor of Growth and Developmental
Disorders in Children and its Surgery
Faculty of Physical Therapy
Cairo University

Prof. Elham EL-Sayed Salem
Professor of Growth and Developmental
Disorders in Children and its Surgery
Faculty of Physical Therapy
Cairo University



Acknowledgements

First of all, I would like to kneel thanking "GOD" for giving me the ability and patience to accomplish this work.

I would like to express my deepest appreciation and respect to *Prof. Amira M. El-Tohamy*, Prof. of Growth and Developmental Disorders in Children and its Surgery, Faculty of Physical Therapy, Cairo University. She suggested this subject, supervised all details of the work, and enriched me with her unlimited experience.

Her continuous guidance and assistance were the key stone in preparing this work; to her I am greatly indebted in heart.

deep would like to express my gratitude Prof. Elham El-Sayed Salem, Prof. of Growth and Developmental Disorders in Children and its Surgery, Faculty of Cairo Physical Therapy. University. Her continuous cooperation, supervision, guidance & kind help in reading, correction, and revision of this work.

Also, I would like to express my deepest thanks to my professors, lecturers & colleagues in the physical therapy department for Growth & developmental disorders in children & its surgery, Faculty of Physical Therapy, Cairo University, for their valuable guidance & consultance.

Heba M. Youssr El Basatiny
2004

Dedication

To

My Husband, Parents and Children
for their Continuous Support and
Encouragement

Abstract

Postural control training: Effect on balance & locomotion in hemiparetic children. By: Heba Mohamed Youssr Mohamed El Basatiny; Supervisors: Prof. Dr. Amira Mohamed El-Tohamy, Prof. Dr. Elham El-Sayed Salem. Doctoral thesis P.T., Physical Therapy Department for Growth and Developmental Disorders in Children and its Surgery, Faculty of Physical Therapy, Cairo University, 2004.

Abstract

The purpose of this work was to study the effect of postural control training on balance & locomotion in spastic hemiparetic children. The study was conducted on forty-two children, ranging in age from 7 to 12 years from both sexes, they include: fifteen normal children served as a control group (A) & twenty-seven ambulant spastic hemiparetic children as study group (B). They received a balance training program on Biodex system in addition to a physical therapy program 3 times/week for 3 months. The post treatment stability indices & gait parameters results showed significant improvement in balance & locomotion of study group (B) after treatment which confirm the importance of postural control training (by using Biodex system) in the rehabilitation program of spastic hemiparetic children.

Key Words: Postural Control, balance, locomotion, hemiplegia.

List of Abbreviations

AP : Anteroposterior.

BOS: Base of support.

cm : Centimeters.

CNS : Central nervous system.

COG: Center of gravity.

COM: Center of mass.

CP: Cerebral palsy.

IQ: Intellectual queint.

kg: Kilograms.

m: Meter.

m/s : Meter/second.

Max: Maximum.

Min : Minimum.

ML: Mediolateral.

NDT: Neurodevelopmental treatment.

No.: Number.

P : Probability value.

S.D: Standard deviation.

sec : Second.

SEM: Standard error of mean.

SI : Stability index.

t : Paired samples t-test.

VOR: Vestibuloocular reflex.

VSR : Vestibulospinal reflex.

 \overline{X} : Mean.

% : Percentage.

(*) : Significant.

List of Tables

<u>Table</u>	:	<u>Page</u>
(1)	Age, height & weight of Normal group (A) (n = 15) &	
	Study group (B) (n = 27)	.69
(2)	Side of lesion, age, and spasticity grades distribution in the	
	study sample (group B)	.69
(3)	Pre-study overall stability index of normal children (group A)	
	& hemiparetic children (group B) at stability level 8	.71
(4)	Pre-study anteroposterior stability index of normal children	
	(group A) & hemiparetic children (group B) at stability	
	level 8	.72
(5)	Pre-study mediolateral stability index of normal children	
	(group A) & hemiparetic children (group B) at stability	
	level 8	73
(6)	The overall stability index of hemiparetic children (study grown)	up B)
	(n = 27) before & after treatment at stability level 8	76
(7)	The anteroposterior stability index of hemiparetic children	
	(study group B) $(n = 27)$ before & after treatment at stability	
	level 8	77
(8)	The mediolateral stability index of hemiparetic children	
	(study group B) $(n = 27)$ before & after treatment at stability	
	level 8	78
(9)	The overall stability index of normal children (group A) (n=15))
	in relation to post treatment values of hemiparetic children	
	(group B) (n= 27) at stability level 8	83
(10)	The anteroposterior stability index of normal children (group A))
	(n=15) in relation to post treatment values of hemiparetic	
	children (group B) (n=27) at stability level 8	84

Table	<u>e</u>	age
(11)	The mediolateral stability index of normal children (group A)	
	(n=15) in relation to post treatment values of hemiparetic	
	children (group B) (n= 27) at stability level 8	85
(12)	Stride length (m) of hemiparetic children (study group B)	
	(n = 27) before and after treatment	88
(13)	Cadence (steps/min) of hemiparetic children (study group B)	•
	(n = 27) before and after treatment	89
(14)	Speed (m/sec) of hemiparetic children (study group B)	
	(n = 27) before and after treatment	90
(15)	Cycle time (sec) of hemiparetic children (study group B)	
	(n = 27) before and after treatment	91
(16)	The percentage of stance & swing phases of the affected	
	lower limb of hemiparetic children (n = 27) before & after	
	treatment	95
	•	
		•
		•

List of Figures

<u>Figur</u>	<u>e</u> <u>Page</u>
(1)	The systems model of postural control illustrates the constant
	cycle that occurs simultaneously at many levels10
(2)	Automatic postural strategies. (A) Ankle strategy. (B) Hip
	strategy. (C) Suspensory strategy. (D) Stepping strategy16
(3)	Open-loop motor control system35
(4)	Elements of an open-loop control system
(5)	Closed-loop motor control system
(6)	A closed-loop control system
(7)	Biodex balance System; (a) display control panel,
	(b) foot platform, (c) safety support bar & (d) Printer43
(8)	Foot platform, the alphabetic letters (on sides), numbers
	(on the lowermost), and the angles (on the uppermost part
	of the platform surface)44
(9)	The Display Panel Keys49
(10)	The Balance System Main Menu49
(11)	The Test Parameter Setup Screen
(12)	Stance Type Screen for Centering Process
(13)	The Stability Test Screen Prior to Starting the Test51
(14)	The Stability Test Screen During Testing51
(15)	Camera system with six proreflex infrared cameras53
(16)	A wand-kit used for caliberation of the proreflex system53
(17)	A PC Computer with the Q Trac Software Installed54
(18)	Reflective markers attached on the subject bilaterally on the
	chosen landmarks59
(19)	Starting test position of child on the balance system61
(20)	Medical balls, roll, wedges, tilting board64
(21)	Mean values of the stability indices for normal children
	(n = 15) & hemiparetic children (Study group B) $(n = 27)$
	before treatment at stability level 874