

BIOLOGY AND CONTROL OF BROWN ROT FUNGI ON SOME STONE FRUITS

By

THAURIA MABRUOK MOHAMED ABO EL WAFa

B.Sc. Agric. Sci. (Plant Pathology), Zagazig University, 1993

M.Sc. Agric. Sci. (Plant Pathology), Ain Shams University, 2012

**A Thesis Submitted in Partial Fulfillment
Of
The Requirements for the Degree of**

**DOCTOR OF PHILOSOPHY
in
Agricultural Sciences
(Plant Pathology)**

**Department of Plant Pathology
Faculty of Agriculture
Ain Sham University**

2018

Approval Sheet

**BIOLOGY AND CONTROL OF BROWN ROT
FUNGI ON SOME STONE FRUITS**

By

THAURIA MABRUOK MOHAMED ABO EL Wafa

B.Sc. Agric. Sci. (Plant Pathology), Zagazig University, 1993

M.Sc. Agric. Sci. (Plant Pathology), Ain Shams University, 2012

This thesis for PhD. degree has been approved by:

Dr. Mohamed Mohamed Baiomy Ammar -----

Prof. Emeritus of Plant Pathology, Faculty of Agriculture, Menofia
University.

Dr. Soad Mohamed Abd -Allah -----

Prof. Emeritus of Plant Pathology, Faculty of Agriculture, Ain Shams
University.

Dr. Medhat Kamel Ali I-Sayed -----

Prof. Emeritus of Plant Pathology, Faculty of Agriculture, Ain Shams
University.

Dr. Ahmed Ahmed Ahmed Mosa -----

Prof. Emeritus of Plant Pathology, Faculty of Agriculture, Ain Shams
University.

Date of Examination: 24 / 12/ 2018

BIOLOGY AND CONTROL OF BROWN ROT FUNGI ON SOME STONE FRUITS

By

THAURIA MABRUOK MOHAMED ABO EL WAFI

B.Sc. Agric. Sci. (Plant Pathology), Zagazig University, 1993

M.Sc. Agric. Sci. (Plant Pathology), Ain Shams University, 2012

Under the supervision of:

Dr. Ahmed Ahmed Ahmed Mosa

Prof. Emeritus of Plant Pathology, Department of Plant Pathology,
Faculty of Agriculture, Ain Shams University (Principal Supervisor).

Dr. Medhat Kamel Ali El-Sayed

Prof. Emeritus of Plant Pathology, Department of Plant Pathology,
Faculty of Agriculture, Ain Shams University.

Dr. Salah EL-Sayed Youssef

Head Researches of Plant Pathology, Department of Postharvest
Diseases Researches, Plant Pathology Institute, Agriculture Research
Center.

CONTENTS

	Page
LIST OF TABLES	VI
LIST OF FIGURES	VII
INTRODUCTION	1
II. REVIEW OF LITERATURE	3
1. The Diseases: Brown Rot of Stone Fruits and its Causal Pathogen	3
2. Identification and Characterization of <i>Monilinia</i> spp.	10
2.1 Cultural and Morphological Characterization	11
2.2. Molecular Characterization	12
3. Pathogenic variations among <i>Monilinia</i> spp.	15
4. Disease Control:	16
4.1 Chemical control:	18
4.2. Biological control:	20
4..3. Promising Control Alternatives Agents:	22
4.3.1. Salt Compounds:	22
4.3.2. Organic Acids:	25
4.3.3. Essential oils (EOs):	26
III. MATERIALS AND METHODS	28
1. Isolation and identification of <i>Monilinia</i> species	28
1.1. Source of samples:	28
1.2. Pathogen isolation	28
1. 3. Identification:	29
1.3.1. Phenotypic and Microscopic Characteristics:	29
1.3.2 Molecular Identification of <i>Monilinia</i> isolates:	32
3.2.1 Fungal growth on duplex media	32
3.2.2. DNA extraction	32
3.2.3. DNA quantification and gel documentation	33

	Page
3.2.3. DNA quantification and gel documentation	33
3.2.4 RAPD-PCR analysis	33
3.2.5. DNA sequencing	34
3.2.5.1. Preparation of DNA	34
3.2.5.2 ITS-PCR conditions	34
3.2.5.3 Gel electrophoresis	34
3.2.5.4. PCR Product purification	34
3.2.5.5. DNA sequencing	35
3.2.5.6. Identification of isolates using BLAST	35
4. Pathogenicity test:	36
II. Control of Brown rot:	37
1. Effectiveness of certain salts and acids on <i>Monilinia</i> spp., <i>in vitro</i>	37
1.1. Source of <i>Monilinia</i> isolates:	37
1.2. Salts and acids	37
1.3. <i>In vitro</i> experiment:	38
1.4. Preharvest spray of salts and acids, <i>in vivo</i> .	39
1.5. Postharvest treatments with certain salts and acids:	40
1.6 Assessment of brown rot:	41
1.7 Quality assay:	41
1.7.1 Total soluble solids (TSS):	41
1.7.2. Firmness:	41
2. Sensitivity of <i>Monilinia</i> isolates to fungicides	42

	Page
2.1 <i>In viro</i> assay:	42
2.2. Preharvest spray of fungicides	43
2.3. Postharvest treatments with fungicides:	44
3. Biological control:	45
3.1. Source of Bioagents	45
3.2. Dual Culture Assay:	45
3.3. Preharvest treatments with Bio-agent	45
4.Effect of Essential oils (ESo):	46
4.1 <i>In vitro</i> sensitivity of <i>Monilinia</i> spp.	46
4.2. Postharvest treatments with of essential oils:	47
5. Experimental design and statistical analysis:	48
IV. RESULTS	49
Part I: - Biology of <i>Monilinia</i> spp. on Stone Fruits in Egypt	49
1.Incidence of brown rot fungi in stone fruit orchards.	49
2.Identification of fungi causing brown rot.	55
2.1.Morphological characterization of <i>Monilinia</i> spp.	55
2.2Molecular characterization of <i>Monilinia</i> spp.	59
2.2.1.Molecular diversity of <i>Monilinia</i> spp. isolates:	59
2.2.2Analysis of internal transcribed spacer (ITS) regions 1 and 4	72
3. Pathogenicity of <i>Monilinia</i> isolates peaches and apricot:	73
Part II:- Control of <i>Monilinia</i> spp., the causal of Brown Rot disease of stone Fruits:	75
1- <i>In Vitro</i> Assays	75
1.1. <i>In vitro</i> effectiveness of certain salts and acids on growth of three <i>Monilinia</i> spp.	75
1.2. <i>In vitro</i> effectiveness of four fungicide on three <i>Monilinia</i> spp.	78
1.3. <i>In vitro</i> effectiveness of Essential oil on <i>Monilinia laxa</i> , <i>M. fructigena</i> and <i>Monilinia fructicola</i> . isolate No. 118:	83

	Page
1.4. Effect of some bio-control agents on growth on <i>Monilinia laxa</i> , <i>M. fructigena</i> and <i>M. fructicola</i> :	87
2.Control of Brown rot disease of:Peaches ,in vivo	89
2.1.Control of brown rot of peaches during cold storage by preharvest applications with salts and acids:	89
2.2 Quality of peaches during cold storage preharvest sprayed with tested salts and acids:	91
2.3. Control of brown rot of peach Florda Prince cv .during cold storage by preharvest application with fungicides:	94
2.4. Quality of peach FlordaPrince cv .during cold storage preharvest sprayed with fungicides:	96
2.5. Control of brown rot of peach during cold storage, by preharvest application with bioagents:	98
2.6. Quality of peach during cold storage, preharvest sprayed with bioagents:	100
2.7. Control of brown rot of peach FlordaPrince cv . by postharvest applications of salts and acids:	102
2.8. Quality of peach FlordaPrince cv . postharvest treated with tested salts and acids during cold storage:	104
2.9. Control of brown rot of peach FlordaPrince cv ., by postharvest applications of natural oils:	106
2.10. Quality of peach FlordaPrince cv .postharvest treated with natural oils during cold storage:	108
3. Control of Brown rot disease on apricot,in vivo	110
3.1 Control of brown rot during cold storage of apricot by preharvest application with certain salts and acids:	110
3.2 Quality of apricot preharvest during cold storage sprayed with tested salts and acids during cold storage:	112
3.3 Control of brown rot of apricot by preharvest applications with fungicides:	115

	Page
3.4 Quality of apricot preharvest sprayed with fungicides during cold storage:	117
3.5. Control of brown rot of apricot by preharvest applications with bioagents:	119
3.6. Quality of apricot preharvest sprayed with bioagents during cold storage:	121
3.7 Control of brown rot of apricot by postharvest applications of salts and acids:	123
3.8 Quality of apricot postharvest treated with tested salts and acids during cold storage:	125
3.9 Control of brown rot of apricot, by postharvest applications of natural oils:	127
3.10 Quality of apricot postharvest treated with natural oils during cold storage:	129
DISCUSSION	132
SUMMARY	147
REFERENCES	155
APPENDIX	173
ARABIC SUMMARY	189

LIST OF TABLES

Table No.		Page
1	Morphological characteristics and germ tube branching distinguishing among different <i>Monilinia</i> and <i>Monilinia</i> spp. (Lane, 2002), Poniatowska <i>et al.</i> (2013) and Lichtemberg (2014).	31
2	RAPD polymorphic decamer primers	33
3	Fungicides used in this study	42
4	Frequency of occurrence of <i>Monilinia</i> spp. Causing brown rot disease on different phenological stages of peaches, nectarines and apricot from flowering until ripe fruit in 4 Governorates during seasons 2013-2014.	50
5	Total no. of isolates obtained from three stone fruit crops during 2013-2014, in Egypt.	51
6	Similarity matrix, in percentage, among the analyzed <i>Monilinia</i> spp. isolates based on RAPD band pattern analysis according to Primer 1.	64
7	Similarity matrix, in percentage, among the analyzed <i>Monilinia</i> spp. isolates based on RAPD band pattern analysis according to Primer 2.	66
8	Gene sequence with universal primer	67
9	Isolates analyzed by PCR amplification of rDNA	68
10	Sequences of <i>Monilinia</i> from NCBI GeneBank used for construction of phylogenetic tree and GeneBank Accession numbers.	69
11	Pathogenicity of different isolates of <i>Monilinia fructicola</i> . on peach cv. FloridaPrince and apricot cv. Canino .	74
12	Effect of sodium bicarbonate, calcium chloride, potassium sorbate, potassium silicate, boric acid and salicylic Acid on linear growth (mm) of <i>Monilinia</i> spp. on PDA .	77
13	Models correlating salts and acids concentrations and their efficacy to inhibit the growth of <i>Monilinia fructicola</i> . on	

Table No.		Page
	PDA medium <i>in vitro</i> and their IC ₅₀	78
14	Effect of pyraclostrobin + boscalid formulation, iprodione, Reynoutria extract and cyprodinil + fludioxonil formulation on growth rate of <i>Monilinia fructigena</i> , <i>M. laxa</i> and <i>Monilinia fructicola</i> . isolate no. 118 <i>in vitro</i> at 22°C for 7 days	80
15	Models correlating fungicide concentrations and their efficacy to inhibit the growth of <i>Monilinia</i> spp. on PDA medium, <i>in vitro</i> , and their IC ₅₀	82
16	Effect of thyme, camphor, cinnamon and carnation on linear growth of <i>Monilinia fructicola</i> . isolates no. 118, <i>M. laxa</i> and <i>M. fructigena</i> on PDA medium at 22°C for 7 days.	84
17	Models correlating plant oils concentrations and their efficacy to inhibit the growth of <i>Monilinia</i> spp. on PDA medium, <i>in vitro</i> , and their IC ₅	86
18	effect of four bioagents, i.e. <i>Trichoderma harzianum</i> (Th5), <i>Trichoderma harizanium</i> (Th20) <i>Bacillus cereus</i> (BC) and <i>Bacillus amyloliquefaciens</i> (BA), on mycelia growth of three isolates of <i>Monilinia</i> spp. grown in dual PDA culture plates and incubated for 7 days at 22 °C	88
19	Effect of preharvest spraying of Peach trees with certain salts and acids on brown rot incidence (DI) and severity (DS) of fruits, collected Qalyubia and Ismailia Governorates, during Seasons 2015 and 2016, and kept after harvest at 0°C and 90% RH during cold storage for up to 30 days*	90
21	Effect of preharvest spraying of Peach trees with different bioagents on Brown rot Incidence (DI) and severity (DS) of fruits, collected Qalyoubia and Ismailia governorates, during seasons 2015 and 2016, and kept after harvest at	

Table No.		Page
	0°C and 90% RH during cold storage for up to 30 days*	99
22	Effect of posth Effect of postharvest dipping of Fruits Peaches with certain salts and acids on Brown rot Disease Incidence (DI) and Diseases severity (DS) kept at 0°C cold storage and 90% RH for up to 30 days during seasons 2016.	103
23	Brown rot incidence and severity percentages on peach fruits, postharvest treated with essential oil during cold storage at 0°C and 90% RH for up to 30 days, during seasons 2015 and 2016.	107
24	Effect of preharvest spraying of Apricot trees with certain salts and acids on brown rot incidence (DI) and severity (DS) of fruits, collected Qalyubia and Ismailia Governorates, during Seasons 2015 and 2016, and kept after harvest at 0°C and 90% RH during cold storage for up to 30 days	111
25	Effect of preharvest spraying of Apricot trees with different Fungicides on Brown rot incidence (DI) and severity (DS) of Fruits, collected Qalyubia and Ismailia governorates during seasons 2015 and 2016, and kept after harvest at 0°C and 90% RH during cold storage for up to 30 days	116
26	Effect of preharvest spraying of Apricots trees with different bioagents on Brown rot incidence (DI) and severity (DS) of fruits, collected Qalyoubia and Ismailia governorates during seasons 2015 and 2016, and kept after harvest at 0°C and 90% RH during cold storage for up to 30 days*	120
27	Effect of postharvest dipping of Fruits Apricot with certain salts and acids on Brown rot Disease Incidence	

Table No.		Page
	(DI) and Diseases severity (DS) kept at 0°C cold storage and 90% RH for up to 30 days during seasons 2016.	124
28	Brown rot incidence and severity percentages on apricot postharvest treated with essential oil during cold storage at 0°C and 90% RH for up to 30 days during seasons 2015 and 2016.	127

LIST OF FIGURES

Fig. No.		Page
1	Frequency of occurrence of different <i>Monilinia</i> spp. Obtained from peach, apricot and nectarine grown at different orchards in four Governorates, during 2013-2014, in Egypt.	53
2	Colony characteristics(A-G) and structures (H-L) of <i>Monilinia</i> spp. isolates, from peach, apricot and nectarine in Egypt, grown on PDA (10 days, 12 h light /12 h dark at 22 °C). (A) Sporulation (abundant); (B) Concentric rings; (C) Lobed margin (upper surface); (D) Lobed margin (lower surface); (E) Rosetting margin (upper surface); (F) Rosetting margin (lower surface); (G) Black or brown arcs; (H) Stromata; (I) Apothecia; (J) Apothecia (close-up); (K) Apothecia 100x; (L) Stromata 100x.	57
3	Colony morphology of four <i>Monilinia</i> species, grown on PDA (10 days, 12 h light /12 h dark at 22 °C. (A): <i>M. fructigena</i> , isolate No112; (B): <i>M. fructicola</i> isolate No 118; (C): <i>M. laxa</i>	58
4	Dendrogram of main obtained <i>Monilinia</i> spp. isolates from stone fruits and twigs	61
5	DNA bands of 10 <i>Monilinia</i> spp. isolates from stone fruits as determined using primer 1 for the RAPD/PCR band pattern.	63
6	Dendrogram obtained by clustering (UPGMA method) based on the band pattern obtained by the RAPD PCR analysis for ten <i>Monilinia</i> fungal isolates when using primer 1.	64
7	DNA bands of 10 <i>Monilinia</i> spp. isolates from stone fruits as determined using primer 2 for the RAPD/PCR band pattern.	65
8	Dendrogram obtained by clustering (UPGMA	

Fig. No.		Page
	method) based on the band pattern obtained by the RAPD PCR analysis for ten <i>Monilinia</i> fungal isolates when using primer 2.	66
9	DNA bands of 4 <i>Monilinia</i> isolates from stone fruits as determined using primer ITS1, ITS2 for the band pattern.	67
10	Sequence alignment of the ITS region of <i>Monilinia</i> species and selected isolates. Isolates and GeneBank Accession numbers	68
11	RAxML tree of the genomic region based on analyses of a combined dataset of ITS sequence data.	70
12	Sequence alignment of the genome area corresponding to the <i>Monilinia fructigena</i> (ARC5,7) 508-bp , <i>Monilinia fructicola</i> (ARC1) 497pb and <i>Monilia laxa</i> (ARC4)509 -pb I TS region product for the three <i>Monilinia fructicola</i> . and Identical nucleotides are represented by stars, and absent nucleotides are shown by hyphens. Spaces at beginning and end of alignment are undetermined sequence	71
13a	Quality of peach preharvest-sprayed ,three times, with certain salts and acids at growth stages during growing seasons 2015 at Qalyoubia and kept after harvest at 0°C and 90% RH	92
13b	Quality of peach preharvest-sprayed, three times, with certain salts and acids at growth stages During Growing seasons 2016 Qalyoubia and kept after harvest at 0°C and 90% RH	92

Fig. No.		Page
14a	Quality of FlordaPrince peaches preharvest-sprayed three times with certain salts and acids at growth stages during growing seasons 2015 Ismailia kept after harvest at 0°C and 90% RH	93
14b	Quality of FlordaPrince peaches preharvest-sprayed three times with certain salts and acids at growth stages during growing seasons 2016 Ismailia kept after harvest at 0°C and 90% RH	93
15	Quality of peaches preharvest-sprayed three times with fungicide at growth stages during growing seasons 2015 in Qalyubia and Ismailia governorates and kept after harvest at 0°C and 90% RH.	97
16	Quality of peaches preharvest-sprayed three times with Bioagent at growth stages during growing seasons 2016 in Qalyubia ismailia and governorates and kept after harvest at 0°C and 90% RH	101
17a	Quality of peaches postharvest-treated with certain salts and acids and kept at 0°C cold storage and 95% RH for up to 30 days during season 2016.	107
17b	Quality of peaches postharvest-treated with certain salts and acids and kept at 0°C cold storage and 95% RH for up to 30 days during season 2016.	107
18a	Quality of FlordaPrince peaches post – harvest treatments with essential oil 0°C and 90% RH during cold storage for up to 30 days during growing seasons 2016.	109
18b	Quality of FlordaPrince peaches post – harvest treatments with essential oil 0°C and 90% RH during cold storage for up to 30 days during growing seasons 2016.	109
19a	Quality of Canino apricot preharvest-sprayed three times with certain salts and acids at growth stages during growing seasons 2015 in Qalyubia	