

**FLUORESCENT PSEUDOMONADS AS PLANT
GROWTH PROMOTERS AND BIOCONTROL
OF ROOT-INFECTING PATHOGENS
ON MAIZE PLANT**

By

OSAMA ELSAYED MOHAMED

B.Sc.Agric.Sc. (Biotechnology), Ain-Shams University, 2012

**A Thesis Submitted in Partial Fulfillment
Of
The Requirement for the Degree of**

**MASTER OF SCIENCE
in
Agricultural Sciences
(Agricultural Microbiology)**

**Department of Agricultural Microbiology
Faculty of Agriculture
Ain-Shams University**

2019

Approval Sheet

**FLUORESCENT PSEUDOMONADS AS PLANT
GROWTH PROMOTERS AND BIOCONTROL
OF ROOT-INFECTING PATHOGENS
ON MAIZE PLANT**

By

OSAMA ELSAYED MOHAMED

B.Sc.Agric.Sc. (Biotechnology), Ain-Shams University, 2012

This thesis for M. Sc. degree has been approved by:

Dr. Mohammed Zakaria Sedik
Prof. Emeritus of Agric. Microbiol, Fac. of Agric., Cairo University

Dr. Shawky Mahmoud Selim
Prof. of Agric. Microbiol, Fac. of Agric., Ain-Shams University

Dr. Enas Abd El-Tawab Hassan.....
Prof. of Agric. Microbiol, Fac. of Agric., Ain-Shams University

Dr. Elshahat Mohamed Ramadan
Prof. Emeritus of Agric. Microbiol, Fac. of Agric., Ain-Shams
University

Date of Examination: 16/12/2018

FLUORESCENT PSEUDOMONADS AS PLANT GROWTH PROMOTERS AND BIOCONTROL OF ROOT-INFECTING PATHOGENS ON MAIZE PLANT

By

OSAMA ELSAYED MOHAMED

B.Sc.Agric.Sc. (Biotechnology), Ain-Shams University, 2012

Under the supervision of:

Dr. Elshahat Mohamed Ramadan

Prof., Emeritus of Agricultural Microbiology, Department of Agric.
Microbiology, Faculty of Agriculture, Ain-Shams University
(Principal Supervision).

Dr. Enas Abd El-Tawab Hassan

Prof. of Agricultural Microbiology, Department of Agric.
Microbiology, Faculty of Agriculture, Ain-Shams University.

Dr. Elhamy Mohamed Mostafa El-Assiuty

Head Research Emeritus of Plant Pathology, Plant Pathology
Research Institute, Agricultural Research Centre, Giza

ABSTRACT

Osama Elsayed Mohamed: Fluorescent Pseudomonads as Plant Growth Promoters and Biocontrol of Root-Infecting Pathogens on Maize Plant. Unpublished M. Sc. Thesis, Department of Agricultural Microbiology, Faculty of Agriculture, Ain Shams University, 2019.

Possibility of manipulating some of the efficient strains of rhizospheric fluorescent pseudomonads to manage the root-infecting pathogens of maize was studied throughout this study. Out of 110 isolates, 24 of *Pseudomonas* species, recovered from the rhizosphere of maize and sugar beet showed high antagonistic effect against two major root-infecting pathogens of maize, namely *Cephalosporium maydis* and *Fusarium verticillioides* *in vitro*. Pot experiment revealed that just 4 isolates could reduce infection with both pathogens and enhance the plant growth as well. Based on the genotypic identifications, they could be identified as: *Pseudomonas putida* Pau9, *P. putida* Pau11, *P. putida* Psf3 and *P. aeruginosa* Psf9. Assay of extracellular enzymes revealed that cellulase was actively produced, only by *P. aeruginosa* Psf9. Chitinase, however was detected in growing media of three strains, but not by *P. putida* Pau11. Assay of antibiotics produced by the bacterial strains showed that phenazine could, only be produced by *P. aeruginosa* Psf9. HCN was found to be excreted by *P. putida* Psf3 and *P. aeruginosa* Psf9. Except *P. putida* Pau9, the IAA could be produced by the other three strains. All strains were able to produce siderophores, and caused availability of P and K. GC-MS analyses revealed that different compounds were detected within the metabolites produced by each of *Pseudomonas* strains under study. Coating maize seed with the mixture of the four strains and seeding in potted-soil infested singly with one of the two target pathogens, or in combination revealed that fresh and dry weights of resulting plants were significantly increased compared with the control. Treatment caused significant increase in root lengths, but insignificant increase in shoot lengths. This is correct in soil infested with

C. maydis. Whereas, in soil infested with *F. verticillioides*, treatment caused significant results in all of the growth parameters of plants. Combining the two pathogens in soil showed that insignificant effect of coating seed with the mix of bacteria on any of shoot or root lengths. Whereas, significant results were found in shoot and root weights compared to the non-treated control.

Keywords: Fluorescent Pseudomonads, *Cephalosporium maydis*, *Fusarium verticillioides*, Maize plant, GC-MS analyses bioagent.

ACKNOWLEDGEMENT

Praise and thanks be to ALLAH, the most merciful for assisting and directing me to the right way.

There are few opportunities in the most people's lives to officially prove their gratitude to people who were mentors and supporters at different stages of our lives. However, we do not forget to seal those feelings on paper.

I would like to express my deep gratitude to **Dr. Elshahat Mohamed Ramadan** Prof. of Agricultural Microbiology, Faculty of Agriculture, Ain Shams University, for his supervision, helpful encouragement, critical reading and revising the manuscript.

I would like to express my sincere thanks, and deep gratitude to **Dr. Enas Abd El-Tawab Hassan** Prof. of Agricultural Microbiology, Faculty of Agriculture, Ain Shams University, for her fruitful discussion, criticism and valuable help throughout of this study.

Deep thanks and appreciation are extended to **Dr. Elhamy Mohamed Mostafa El-Assiuty**, Chief of Research, Plant Pathology Research Institute, Agricultural Research Centre, for his encouragement, motivation and providing all needs and facilities during the preparation of this thesis.

Cordially, it is my pleasure to express my deep gratitude and sincere appreciation to **Dr. Zeinab Mohamed Fahmy** Chief of Research, Plant Pathology Research Institute, A.R.C, for her support and valuable encouragement throughout executing this study.

I would, also like to thank all members of Maize and Sugar Crops Research Department, Plant Pathology Research Institute, A. R. C. for their cordial assistance.

Finally, I would like to thank my parents, my wife, brothers and sister for their love, patience, interest, support and continuous encouragement.

CONTENTS

	Page
LIST OF TABLE.....	
LIST OF FIGURES.....	
INTRODUCTION.....	1
REVIEW OF LITERATURE.....	3
1. Maize plant as an economical crop.....	3
2. Pathogens infected maize plant	3
3. Fluorescent pseudomonads involved in suppression of plant root pathogens and improving plant growth	4
A. Antagonism by production of lytic enzymes.....	5
B. Phytohormones production	6
C. Production of siderophores	7
D. Antibiotic production	7
E. Production of Hydrogen cyanide (HCN)	8
F. Phenolic compound	9
G. Phosphate solubilization	10
H. Potassium solubilization	10
4. Performance of plant affected by microbial inoculants	11
MATERIAL AND METHODS.....	13
1. Isolation and identification of maize infected pathogens.....	13
A. <i>Cephalosporium maydis</i> (<i>Harpophora maydis</i>).....	13
B. <i>Fusarium verticillioides</i> (<i>F.moniliforme</i>).....	13
2. Isolation of rhizospheric fluorescent pseudomonads	14
3. Assessment the antagonistic activity of fluorescent pseudomonads.....	14
A. <i>In vitro</i> antagonistic activity.....	14
B. <i>In vivo</i> evaluation of antagonist bacterial isolates	15
4. Identification of the most efficient <i>Pseudomonas</i> isolate.....	15
4.1. Biochemical characterization of <i>Pseudomonas</i> spp.....	15
4.2. Molecular identification of <i>Pseudomonas</i> spp.....	15

II

5. Molecular identification of <i>Fusarium</i>	16
5.1. DNA extraction.....	16
5.2. PCR amplification	17
6. Molecular identification of <i>Cephalosporium</i>	17
6.1. DNA extraction	17
6.2. DNA extraction	17
7. Phylogenetic analysis	18
8. Characterization of antagonistic bacteria	18
8.1. Production of bacterial lytic enzymes.....	18
A. Production of cellulase.....	18
B. Pectinase activity.....	19
C. Chitinase activity.....	19
8.2. Detection of antibiotic production genetically.....	19
8.3. Phosphate solubilization.....	21
8.3.1. Qualitative assay.....	21
8.3.2. Quantitative assay.....	21
8.4. Potassium solubilization.....	21
8.5. Production of indole 3- acetic acid (IAA).....	22
8.6. Production of HCN.....	22
8.7. Quantitative assessment of siderophore production.....	22
1- Preparation of standard curve of deferoxamine mesylate	22
2- Preparation of culture for examination.....	23
3- Quantitative siderophores.....	23
9. Characterization of antifungal metabolites by Gas Chromatography Mass Spectrometry (GC-MS).....	24
10. Effect of bioagent on fungal pathogen infected maize plant, under pot experiment.....	25
10.1. Parameters measured.....	25
A. Vegetative parameters.....	25
B. Chemical analysis	25
1- Nitrogen, phosphorus and potassium content	25
2- Chlorophyll and beta carotenoid contents	26

III

a. Enzyme activity assay	26
11. Statistical analysis.....	26
RESULTS	27
1. Identification of pathogenic fungus of maize plant.....	27
1.1. Phenotypic identification of <i>Cephalosporium maydis</i>	27
1.2. Genotypic identification of <i>Cephalosporim maydis</i>	27
1.3. Phenotypic identification of <i>Fusarium</i>	27
1.4. Phylogenetic analysis of <i>Fusarium</i>	27
2. Isolation of the fluorescent pseudomonads.....	30
3. Antifungal activity.....	30
4. Evaluation of selected cultures as biocontrol agent <i>in vivo</i> ...	31
5. Biochemical characterization of fluorescent pseudomonads...	33
6. Genotypic identification for the most potent antagonist bacterial isolates.....	34
7. Biological activities of the efficient bacterial strains.....	37
7.1. Production of extracellular enzymes	37
7.2. Antibiosis of <i>Pseudomonas</i> strains	38
7.3. Production of hydrogen cyanide (HCN).....	38
7.4. Production of siderophores.....	38
7.5. Production of phytohormones by <i>Pseudomonas</i> strains.....	39
7.6. Phosphate solubilization.....	40
7.7. Potassium availability.....	40
8. Analyses of secondary metabolites produced by the four efficient <i>Pseudomonas</i> strains.....	41
9. Performance of maize plant in soil infested with root-infecting pathogen and biocontrol agent.....	46
9.1. Vegetative parameters.....	46
9.2. Accumulation of pigment and enzyme activity of mize plant	51
DISCUSSION	54
SUMMARY	60
REFERENCES	63
ARABIC SUMMARY	

LIST OF TABLES

No.		Page
1	Sequence of specific primer used for <i>Pseudomonas</i> spp identification.....	16
2	Primers for PCR analysis of tested antibiotics.....	20
3	Number of fluorescent pseudomonads isolates recovered from different sources.....	30
4	Efficiency of coating maize seed with fluorescent pseudomonad isolates in managing root-infecting pathogens of maize under greenhouse conditions.....	32
5	Biochemical characterization for identification of the most potent <i>Pseudomonas</i> isolates.....	33
6	Assessment of enzymatic activity of selected <i>Pseudomonas</i> strains.....	37
7	Potency of <i>Pseudomonas</i> strains for producing antimicrobial compounds.....	39
8	Production of IAA and gibberellins by tested strains.....	39
9	GC-MS analyses for the metabolites produced by <i>Pseudomonas putida</i> psf 3 in liquid culture.....	42
10	GC-MS analyses for the metabolites produced by <i>Pseudomonas aeruginosa</i> psf 9 in liquid culture.....	43
11	GC-MS analyses for the metabolites produced by <i>Pseudomonas putida</i> Pau 9 in liquid culture.....	44
12	GC-MS analyses for the metabolites produced by <i>Pseudomonas putida</i> Pau11 in liquid culture.....	45
13	Effect of the combination of the four bacterial mixture and <i>C.maydis</i> on parameters of maize plant growth under greenhouse conditions.....	47
14	Effect of the combination of the four bacterial mixture and <i>F. verticillioides</i> on growth parameters of maize plants under greenhouse conditions.....	48

No.		Page
15	Effect of the combination of the four bacterial mixture and <i>C.maydis</i> + <i>F. verticillioides</i> on growth parameters of maize plants under greenhouse conditions.....	48
16	Effect of the combination of four bacterial mix and fungal pathogens on chlorophyll and carotenoides contents of mize plant under greenhouse condition.....	51
17	Effect of combination of bacterial mixture and fungal pathogens on peroxidase activity of maize plant under greenhouse conditions.....	52