

بسم الله الرحمن الرحيم



-C-02-50-2-





شبكة المعلومات الجامعية التوثيق الالكتروني والميكرونيلم





جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة يعيدا عن الغيار







بالرسالة صفحات لم ترد بالأصل









EVALUATION OF THE EFFECT OF A PLANT ALKALOID (BERBERINE DERIVED FROM BERBERIS ARISTATA) ON T. VAGINALIS IN VITRO

Ву

GHADA ABDEL RAHMAN SAAD M. B, B. Ch.

Demonstrator of Parasitology, Parasitology Department, Faculty of Medicine, Ain Shams University

Thesis
Submitted for the partial fulfillment
of

Master degree of Parasitology

Supervised by

65874

PROF. DR. SALWA AHMED SOFFAR

Professor of Parasitology, Parasitology Department, Faculty of Medicine, Ain Shams University.

DR. DOREYA MOHSEN MAHMOUD

Assistant Prof. of Parasitology, Parasitology Department, Faculty of Medicine, Ain Shams University.

DR. SAHAR SAFWAT ABDEL AZIZ

Lecturer of Parasitology, Parasitology Department, Faculty of Medicine, Ain Shams University.

مال مرور

Gh. A

Faculty of Medicine Parasitology Departmen®

Ain Shams University

2000

wift.

p

* Pag

بسم الله الرحمن الرحيم

TEXT.

A. C.

.** _,†

Acknowledgement

I would like to express my supreme gratitude and appreciation to Prof. Dr. Salwa Ahmed Soffar, Professor of Parasitology, Faculty of Medicine, Ain Shams University for the enormous effort she did in supervising this thesis, besides her precious help, constant guidance and sincere encouragement.

My special thanks and appreciation to Dr. Doreya Mohsen Mohamoud, assistant Prof. of Parasitology, Faculty of Medicine, Ain Shams University, for her guidance, assistance, generous cooperation, and meticulous supervision on every part of this thesis.

I would like to express my deepest thanks to Dr. Sahar Safwat Abdel Aziz, Lecturer of Parasitology, Faculty of Medicine, Ain Shams University, for her enormous effort, indispensable assistance, and sincere help for revision and fulfillment of this work.

I am deeply grateful to Prof. Dr. Magda El-Sayed Azab, Professor and Head of Parasitology Department, Faculty of Medicine, Ain Shams University for her assistance and precious help in providing the samples needed for this work.

Special words of thanks go to Dr. Hala Salah Ismial El-Wakil, Lecturer of Parasitology, Faculty of Medicine, Ain Shams University, who gave me free use of the materials

needed for this work, and for her assistance during culture maintenance

Last but not least, I would like to thank all members of the Parasitology Department, Faculty of Medicine, Ain Shams University for their kind help.

CONTENTS

	Page No.
INTRODUCTION	1-2
REVIEW OF LITRATURE	3-48
AIM AND PLAN OF THE WORK	49-50
MATERIALS AND METHODS	51-60
RESULTS	61-117
DISCUSSION	118-127
SUMMARY -	128-129
REFERENCES	130-157

·
·
·
·
·
· . .

List of Tables and Figures

Table (1):	Antigens of <i>T. vaginalis</i> and their characteristics	_18
Table (2):	Differential diagnosis of vaginal discharge	. 20
Table (3):	The mean count of <i>T. vaginalis</i> organisms in tubes of culture media after exposure to various concentrations of BS in comparison to control tubes on 3 successive days.	_ 99
Table (4):	The mean count of <i>T. vaginalis</i> organisms previously exposed to various concentrations of BS upon reculture in drug free culture tubes in comparison with control tubes.	100
Table (5):	The mean count of <i>T. vaginalis</i> organisms in tubes of culture media after exposure to various concentrations of metronidazole in comparison to control tubes on 3 successive days.	-101
Table (6):	The mean count of <i>T. vaginalis</i> organisms previously exposed to various concentrations of metronidazole upon reculture in drug free culture tubes in comparison with control tubes.	102
Table (7):	The mean count of <i>T. vaginalis</i> organisms in tubes of culture media after exposure to various concentrations of BS in comparison to mean count of organisms in tubes exposed to various concentrations of metronidazole.	103

Table (8):	Percentage of motility and viability of <i>T. vaginalis</i> on exposure to different concentrations of BS in comparison to control tubes on 3 successive days. — — — — — — — —	105
Table (9):	Percentage of motility & viability of <i>T. vaginalis</i> on exposure to different concentrations of metronidazole in comparison to control tubes on 3 successive days	107
Fig., (1):	Structure of T. vaginalis	8
Fig., (2):	Chemical structure of Metronidazole	34
Fig., (3):	Chemical structure of berberine sulphate	41
Fig., (4):	Curve illustrating relationship between various concentrations of BS and count of <i>T. vaginlis</i> in culture.	108
Fig., (5):	Curve illustrating relationship between various concentrations of metronidazole and count of <i>T. vaginalis</i> in culture.	109
Fig., (6):	Curve illustrating relationship between various concentrations of BS and percentage of motility of <i>T. vaginalis</i> .	llo
Fig., (7):	Curve illustrating relationship between various concentrations of BS and percentage of viability of T. vaginalis.	111