


سامية محمد مصطفى



شبكة المعلومات الجامعية

بسم الله الرحمن الرحيم




سامية محمد مصطفى



شبكة المعلومات الجامعية



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



سامية محمد مصطفى



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم


قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار




سامية محمد مصطفى



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة




سامية محمد مصطفى



شبكة المعلومات الجامعية



بالرسالة صفحات لم ترد بالأصل



“Cone Beam Computed Tomography Comparative Assessment of Maxillofacial Morphology in Patients with Bilateral Cleft Lip and Palate”

Thesis

Submitted to the faculty of Dentistry, Ain Shams University,

In partial fulfillment of the requirements for the Master degree

In Oral and Maxillofacial Radiology

Presented by

Yomna Osama Dardeer Abdelmalek Elhoseiny

Yomna.elhoseiny@gmail.com.01011102765

B.D.S, Ain Shams University- 2010

Under the supervision of

Prof.Dr. Mona Abo-El-Fotouh

Professor of Oral and Maxillofacial Radiology

Head of department of Oral Medicine, Periodontology and Oral Radiology

Faculty of Dentistry- Ain Shams University

Prof.Dr.Marwa Abdel Wahab Elkassaby

Professor of Oral and Maxillofacial Surgery

Head of department of Oral and Maxillofacial Surgery

Faculty of Dentistry- Ain Shams University

Dr.Raghdaa Abo Elkhair Bayoumy Mostafa

Lecturer of Oral and Maxillofacial Radiology

Faculty of Dentistry- Ain Shams University

Faculty of Dentistry

Ain Shams University (2019)

Dedication

A special feeling of gratitude to my loving, caring and supporting parents and family

Acknowledgment

First of all, Praise to ALLAH, who without HIS support, I could not finish this work.

I would like to express my sincere gratitude and deepest thanks to my supervisor **Prof. Dr. Mona Abo-El-Fotouh**, professor and Head of Oral and Maxillofacial Radiology Department, Faculty of Dentistry, Ain Shams University, for her help, direct supervision, sincere encouragement and patience, during this work.

I would like to express my gratefulness and appreciation for **Prof. Dr. Marwa Elkassaby**, professor and Head of Oral and Maxillofacial Surgery Department, Faculty of Dentistry, Ain Shams University, who gave up much of her time and deep experience to allow the pursuit of this work.

I am greatly thankful to my friend and supervisor **Dr Raghdaa Abo Elkair**, lecturer of Oral and Maxillofacial Radiology, Faculty of Dentistry, Ain Shams University, for her valuable advices and continuous support. I am deeply grateful to her for her skillful supervision and valuable opinions.

I would like to thank my colleagues in the Oral Radiology Department for their support and help whenever asked for.

List of contents

Dedication.....	i
Acknowledgment.....	ii
List of figures.....	iii
List of tables.....	ix
List of abbreviations.....	xi
Introduction.....	1
Review of literature.....	4
Aim of study.....	39
Patients and methods.....	40
Results.....	63
Case (1) presentation.....	79
Case (2) presentation.....	89
Discussion.....	99
Summary.....	111
Conclusion.....	113
Recommendations.....	114
References.....	115
Arabic summary.....	

List of figures

<u>Figure number</u>	<u>Title</u>	<u>Page number</u>
1	Facial primordia responsible for development of the face	6
2	Figure 2: Development of the secondary palate T = tongue, P = palatal shelf, a) formation of the palatal shelves, b) elevation of the shelves to a horizontal orientation, c) Fusion of the shelves and separation of the nasal and oral cavities	7
3	Veau classification	8
4	At 24 weeks, 3D surface-rendered imaging (coronal plane) showed bilateral cleft lip with protrusion of the premaxilla	14
5	Three-dimensional US image in a 32 weeks fetus showing left unilateral cleft lip	15
6	Sonographic coronal view showing the retranasal triangle with higher echogenicity than surroundings	16
7	Basic principle of CBCT	18
8	Comparison of volume data sets obtained isotropically (left) and anisotropically (right)	19
9	Different FOV used in CBCT a) large FOV b) medium FOV c) restricted FOV	21
10	Component of image intensifier tube/charge-coupled device	23
11	Types of FPD	24
12	Standard display modes of CBCT	25
13	Oblique planar reformation used for examination of mandibular condyle	26
14	Oblique reformation. Lines in the axial plane (left) indicate the rotated sagittal (middle) and coronal (right) planes	27
15	Reconstructed panorama	27
16	a)Position of cross-sectional images, perpendicular to the panoramiccurve, displayed on an axial slice, b) Cross-sectional images at various positions along the curve	28

17	Different 3-D visualization techniques applied to CBCT dataset of a patient with a craniofacial deformity	30
18	Sagittal slice showing different anatomical landmarks.	42
19	Adjusting sagittal plane in coronal image such that it passes through middle of crista galli posteriorly (A) and lower central incisors anteriorly (B)	43
20	Sagittal image showing SNA angle measured using angle tool S = sella; N = nasion; ; A = point A, SNA = 89.8°	44
21	a) Coronal slice showing tip point of nasal septum and point O, b) Measuring ASD using angle tool , point O = most deviated part of the nasal septum ,ASD = 10.1°	46
22	Sagittal slice showing ANB angle measured using angle tool, ANB = 12.6°	47
23	Sagittal image showing AUFH measured as straight line from N =nasion to ANS = anterior nasal spine using ruler tool, AUFH = 39.8mm	48
24	Sagittal slice showing ALFH measured using ruler tool, ANS = anterior nasal spine; Me = menton, ALFH = 78.4mm	49
25	Sagittal slice showing nasal tip projection measured as the linear horizontal distance from tip of nose to a straight line tangent to nasion (N) and pgonion (Pg), TP = 27.3mm	50
26	Sagittal slice demonstrating the nasal bone length measured from the most anterior point of the frontonasal suture to end point of the nasal bone, nasal bone length = 21.6	51
27	Coronal slice demonstrating the width of nasal bone aperture measured at the widest points. width of nasal bone aperture = 26.5 mm	52
28	a) Locating ANS = anterior nasal spine on axial slice, b) measuring distance from the point, corresponding to the ANS at the level of maxillary tuberosity, to a horizontal line joining the most posterior parts of maxillary tuberosity = 46.1mm	53

29	Axial slice showing Zygomatic prominence on right and left side, R =8.02mm, L =8.61mm	54
30	ABD measured apical to central incisor, ABD = 45.3	55
31	Selecting VOI overlay tool and adjusting borders around the maxillary sinus in axial, sagittal and coronal planes	56
32	Adjusting the rendering histogram	57
33	Object tool selected, starting points placed on coronal image and the desired amount of growth entered	57
34	Changing amount of growth	58
35	Shifting to [Expand/Shrink] tool and entering value in voxel unit	58
36	Choosing select as a new object	59
37	Segmented maxillary with volume calculated, maxillary sinus volume = 4.89cc	59
38	a) Selecting target position b) Adjusting the cube size	60
39	3D zoom function used for teeth assessment	60
40	Assessment of the right side	61
41	Assessment of the left side	61
42	Bar graph showing difference in ASD among the two groups	64
43	Bar graph showing difference in ANB among the two groups	65
44	Bar graph showing difference in AUFH/ALFH among the two groups	66
45	Bar graph showing difference in TP among the two groups	67
46	Bar graph showing difference in nasal bone length among the two groups	68
47	Bar graph showing difference in width of nasal aperture among the two groups	69
48	Bar graph showing difference in the relation of premaxilla to maxilla among the two groups	70
49	Bar graph showing difference in zygomatic prominence among the two groups	71

50	Bar graph showing difference in zygomatic prominence between right and left side in group P (a) and in group R (b)	72
51	Bar graph showing difference in ABD among the two groups	73
52	Bar graph showing difference in maxillary sinus volume (MSV) among the two groups	74
53	Bar graph showing difference in maxillary sinus volume (MSV) among right and left side of group P (a) and group R (b)	75
54	Bar graph showing significant difference in number of teeth in premaxilla among the two groups	76
55	Sagittal image showing SNA angle; S = sella; N = nasion; A = point A, SNA= 89.6°	79
56	Coronal image showing ASD; point 0 = most convex point of the deviated nasal septum, ASD = 9°	80
57	Sagittal image showing ANB angle; A = point A; N = nasion; B = point B, ANB=13.7°	80
58	Sagittal image showing AUFH; N = nasion; ANS = anterior nasal spine, AUFH = 39.7 mm	81
59	Sagittal image showing ALFH; ANS = anterior nasal spine; Me = menton, ALFH = 78.6 mm	81
60	Sagittal image showing nasal tip projection; N = nasion; Po = pogonion, TP = 27.7 mm	82
61	Sagittal image showing nasal bone length = 21.6 mm; N = nasion	82
62	Coronal image showing width of nasal aperture = 21.9 mm	83
63	Coronal image showing distance from point corresponding to ANS to a horizontal line joining the most posterior parts of maxillary tuberosity = 45.9mm	83
64	Coronal image showing zygomatic prominence Right side = 6 mm, Left side = 6.6 mm	84
65	Sagittal image showing ABD measured apical to central incisor	84

66	Right maxillary sinus volume = 4.89 cc	85
67	Left maxillary sinus volume = 4.15 cc	85
68	a) Axial image showing cube size adjusted to include the whole premaxilla, b) teeth in premaxilla as viewed from labial aspect	86
69	a) Axial image showing cube size adjusted to include posterior segment on left side and corresponding zoomed cube (b)	87
70	a) Axial image showing cube size adjusted to include posterior segment on right side and corresponding zoomed cube (b)	88
71	Sagittal image showing SNA angle; S = sella; N = nasion; A = point A, SNA= 73.7°	89
72	Coronal image showing ASD; point 0 = most convex point of the deviated nasal septum, ASD = 8.1°	90
73	Sagittal image showing ANB angle; A = point A; N = nasion; B = point B, ANB=2.7°	90
74	Sagittal image showing AUFH; N = nasion; ANS = anterior nasal spine, AUFH = 46.69 mm	91
75	Sagittal image showing ALFH; ANS = anterior nasal spine; Me = menton, ALFH = 65.19 mm	91
76	Sagittal image showing nasal tip projection; N = nasion; Po = pogonion, TP = 21.86 mm	92
77	Sagittal image showing nasal bone length = 20.3 mm; N = nasion	92
78	Coronal image showing nasal aperture = 22.81 mm	93
79	Coronal image showing distance from point corresponding to ANS to a horizontal line joining the most posterior parts of maxillary tuberosity = 35.77mm	93
80	Coronal image showing zygomatic prominence Right side = 7.87 mm, Left side = 9.61mm	94
81	Sagittal image showing ABD measured apical to central incisor	94
82	Right maxillary sinus volume = 8.86 cc	95

83	Left maxillary sinus volume = 8.23 cc	95
84	a) Axial image showing cube size adjusted to include the whole premaxilla, b) teeth in premaxilla as viewed from labial aspect	96
85	a) Axial image showing cube size adjusted to include posterior segment on left side and corresponding zoomed cube (b)	97
86	a) Axial image showing cube size adjusted to include posterior segment on right side and corresponding zoomed cube (b)	98