

بِسْمِ اللّٰهِ الرَّحْمٰنِ الرَّحِیْمِ



HOSSAM MAGHRABY



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



HOSSAM MAGHRABY

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم
قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغييرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HOSSAM MAGHRABY



بعض الوثائق

الأصلية تالفة



HOSSAM MAGHRABY



بالرسالة صفحات

لم ترد بالأصل



HOSSAM MAGHRABY

B1XV9.

**HEALTH EDUCATION PROGRAM FOR
MOTHERS OF CHILDREN SUFFERING FROM
IRON DEFICIENCY ANAEMIA IN
UNITED ARAB EMIRATES**

Thesis

Submitted to the High Institute of Public Health
University of Alexandria
In partial fulfilment of the
Requirements for the

**Degree of Master of Public Health Science
(Health Education and Behavioral Sciences)**

By

Aisha Essa Hassan Abdulla

B. Education and Psychology, University of United Arab Emirates, 1992.
DPH (Health Education and Behavioral Sciences),
High Institute of Public Health,
University of Alexandria, 1999

**High Institute of Public Health
University of Alexandria**

2004

SUPERVISORS



Prof. Dr. Mofida Mahmoud Kamal
Professor of Health Education and Behavioral sciences,
High Institute of public Health
University of Alexandria

Mofida Kamal

Dr. Ebtisam Mohamed Fetohy
Assistant Prof. of Health Education and Behavioral sciences,
High Institute of public Health
University of Alexandria

ع-ب-ف

Dr. Gamil Mohamed Tourkey
Epidemiology Consultants Sharjh Medicine District
Preventive Medicine Department

ع-ب-ت

ACKNOWLEDGMENT

First of all, thanks to **GOD** for help and strength offered to me during this work. I express my deepest gratitude and appreciation to *Prof. Dr. Mofida Mahmoud Kamal*, Professor of Health Education and Behavioral Science, High Institute of Public Health, University of Alexandria for her great willing help, supports keen supervision and encouragement through at the whole period of this study. Prof. Mofida contributed greatly to this work by her vigorous critical suggestions and tremendous experience.

I would like to thank the sincere help, guidance and supervision of *Dr. Ebtisam Mohamed Fetohy*, assistant Professor of Health Education and Behavioral Sciences, High Institute of Public Health, University of Alexandria for her precious advice, constructive guidance, assistance, kindness, valuable support, encouragement and unlimited help which made her supervision a great pleasure to me.

My sincere thanks and appreciation are due to *Dr. Gamil Mohamed Turkey*, Deputy manegar of Sharjah Medicine District, for his help, fatherly supervision and guidance. Indeed, without the facilities Dr. Gamil Mohamed Turkey has given me from his precious time, this work would never have been possible.

I would like to thank the sincere help of Miss **Noha M. Talat**.

Last but not least, I owe special thanks and gratitude to my parents and who nurtured, loved and Faithfully supported me throughout this entire work.

CONTENTS

Chapter	Page
Introduction	1
Aim of the work	37
Subject and Methods	38
Results	52
Discussion	117
Conclusion	129
Recommendations	132
Summary	134
References	139
Appendix	
Protocol	
Arabic summary	

LIST OF TABLES

	Table	Page
I	Distribution of the sample according to the age and sex of the children	53
II	Parent education and occupation of the two studied groups	55
III	Socioeconomic characteristics of the two studied groups.	57
IV	History of iron deficiency of the two studied groups.	59
V	Distribution of the two studied groups regarding the total score of their knowledge before and after the conduction of health education program.	61
VI	Distribution of the studies samples in the two groups regarding to their knowledge about the iron deficiency anaemia symptoms.	66
VII	Distribution of the studied samples in the two groups regarding to their knowledge about the causes of iron deficiency anaemia.	67
VIII	Distribution of the two studies groups according to their perceived susceptibility before and after the conduction of the health education program.	70

	Table	Page
IX	Distribution of the two studies group according to their perceived severity before and after the conduction of the health education program.	74
X	Distribution of the two studies groups according to their perceived benefits before and after the conduction of the health education program	79
XI	The distribution of the two studies groups according the perceived barriers of the mothers of group I and II before and after the program.	84
XII	Distribution of the studies samples in the two groups regarding to their barrier to get proper treatment of IDA.	87
XIII	Distribution of the two studies groups according to their practice before and after the conduction of program.	90
XIV	Distribution of the studies samples in the two groups regarding to their practice toward food intake for group I.	94
XV	Distribution of the studies samples in the two groups regarding to their practice toward breast feeding and when you give first food for your children.	96
XVI	Distribution of the studied samples in the two groups regarding to their practice towards type of food except milk at beginning weaning.	97

	Table	Page
XVII	Distribution of the sample according to their level of knowledge and perceived suseptability in group I after the program.	99
XVIII	Distribution of the sample according to their level of knowledge and perceived severity in group I after the program.	101
XIX	Distribution of the sample according to their level of knowledge and perceived benefits among group I after the program.	103
XX	Distribution of the sample according to their level of knowledge and perceived barriers in group I after the program.	105
XXI	Distribution of the sample according to their scio-economic score and total score of knowledge in group I after the program.	106
XXII	Distribution of the sample according to their scio-economic score and total score of practice in group I after the program.	108
XXIII	Distribution of the studied groups according to anthropometric measurements of the children	110

Table

Page

XXIV Distribution of the studied groups according to the levels of the blood sample before and after conduction of the health education program. 113

XXV Distribution of the two studied groups according to blood picture before and after conduction of the health education program. 116

LIST OF FIGURES

	Figure	Page
I	Distribution of the two studies group according to the total score of their knowledge about iron deficiency anaemia before program.	62
II	Distribution of the two studies groups as regard to their knowledge after program	63
III	Perceived susceptability of group I and II before program.	71
IV	Perceived susceptability of group I and II after program.	72
V	Perceived severity of group I and II before the program.	75
VI	Perceived severity of group I and II after the program.	76
VII	Distribution of the two studied groups according to their perceived benefit of the treatment of IDA before the conduction of HE program.	80
VIII	Distribution of the two studied groups according to their perceived benefits after the conduction of the health education program	81
IX	Perceived barrier between group I and II after the program	85

Figure**Page**

X	Perceived barrier between group I and II after the program.	86
XI	Practices of group I and II before the program.	91
XII	Practices of group I and II after the program.	92