

Dept. of Microbiology



## Effect of Bacterial Proteolytic enzymes on Virulence and Pathogenicity of Avian Influenza Virus

#### A Thesis Presented By Sherein Galal Mohamed Khoulosy

(M.V.Sc, Cairo University)
For PhD Degree in Veterinary Medical Sciences, Microbiology
(Bacteriology, Immunology and Mycology)

Under the Supervision of

#### Prof. Dr. Salah El-Dein Abdel Kareim Selim

Professor of Microbiology Faculty of Veterinary Medicine Cairo University

Prof. Dr. Ahmed S. Shehata
Professor of Microbiology Faculty
of Veterinary Medicine
Cairo University

Prof. Dr. Soad A.A. Abdel Wanis Chief Researcher of Poultry Diseases Animal Health Research Institute

# 

## " وقل رب زوني علما "



(سورة طه – اية ١١٤)

Cairo University
Faculty of Veterinary Medicine
Department of Microbiology

Name: Sherein Galal Mohamed Khoulosy

<u>Degree:</u> PhD Degree in Veterinary Medical Sciences, Microbiology (Bacteriology, Immunology and Mycology)

<u>Title of Thesis</u>: Effect of Bacterial Proteolytic enzymes on Virulence and Pathogenicity of Avian Influenza Virus

#### **Supervisors:**

**Prof. Dr. Salah El-Dein Abdel Kareim Selim , Professor of Microbiology, Dept.** of Microbiology, Faculty of Veterinary Medicine, Cairo University

Prof. Dr. Ahmed Samir Shehata

Professor of Microbiology, Faculty of Veterinary Medicine, Cairo University

Prof. Dr. Soad Abdel Aziz Abdel Wanis

Chief Researcher of Poultry Diseases, Animal Health research Institute

#### **Abstract**

Two thousands cloacal swabs were collected from 100 poultry farms showing respiratory and diarrheic sings. All samples were examined bacteriologically for isolation of *E.coli & Salmonella*. The results revealed detection of *Salmonella* in 31 farms out of 100 examined farms with percentage 31%. Isolation of *E.coli* was carried out and the results revealed isolation of *E.coli* from 67 farms with percentage 67% of examined 100 poultry farms. Detection of the proteolytic enzymes produced by bacterial isolates was done using Protease Agar Casein Assay. All collected samples were examined for detection of Avian Influenza viruses using RRT–PCR. HPAI (H5N1) virus was detected in 10 farms while LPAI (H9N2) was detected in 31 farms out of 100 examined farms. Effect of co-infection with HPAI & LPAI and proteolytic bacteria (*Salmonella Enteritidis* and *E.coli* O158) was carried out in SPF chicken. Morbidity and mortality were recorded as well as re-isolation and virus shedding were done

## Dedication

Formy
Father

Mother

Brothers (Ahmed and Mustafa)

Sister (Azza)

My Friends

I hope that they will be happy, with my deep thanks for their support during the work.

## **Acknowledgment**

First and always, my sincere gratitude and thanks for **[ALLAH]** for all his gifts allover my life, his guidance and care made this study came to light

I would like to express my deep gratitude and appreciation to **Prof. Dr. Salah El-Dein Abdel Kareim Selim**, Professor of Microbiology, Dept. of Microbiology, Faculty of Veterinary Medicine, Cairo University for his valuable help, sincere advice and great support to produce this research

I would like to express my thanks to **Prof. Dr. Ahmed Samir,** Professor of Microbiology, Dept. of Microbiology, Faculty of Veterinary Medicine, Cairo University for his assistance, advice to produce this research

I would like to express my appreciation to **Prof. Dr. Soad Abdel Aziz** Technical Manager for Bacteriology, Reference
Laboratory for Veterinary Quality Control on Poultry Production,
for her assistance, sincere advice and great support during my
research

I would like to thank **Prof. Dr Mona Mehrez**, **Vice Minister of Agriculture**, **Prof. Dr. Momtaz Shahin**, **Director of Animal Health Research Institute** for their support, all the members of Bacteriology Dept. Biotechnology Dept., Animal House Unit with special reference to **Dr. Walid Hamdy Kilany**, Head of Animal House Unit, **Dr. Ahmed Shawky**, staff of Animal House Unit, all staff of epidemiology Unit and also all the members of National Laboratory for Veterinary Quality Control on Poultry Production with reference to **Dr. Abdullah Selim**, Technical Manager for Virology for their help, efforts and continuous assistance.

## **List of Contents**

| Title  | Page       |
|--|------------|
| 1- Introduction  | 1-6        |
| 2-Review of literature   | 7-47       |
| 2-2 Viral –Bacterial association infections  | 7          |
| 2-3 Incidence of Salmonella species in poultry farms   | 14         |
| 2-4 Incidence of <i>E.coli</i> species in poultry farms  | 27         |
| 2-4 Detection of the proteolytic enzymes produced by bacterial isolates using Protease Agar Casein Assay | 34         |
| 2-5 Incidence of Avian Influenza viruses in poultry farms  | 37         |
| 2-6 Cytopathic effect of LPAI virus in cell culture in the   | 45         |
| presence of bacterial protease   |            |
| 3- Material and Methods  | 48-99      |
| 3-1 Material   | 48         |
| 3-1-1 Samples  | 48         |
| 3-1-2 Media used for isolation of Salmonella spp.  | 49         |
| 3.1.3. Media used for biochemical identification of <i>Salmonella</i> spp.                               | 50         |
| 3.1.4. Chemicals and reagents used for biochemical   | 51         |
| identification of Salmonellae.   | <i>E</i> 1 |
| 3.1.5. Stain used  | 51         |
| 3.1.6. Diagnostic antisera   | 51<br>51   |
| 3.1.7. Equipment used for the isolation, identification and detection of <i>Salmonella</i>               |            |
| 3-1-8 Materials used for PCR technique   | 52         |
| 3-1-9 Equipment and apparatus used for PCR Technique   | 56         |
| 3-1-10 Standard bacterial strains  | 57<br>57   |
| 3-1-11. Media used for isolation of <i>E.coli</i> spp.   | 57<br>59   |
| 3-1-12.Media used for biochemical identification of <i>E.coli</i>  | 58         |
| 3-1-13. Diagnostic antisera  | 59         |
| 3-1-14. Materials and equipment used for PCR   | 60         |
| amplification of <i>E.coli</i> isolates  | 65         |
| 3-1-15 Media used for detection of the proteolytic enzymes   | 65         |
| produced by bacterial isolates   | ((         |
| 3-1-16 Material used for detection of Avian Influenza virus  | 66         |
| by Real Time PCR   | 70         |
| 3-1-17. Cytopathic effect of Low Pathogenic Avian  | 70         |
| influenza in cell culture in the presence of bacterial   |            |
| proteolytic enzymes  |            |

| Title   | Page           |
|---|----------------|
| 3-1-18. Materials used to study the effect of proteolytic   | 72             |
| bacteria on pathogenicity of AI viruses (H5N1 &   |                |
| H9N2)   | <b>-</b> 4 00  |
| 3-2 Methods   | 74-99          |
| 3-2-1 Handling and preparation of collected cloacal swabs   | 74             |
| for isolation of Salmonella Spp.  | 7.4            |
| 3-2-2 Procedure for detection of Salmonella spp.  | 74<br>75       |
| 3-2-3 Selection of colonies for confirmation of <i>Salmonella</i>                                 | 75<br>75       |
| 3-2-4 Microscopic identification of Salmonella  | 75<br>76       |
| 3-2-5 Biochemical identification of <i>Salmonella spp</i> .                                       | 76<br>76       |
| 3-2-6 Serotyping of <i>Salmonella</i> spp. 3-2-7 Polymerase Chain Reaction (PCR) for detection of | 78<br>78       |
| Salmonella  | 78             |
| 3-2-8 Isolation and identification of <i>E.coli</i>   | 82             |
| 3-2-8-1. Procedure for detection of <i>E.coli</i>   | 82             |
| 3.2.8.2. Microscopical examination  | 83             |
| 3.2.8.3. Biochemical identification of <i>E.coli</i> isolates:                                    | 83             |
| 3.2.8.4. Serotyping of <i>E.coli</i> isolates   | 85             |
| 3.2.8.5 Polymerase Chain Reaction (PCR) for <i>E.coli</i>   | 86             |
| identification  |                |
| 3-2-9 Protease Agar Casein Assay used for detection of the  | 89             |
| proteolytic enzymes produced by bacterial isolates  |                |
| 3-2-10 Method used for detection of Avian Influenza virus   | 90             |
| by REAL TIME PCR  |                |
| 3-2-11. Cytopathic effect of Low Pathogenic Avian   | 95             |
| influenza in cell culture in the presence of bacterial  |                |
| proteolytic enzymes   |                |
| 3-2-12. Procedures for experimental infection of chicken to                                       | 97             |
| study the effect of proteolytic bacteria on   |                |
| pathogenicity of AI viruses (H5N1 & H9N2)   |                |
| 4- Results  | 100-127        |
| 5- Discussion   | 128-144        |
| 6- Summary  | 145-147        |
| 7- Conclusion   | 148            |
| 8- References   | 149-177<br>۳-۱ |
| Arabic summary  | 1 – 1          |
| Arabic abstract   |                |

## **List of Tables**

| No         | Title   | Page       |
|------------|---|------------|
| 1          | Number of examined chicken farms from 7   |            |
|            | governorates  | 49         |
| 2          | Oligonucleotide Primers sequences used for  |            |
|            | amplification of DNA for the detection of Salmonella  |            |
|            | species   | 53         |
| 3          | Reference bacterial strains cultures used for quality   |            |
| 4          | assurance and PCR specificity   | 57         |
| 4          | Monovalent specific antisera used in the serotyping of  | <b>5</b> 0 |
| 5          | E.coli serovars   | 59         |
| 5          | Oligonucleotide primers sequences used for  | 62         |
| 6          | amplification of DNA for the detection of <i>E.coli</i> The oligonucleotide for amplification of part of M gene | 02         |
| U          | for AIV by REAL TIME PCR  | 68         |
| 7          | The oligonucleotide for amplification of part of H5 gene  | 00         |
| ,          | for AIV-H5 by REAL TIME PCR   | 68         |
| 8          | The oligonucleotide for amplification of part of H9 gene  | 00         |
|            | for AIV-H9 by REAL TIME PCR   | 69         |
| 9          | Biochemical confirmation of Salmonella spp.   | 76         |
| 10         | Thermal cycler steps of Salmonella DNA amplifications   | 80         |
| 11         | Biochemical character of <i>E.coli</i> isolates   | 85         |
| 12         | Thermal cycler steps of <i>E.coli</i> DNA amplifications  | 88         |
| 13         | Real-Time PCR Reaction Mix Volumes  | 91         |
| 14         | Thermal cycling conditions for gene-specific Probe and  |            |
|            | Primer sets for influenza virus Type A  | 92         |
| 15         | Thermal cycling conditions for gene-specific Probe and  | 0.0        |
| 1.6        | Primer sets for influenza virus H5  | 92         |
| 16         | Thermal cycling conditions for gene-specific Probe and  | 0.2        |
| 17         | Primer sets for influenza virus H9  | 93         |
| 17         | Experiment groups and challenge doses   | 98<br>99   |
| 18<br>19   | Experiment Sampling  Providence of Salmonella among broiler forms   | 99<br>100  |
| 20         | Prevalence of <i>Salmonella</i> among broiler farms Prevalence of <i>Salmonella</i> among layer farms           | 100        |
| 21         | Total prevalence of <i>Salmonella</i> in examined chicken   | 101        |
| <i>4</i> 1 | farms   | 101        |
| 22         | Antigenic structure of isolated <i>Salmonella</i> Serovars  | 103        |

| No | Title  | Page |
|----|--|------|
| 23 | Percentage of Salmonella serovars isolated from cloacal      |      |
|    | samples collected from poultry flocks                        | 104  |
| 24 | Most prominent signs appeared in poultry flocks              |      |
|    | infected with Salmonella spp.                                | 105  |
| 25 | Prevalence of <i>E.coli</i> among broiler farms              | 108  |
| 26 | Prevalence of <i>E.coli</i> among layer farms                | 108  |
| 27 | Total prevalence of <i>E.coli</i> in examined chicken farms  | 110  |
| 28 | Percentage of <i>E.coli</i> serogroups isolated from poultry |      |
|    | flocks   | 110  |
| 29 | Serogrouping of <i>E.coli</i> isolates                       | 111  |
| 30 | Most prominent signs appeared in poultry flocks              |      |
|    | infected with E.coli serogroups                              | 112  |
| 31 | Results of detection of stx1 gene of E.coli isolates         |      |
|    | recovered from chickens                                      | 113  |
| 32 | Results of detection of eaeA gene of E.coli isolates         |      |
|    | recovered from chickens                                      | 114  |
| 33 | Results of detection of iss gene of E.coli isolates          |      |
|    | recovered from chickens                                      | 116  |
| 34 | PCR results for detection of viral infection in samples      |      |
|    | from Broiler farms   | 119  |
| 35 | PCR results for detection of viral infection in samples      |      |
|    | from Layer farms   | 120  |
| 36 | Total prevalence of AI viruses in examined poultry           |      |
|    | farms  | 120  |
| 37 | The daily mortalities of LPAI H9N2 challenged groups         | 123  |
| 38 | The daily mortalities of HPAI H5N1 challenged groups         | 124  |
| 39 | Specific cumulative mortalities in all groups challenged     |      |
|    | with A1 viruses  | 125  |
| 40 | Overall percentage of re-isolation of <i>E.coli</i> and      |      |
|    | Salmonella from different organs of SPF chicken,             |      |
|    | necropsied on day 6, 8 and 10th day post-viral               |      |
|    | infection  | 126  |
| 41 | Mean of virus shedding in samples collected at 6th, 8th      |      |
|    | and 10th day post viral inoculation                          | 127  |

## **List of Figures**

| No | Title   | Page |
|----|---|------|
| 1  | Prevalence of Salmonella among examined commercial  |      |
| 1  | chicken farms   | 102  |
| 2  | Percentage of Salmonella serovars isolated from poultry   |      |
| _  | flocks  | 104  |
|    | Agarose gel electrophoresis showing positive  |      |
| 3  | amplification of 284 bp fragments using 139 – 141   |      |
|    | primers specific for the <i>invA</i> gene from extracted DNA of <i>Salmonella Enteritidis</i>                 | 106  |
|    | Agarose gel electrophoresis showing positive  | 100  |
|    | amplification of 488 bp fragments using A058 - A01  |      |
| 4  | primers specific for the sefA gene found in Salmonella  |      |
|    | Enteritidis from extracted DNA of one standard strains  |      |
|    | of Salmonella Enteritidis   | 107  |
| 5  | Prevalence of <i>E.coli</i> among examined commercial   |      |
| 3  | chicken farms   | 109  |
| 6  | Percentage of <i>E.coli</i> serogroups isolated from poultry  | 111  |
|    | farms   | 111  |
|    | Agarose gel electrophoresis showing amplification of 614 bp fragment using PCR, performed with primer         |      |
| 7  | specific for <i>stx</i> 1genegene of <i>E.coli</i> isolated from  |      |
|    | chickens.   | 113  |
|    | Agarose gel electrophoresis showing amplification of  | 115  |
| 8  | 890 bp fragment using PCR, performed with primer  |      |
|    | specific for eae A gene of E.coli isolated from chickens  | 115  |
|    | Agarose gel electrophoresis showing positive  |      |
| 9  | amplification of 266 bp fragment using PCR, performed   |      |
|    | with primer specific for iss gene of E.coli isolated from   | 116  |
| 10 | chickens.   | 116  |
| 10 | Agar Casein Test protease positive bacteria isolates<br>Cytopathic effect of the Avian Influenza virus (H9N2) | 117  |
| 11 | with proteolytic enzymes on MDCK cells (20X)  | 121  |
|    | with proteorytic chizymes on MDCR cens (20A)  | 141  |