

سامية محمد مصطفى



شبكة المعلومات الجامعية

بسم الله الرحمن الرحيم



سامية محمد مصطفى



شبكة المعلومات الجامعية



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



سامية محمد مصطفى



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



سامية محمد مصطفى



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



سامية محمد مصطفى



شبكة المعلومات الجامعية



بالرسالة صفحات لم ترد بالأصل



MICROBIOLOGICAL STUDIES ON SOME MEDICAL PLANTS

A Thesis Submitted to

Dept. of Nutrition & Food Science

Home Economic Faculty of Minufiya University

For Degree of Master

In Nutrition & Food Science

By


Ashraf Abo El-Azem Abed El-Fatah Abo El-Azem

Supervised By

Prof. Dr. Mohamed Fahmi Sadiq

Professor of Food Hygiene, Nutrition


Institute, Cairo.



Prof. Dr. Mohamed Reda Gomma

Professor of Food Microbiology, Nutrition


Institute, Cairo.



Dr. Sherif Sabry Ragab

Associate Professor of Nutrition & Food Science,

Faculty of Home Economic, Minufiya University.



1999

B
14914

APPROVAL SHEET

Name

Ashraf Abo El-Azem Abed El-Fatah Abo El-Azem

Title of Thesis

Microbiological Studies on Some Medical Plants

Degree

Master in Nutrition and Food Science

This Thesis has been approved by:

Prof. Dr.

Samira Lancel

Prof. Dr.

Fahs

Prof. Dr.

M.M. El-Sayel

Date: 12/6/1999

CONTENTS

	Page
1. INTRODUCTION AND AIM OF WORK	1
<i>1.1. Introduction</i>	<i>1</i>
<i>1.2. Aim of work</i>	<i>2</i>
2. REVIEW OF LITERATURE	3
<i>2.1. Definitions</i>	<i>3</i>
<i>2.2. Microbiolgoical contamination</i>	<i>3</i>
<i>2.2.1. Aerobic plate count</i>	<i>3</i>
<i>2.2.2. Coliform organisms</i>	<i>4</i>
<i>2.2.3. Mold and yeast</i>	<i>5</i>
<i>2.3. Antibiotic potency</i>	<i>8</i>
<i>2.4. Effect of the chemical composition of the species and herpes for the activity organism</i>	<i>21</i>
3. MATERIALS AND METHODS	24
<i>3.1. Materials</i>	<i>24</i>
<i>3.2. Methods</i>	<i>28</i>
4. RESULTS AND DISCUSSIONS	34
<i>4.1. Microbiological contamination</i>	<i>34</i>
<i>4.2. Antibiotic potancy</i>	<i>95</i>
5. SUMMARY	139
6. REFERENCES	144

ARABIC SUMMARY

ACKNOWLEDGMENT

I wish to express my hearty appreciation to Prof. Dr. Mohamed Fahmi Sadiek, Professor of Food hygiene-Nutrition Institute, Cairo for his valuable advice and easement of the practical section of this study in addition to his supportive action for me.

My sincere gratitude and due to Prof. Dr. Mohamed Reda Gomma, Professor of Food Microbiology, Nutrition Institute, Cairo for his interest indispensable advice, valuable comments and criticism in addition to his continous follow up of this research.

I would like to thank Dr. Sherif Sabry Ragab, Associate Professor of Nutrition and Food Science, Home Economic Faculty, Menofiya University for his helpful action and criticism.

Finally, I am also very thankful to my Parents, my Wife, my Daughter and my Brothers for their patience, love and encouragement, they gave me during my study.

1. INTRODUCTION AND AIM OF WORK

1. INTRODUCTION AND AIM OF WORK

1.1. Introduction:

Spices and condiments are consumed in large quantities in Egypt and many countries in the world since very ancient times.

Spices were used as embalming materials in ancient Egypt, as sweet scents in religious ceremonies and to mask the putrid flavour of meat in hot climates lacking refrigeration.

A few spices have an antimicrobial effect at the concentrations used in food and thus serve as preservatives.

In Egypt, most of the aromatic plants are cultivated to produce some of their extracts, i.e., volatile oils, concretes, absolutes, etc.

Many volatile oils are now being processed or extracted for industrial uses. Their wide use in perfumes and cosmetics and for the scenting of soap and similar products. These oils are also used for masking objectionable odors inherent or acquired during industrial processing such as detergents and insecticides. Also, different medical preparations containing aromatic plants extracts are now produced at mass production scale.

Spices are interest to microbiologists for four principal reasons:

1. They contain large numbers of microorganisms that occasionally may cause spoilage or disease when introduced into food.
2. They exhibit antimicrobial activity and occasionally aid preservation and
3. They stimulate microbial metabolism (I.C.M.S.F., 1980).

1.2. Aim of work:

This study was conducted to evaluate the microbial contamination as well as antibiotic potency for different aqueous extract of selected Medical plants.

These plants were obtained from different areas in Cairo and Giza.

Thus the overall objects of the present work were to:

1. Evaluate microbiological contamination of plants through determination of:
 - Aerobic colony count.
 - Molds and yeasts count.
 - Most probable number for coliform organisms.
 - Presence of *E. coli* as an indicator for faecal contamination.
2. Assess the antibiotic potency for aqueous extract of selected plants (Anise, Comomela, Caraway, Cinnamon, Fennel, Hibiscus, Liquorice, Peppermint, Sweet Basil and Thyme) on *Stap aureus*, *Bacillus cereus*, *Bacillus subtilis*, *Proteus mirabilis*, *Salmonella typhimurium* and *E. coli*.

2. REVIEW OF LITERATURE

2. REVIEW OF LITERATURE

2.1. Definitions:

Herbs are the soft-stemmed plant materials used in seasoning food. All other aromatic plant producing products used for a similar purpose are called spices, although this broad definition admits of several exceptions.

Spices are usually only parts of plants and may be either roots, rhizomes, barks, seeds, fruits, flower buds, or others. Unlike the herbs, the spices are very aromatic and may contain large percentages of essential oils as well as other powerful nonvolatile flavouring components. They are normally derived from the semi tropical or tropical regions of the world, are harvested, and usually sun-dried to form the spice of commerce.

Condiments are seasonings which are added to food after it has been served. In this category the most popular and widely used are salt, mustard, pepper, and ginger.

Those definitions previously mentioned are according to **Heath *et al.* (1978)**.

2.2. Microbiological contamination:

2.2.1. Aerobic plate count:

Heath (1964) said that the source of such pathogens can be soil, fecal material from birds and other animals and non potable water used for soaking some spices.

A wide variety of nonsporing bacteria may also be present (Elmossalimi and Youssif, 1965).

Goto *et al.* (1971) found that most of the aerobic plate count colonies are *Bacillus* spp. example, *B. subtilis*, *B. megaterium*, *B. pumilus*, *B. firmis* and *B. brevis*.

Some spices may contain spores from mesophilic aerobes, mesophilic anaerobes and flat sour thermophiles. Some spices sometimes contain bacteria that can cause food-born infections (ICMSF, 1980).

Khairy *et al.* (1992) found that total bacterial count in anise was 1.43×10^2 , in caraway 3.40×10^2 and in fennel 78.50×10^2 / gm.

Mimica *et al.* (1993) reported that Chamomile (*chamomilla recutita*) is widely used as a herbal tea and the therapeutic value depends mainly on the essential oil content. Commercial chamomile tea was sampled at monthly intervals for 18 months. The main oil constituent was bisabolil oxide A which varied between 33.46 and 48.48%. The total bacteria count ranged from 4.0×10^4 to 4.6×10^6 /g DM and yeasts from 3.0×10^3 to 5.8×10^5 / g DM.

2.2.2. Coliform organisms:

Kadis *et al.* (1971) reported that coliform are often found in spices, but *E. coli* is infrequent.