

بسم الله الرحمن الرحيم



-Cardon - Cardon - Ca





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار







بعض الوثائق

الأصلية تالفة







بالرسالة صفحات

لم ترد بالأصل



PREVALENCE OF COMMON PARASITIC INFECTIONS AMONG HEPATIC PATIENTS

B18606

THESIS SUBMITTED IN PARTIAL FULFILLMENT OF M.Sc. DEGREE IN MEDICAL PARASITOLOGY

By Yousry Abd El-Hameid Hawash

(M.B., B.Ch.)

Demonstrator of Parasitology,

National Liver Institute, Menoufiya University

Supervised by

Prof. Dr. SORAYA ABD EL-AZIZ SHARAF

Chairman – Parasitology Dept., National Liver Institute, Menoufiya University

アンソ

Dr. OMAYMA KAMEL EL-SHAFEI

Assistant Professor of Parasitology, Faculty of Medicine, Menoufiya University


الممترثهس

Dr. ABD EL-GAWAD EL-SAYED A.

Lecturer of Parasitology,
National Liver Institute, Menoufiya University

FACULTY OF MEDICINE MENOUFIYA UNIVERSITY 2002

~~. ~~**~**




ڰڹڿڮٵٚٳڲڹڷڰڷٳڐٵؿٳڲڹڮٳڮڮ ٳٷڸٷٳٳٷٳڲٷڮٳٳڲڮڮٷٳ ٳٷڸٷٳٷٳٷٳٷٳٷٳٷٳٷٳٷ

البقرة - ٣٢



To my Beloved Family



ACKNOWLEDGEMENT

I would like to introduce my heartly appreciation to *Prof.*Dr. Nashaat El-Sayed Nassef, Professor and Head of Parasitology Department, Faculty of Medicine, Menoufiya University, for her kind moral support, continuous encouragement and sincere help.

I would like to express my deepest gratitude as I am deeply indebted to my professor and supervisor, *Prof. Dr. Soraya Abd El-Aziz Sharaf*, Professor and Head of Parasitology Department, The National Liver Institute, Menoufiya University, for her kind moral support, enthusiastic encouragement and help to pursue this work.

I would also like to express my very sincere thanks to *Dr. Omaima Kamel El-Shafei*, Assistant Professor of Parasitology, Faculty of Medicine, Menoufiya University, for her patience, tremendous effort and guidance throughout this work.

I wish to express my deep indebtedness to *Dr. Abd El-Gawad El-Sied Abd El-Gawad*, Lecturer of Parasitology, The National Liver Institute, Menoufiya University, for his close supervision, encouragement and sincere follow up.

I am deeply indebted to *Dr. El-Amir Mahmoud Amir*, Lecturer of Parasitology, The National Liver Institute, Menoufiya University, for his sustained support, continuous supervision and unlimited help.

. •

CONTENTS

Introduction	1
Aim of the work	5
Review of Literatures	6
SCHISTOSOMIASIS	6
History	6
Classification	7
Geographical distribution and epidemiology	9
Immunopathology	11
Prevention and control	13
<i>FASCIOLIASIS</i>	15
Classification and life cycle	15
Geographical distribution and epidemiology	16
Pathology of fascioliasis	20
Diagnosis of Fasciola	23
ENTAMOEBA HISTOLYTICA	32
Epidemiology	32
Life cycle	33
Clinical aspects	33
Treatment of amoebiasis	39
ECHINOCOCCUS GRANULOSUS	41
Historical notes	41
Geographical distribution and epidemiology	41
Morphology and life cycle	43
Pathology and pathogenesis	46
Diagnosis of hydatid disease	47
PATIENTS AND METHODS	54
RESULTS	.68
DISCUSSION	78
SUMMARY & CONCLUSION	84
RECOMMENDATIONS	87
REFERENCES	88
Arabic summary	

LIST OF TABLES

- Table (1): Characteristics of studied hepatic disease patients and controls
- Table (2): Clinical data of studied hepatic disease patients.
- Table (3): Laboratory data of studied hepatic patients and controls
- Table (4): Abdominal ultrasonography of hepatic patients.
- Table (5): Abdominal CT findings for selected hepatic patients.
- Table (6): Results of stool analysis by formol-ether oncentration method in hepatic disease patients and controls.
- Table (7): Comparison of the results of IHA method for detection of schistosomiasis in studied hepatic disease patients and controls.
- Table (8): Result of stool analysis for *Schistosoma mansoni* by Kato technique in hepatic disease patients.
- Table (9): Results of rectal snip analysis for S. mansoni in hepatic disease patients.
- Table (10): Results of analysis for detection of *S. mansoni* by different techniques in hepatic disease patients.
- Table (11): Comparison of the results of IHA for detection of fascioliasis between hepatic disease patients and controls.
- Table (12): Comparison of the prevalence of amoebiasis as detected by IHA between hepatic disease patients and controls.
- Table (13): Comparison of the results of detection of hydatid disease by (IHA) technique between hepatic disease patients and controls.