

سامية محمد مصطفى



شبكة المعلومات الجامعية

# بسم الله الرحمن الرحيم



سامية محمد مصطفى



شبكة المعلومات الجامعية



# شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





سامية محمد مصطفى



شبكة المعلومات الجامعية

# جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم


## قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها  
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



## يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



سامية محمد مصطفى



شبكة المعلومات الجامعية



# بعض الوثائق الأصلية تالفة





سامية محمد مصطفى



شبكة المعلومات الجامعية



# بالرسالة صفحات لم ترد بالأصل



GEOLOGICAL AND GEOCHEMICAL STUDIES ON  
THE PLIOCENE-QUATERNARY SEDIMENTS OF  
EL-HAMRAWIN-MERSA-ALAM DISTRICT, RED  
SEA COAST, EGYPT

*A Thesis*

Submitted By

AMAN ABD EL-HAKEEM ZALAMAH

For The Degree Of  
DOCTOR OF PHILOSOPHY IN SCIENCE  
(GEOLOGY)

To :

*The Geology Department  
Faculty of Science  
Zagazig University*

Zagazig-Egypt

1994

B

17951

## ACKNOWLEDGEMENT

The author wishes to express the deep appreciation to Prof. Dr. **M. A. Shaaban** Head of Geology Department, Zagazig University for supervision, kind support, encouragement and consultation. And also I wish to express my deep thanks and gratitude to Ass. Prof. Dr. **A.E. Ziko** Geology Department Faculty of Science, Zagazig University for supervising, sincere help and valuable discussions.

The author is greatly indebted to Prof. Dr. **M. M. Hassaan**, Prof of Economic Geology and Geochemistry, Geology Department, Al-Azhar University who proposed the point of research, planned and supervised all steps of this research, samples collection, constant guidance, critical reading as well as the keen revision of the manuscript and consultation.

Many thanks are extended to Prof. Dr. **M. Y. Attawiya** Head of Radioactive Minerals Department, Nuclear Materials Corporation, Cairo, Egypt for supervising and funding facilities of chemical analysis and X-ray diffraction analysis.

Special thanks due to Dr. **F. M. Abu El-Enin**, Geology Department, Faculty of Science, Zagazig University for Sincere help during collection of samples, supervising and critical reading of the manuscript.

Thanks are also due to Dr. **H. A. Hamed**, Geology Department, Zagazig University for his help. Thanks and deep gratitude to My Husband for his encouragement, his help and association to me in the field.

I am very grateful to all who assisted me and made this work come true.

# CONTENTS

	Page
<b>TABLE OF CONTENTS</b>	i
<b>LIST OF FIGURES</b>	iii
<b>LIST OF TABLES</b>	xii
 <b>CHAPTER I: INTRODUCTION AND PREVIOUS WORK</b>	 1
<b>CHAPTER II: LITHOSTRATIGRAPHY</b>	19
Wadi Quseir El-Kadium	25
Wadi Essel	32
Wadi Sherm El-Bahari	36
Wadi Um-Gheig	39
Wadi Abu-Dabbab	43
Ras Shagra Area	47
 <b>CHAPTER III: PETROLOGY</b>	 53
<b>GRAIN SIZE ANALYSIS</b>	53
Grain Size Parameters	63
Interrelationship Between Grain Size Parameters	76
Paleoenvironmental Interpretations	80
<b>PETROGRAPHY</b>	101
A- Clastic Group of Gabir Formation	102
B- Carbonate Group of Gabir Formation	114
C- Clastic Group of Shagra Formation	118
D- Carbonate Group of Shagra Formation	126
E- Raised Reefs of Abu Shegeili Formation	139
<b>DIAGENESIS</b>	147
1- Diagenesis in Sandstones	147
2- Diagenesis in Carbonate	152



<b>CHAPTER IV: MINERALOGICAL COMPOSITION</b>	173
A- HEAVY MINERALS ANALYSIS	173
Light Fraction	174
Heavy Fraction	175
Genetic Significance of Certain Heavy Minerals	185
B- X-RAY DIFFRACTION ANALYSIS	190
1- Mineral Composition of Bulk Samples	190
2- Clay Minerals	202
3- X-ray Diffraction Analysis of Fossils and Recent Equivalents	205
 <b>CHAPTER V: GEOCHEMISTRY</b>	 213
A- GEOCHEMISTRY OF MAJOR ELEMENTS	214
Quseir El-Kadium Locality	215
Essel Locality	219
Sherm El-Bahari Locality	223
Um-Gheig Locality	225
Abu-Dabbab Locality	226
Ras-Shagra Locality	228
B- GEOCHEMISTRY OF TRACE ELEMENTS	232
1- Abundance	233
2- Vertical Distribution	
Quseir El-Kadium Locality	252
Essel Locality	252
Sherm El-Bahari Locality	252
Um-Gheig Locality	252
Abu-Dabbab Locality	253
Ras-Shagra Locality	253
 <b>CHAPTER VI: DEPOSITIONAL ENVIRONMENT</b>	 254
<b>SUMMARY AND CONCLUSION</b>	267
<b>REFERENCES</b>	279

# LIST OF FIGURES

Figure	Page
(1): Geological map of the studied area	2
(2): Correlation chart of the studied sections	20
(3): Lithologic section of Wadi Quseir El-Kadium	26
(4): Showing the contact between silty sandstone bed (micaceous, bed 11), and fossiliferous sandy limestone (bed 12, dolomitic), Gabir Formation, Wadi Quseir El-Kadium	28
(5): Raised reefs, corals and spines, Raised Reefs, Wadi Quseir El-Kadium	28
(6): The First and the Second Raised Reefs, Raised reef, Abu Shegeili Formation Wadi Quseir El-Kadium	29
(7): Panoramic view of Wadi Quseir El-Kadium	30
(8): Lithologic section of Wadi Essel	33
(9): Shell accumulations and coralline limestones bed no. 20, Raised reefs of Abu Shegeili Formation, Wadi Essel	34
(10): Siltstone at the base, graded to cyclic conglomerate, topped by coralline limestone (bed no. 16), Raised reefs, Abu Shegeili Formation, Wadi Sherm El-Bahari	34
(11): Lithologic section of Wadi Sherm El-Bahari	37
(12): Lithologic section of Wadi Um-Gheig	40
(13): Interance of Wadi Um-Gheig	42
(14): Coralline limestone at the interance of Wadi Abu-Dabbab. Raised Reef of Abu Shegeili Formation	42
(15): Lithologic section of Wadi Abu-Dabbab	44
(16): The conglomeratic bed at the interance of Wadi Abu-Dabbab, Abu Shegeili Formation	46
(17): Lithologic section of Ras-Shagra area	48
(18): Apart of panoramic view of Ras Shagra	51
(19): A second part of panoramic view of Ras Shagra	52
(20): Cumulative curves of the studied sand samples of Pliocene Wadi Essel Locality	55
(21): Cumulative curves of the studied sand samples of Pliocene Wadi Essel locality	56

(22): Cumulative curves of the studied sand samples of Pliocene Wadi Sherm El-Bahari locality	57
(23): Cumulative curves of the studied sand samples of Pliocene Wadi Um-Gheig locality	58
(24): Cumulative curves of the studied sand samples of Pliocene Wadi Abu-Dabbab locality	59
(25): Cumulative curves of the studied sand samples of Pliocene Wadi Abu-Dabbab locality	60
(26): Cumulative curves of the studied sand samples of Pliocene Wadi Abu-Dabbab locality	61
(27): Cumulative curves of the studied sand samples of Pliocene Ras Shagra locality	62
(28): The scatter diagram of mean size (MZ) versus each of sorting ( $\sigma_1$ ), Skewness (SKI) and Kurtosis (KG) of the studied sandstones (after <b>Mason and Folk, 1958</b> )	78
(29): Average Scatter diagram of sorting ( $\sigma_1$ ) versus each of skewness (SKI) and kurtosis (KG) of the studied sandstones, <b>Mason and Folk (1958)</b>	79
(30): Scatter diagram of skewness (SKI) versus kurtosis (KG) of the studied sandstones. <b>Mason and Folk (1958)</b>	81
(31): Apolt of inclusive graphic skewness (SKI) versus graphic kurtosis (KG) of the studied sandstones. <b>Mason and Folk (1958)</b>	81
(32): Bivariant plots of Mean-Size versus standard deviation, (Dune and River plot), after <b>Moiola and Weiser (1968)</b>	83
(33): Bivariant plots of mean-size versus standard deviation (Beach and River plot), after <b>Moiola and Weiser (1968)</b>	83
(34): Bivariant plot of Mean size versus inclusive graphic standard deviation, after <b>Friedman (1961)</b>	83
(35): Bivariant polt of skewness versus standard Deviation after <b>Friedman (1961)</b>	85
(36): Bivariant polt of skewness versus standard Deviation after <b>Friedman (1967)</b>	85
(37): Bivariant polt of first percentile versus standard Deviation after <b>Friedman (1967)</b>	86



(38): Bivariant plot of skewness versus mean size, after <b>Moiola and Weiser (1968)</b>	86
(39): Bivariant plot of skewness versus mean size after <b>Moiola and Weiser (1968)</b>	88
(40): Bivariant plot of skewness versus kurtosis after <b>Moiola and Weiser (1968)</b>	88
(41): Calcareous lithic arenite with pelecypod shell fragments, Gabir Formation, Wadi Essel, SN: 7 pp, x 2.5	103
(42): Calcareous lithic arenite with partly altered feldspar with sparitic cement, Gabir Formation, Wadi Essel SN: 7, CN. x 2.5	103
(43): Calcareous lithic arenite with algae ( <i>lithophyllum. sp</i> ). Gabir Formation, Sherm El-Bahari locality, SN: 2, pp., x 4	105
(44): Calcareous Feldspathic quartz arenite, notice the polycrystalline quartz grains and sparitic cement. Gabir Formation, Wadi Abu-Dabbab, SN: 3, CN., X 2.5	105
(45): The same with plagioclase and clear sparry calcite. CN., X 2.5	106
(46): Calcareous feldspathic quartz arenite with partially sericitized microcline and polycrystalline quartz grains embedded in micrite and microsparite. Gabir Formation, Wadi Abu-Dabbab, SN: 5, CN., X 2.5	106
(47): The same with micritic rims (envelope) around the plagioclase and polycrystalline quartz grains. CN., X 4	107
(48): The same with micritic and microsparitic rim around the grains. Notice the recrystallization of cement. CN., X 4	107
(49): The same with phosphatic particle (well rounded, dark, grain at the upper part right), CN., X 4	108
(50): Gypseous calcareous feldspathic quartz arenite with fine grained quartz and clear gypsum veinlet. Gabir Formation, Wadi Abu-Dabbab, SN: 22, CN., X 4	108
(51): Calcareous Feldspathic quartz arenite with plagioclase and monocrystalline quartz grains embedded in sparite and microsparite. Gabir Formation, Wadi Abu-Dabbab SN: 19, CN., X 2.5	109

- (67): Gypseous arkosic arenite with fibrous gypsum. Shagra Formation, Wadi Essel, SN: 14, CN., X 2.5 124
- (68): Other part of the same section with partially altered feldspar (orthoclase) and monocrystalline quartz grains. SN: 14, pp., X 2.5 124
- (69): Calcareous arkosic arenite with partially altered feldspar. Shagra Formation, Abu-Dabbab, SN: 30, CN., X 4 125
- (70): Other part of the same thin section clear hornblende grains with its clear 2 sets of cleavage. CN., X 4 125
- (71): Calcareous arkose with partially altered feldspar (sericite). Notice the sparite cement. Shagra Formation, Abu-Dabbab locality SN: 33, CN., X 2.5 128
- (72): Sandy fossiliferous grainstone (sandy biosparite) with algae (*Lithophyllum sp.*) and Foraminifera (millioids), Shagra Formation, Wadi Essel, SN: 13, pp., X 2.5 128
- (73): Sandy grainstone with foraminiferal tests and shell fragments. Notice the recrystallization of the shell fragments and the filling of the fossil interspaces with sparitic calcite. Shagra Formation, Wadi Essel, SN: 13, CN., X 2.5 129
- (74): Molluscan biomicrite (Packstone/grainstone), notice the original structure of the Oyster shell fragment and the beginning of the pore filling with drusy calcite at the periphery of the pore spaces, Shagra Formation, Wadi Essel, SN: 18, pp., X 2.5 129
- (75): Sandy coralline grainstone/packstone (coralline algal biomicrite) with large coralline bioclasts and *Lithophyllum Sp.* Shagra Formation, Wadi Essel, SN: 19, pp., X 2.5 130
- (76): Other part of the same section with large bioclasts of *Lithophyllum sp.* pp., X 2.5 130
- (77): Sandy grainstone/packstone (biomicrite) with phosphatic particle (collophane), Shagra Formation, Wadi Essel, SN: 21, pp., X 2.5 131
- (78): Intraclastic grainstone (intra-sparite) with onchoids. Notice the epitaxial and blocky cement Shagra Formation, Wadi Sherm El-Bahari SN: 9, pp., X 2.5 131
- (79): Algal sandy wackestone/packstone (sandy biomicrite) Shagra Formation, Wadi Sherm El-Bahari, SN: 13, pp., X 2.5 133

(80):	Sandy bioclastic (miliolid) wackestone. Shagra Formation, Wadi Sherm El Bahari, SN: 15, pp., x 2.5.	133
(81):	Sandy porous grainstone (caverneous limestone) with caves. Shagra Formation, Wadi Essel, SN: 22, CN., X 2.5	135
(82):	Algal coralline framestone. Shagra Formation, Wadi Sherm El-Bahari, SN: 13, pp., X 2.5	135
(83):	Framestone (coralline). Shagra Formation, Wadi Sherm El-Bahari, SN: 16, pp., X 2.5	136
(84):	Bufflestone. Shagra Formation, Wadi Abu-Dabbab, SN: 35, CN., X 2.5	136
(85):	Algal coralline bindstone, Shagra Formation, Wadi Sherm El-Bahari, SN: 14, pp., X 2.5	138
(86):	Oolitic grainstone/packstone. Shagra Formation Wadi Quseir El-Kadium, SN: Q 8, CN., X 6	138
(87):	Enlarged of the same section X 20	140
(88):	Bufflestone of Raised reefs (Abu Shegeili Formation), Wadi Abu Dabbab, SN: 42', pp., X 2.5	140
(89):	The same with pryozoa and shell fragments, pp., X 2.5	141
(90):	Bufflestone with fine to medium-grained, monocrystalline quartz grains. Pleistocene Raised reefs of Abu Shegeili Formation, Abu-Dabbab, SN: 42, pp., X 2.5	141
(91):	Goniastrean Framestone. Pleistocene Raised reefs, of Abu-Shegeili Formation, Wadi Quseir El-Kadium, SN: 26, CN., X 6	143
(92):	Favian Framestone. Pleistocene Raised reefs, of Abu-Shegeili Formation, Wadi Sherm El-Bahari, SN: 19, CN., X 6	143
(93):	Sandy wackestone. Pleistocene Raised reefs of Abu-Shegeili Formation, Wadi Quseir El-Kadium, SN: 12, CN., X x 6	144
(94):	Enlarged part of the same. CN., X 20	144
(95):	Molluscan packstone, Pleistocene Raised reefs, of Abu Shegeili Formation, Wadi Quseir ElKadium, SN : Q 11, CN., X 6	146
(96):	Enlarged of the same section. CN., X 20	146
(97):	Fossiliferous/algal (Red algae) grainstone. Pleistocene Raised reefs, of Abu-Shegeili Formation, Wadi Quseir El-Kadium, SN: Q5, CN. X 6.	154



- (98): Sandy Fossiliferous packstone (biomicrite) with bioclastics (bivalve, Foraminifera and red algae) embedded in micritic groundmass, notice the partial and complete micritization Shagra Formation, Wadi Essel SN: 18, pp. X 2.5. 154
- (99): Other part of the same with red algae, foraminifera, echinoids and coarse quartz grains and partial recrystallization of micrite and micritization of skeletal particles. pp. X 2.5. 155
- (100): Otherpart of the same thin section, notice the filling of the bivalve by the same micritic and sandy matrix of the whole rock,pp.X2.5. 155
- (101): Sandy packstone (biomicrite), notice the partial recrystallization of the micrite and the shell fragment, Shagra Formation, Wadi Essel, SN: 23, pp., X 2.5. 156
- (102): Algal sandy wackestone, notice the micritization of the bioclastic. Shagra Formation, Wadi Um- Gheig, SN : 9, CN., X 2.5. 156
- (103): Sandy wackestone with polycrystalline quartz grains, notice the partial recrystallization of micrite, Shagra Formation, Wadi Abu-Dabbab, SN: 36, pp., X 2.5. 157
- (104): Sandy packstone (biomicrite), notice the recrystallization of the shell fragments and the filling of the fossil intraspaces with sparite. Shagra Formation, Wadi Essel, SN: 13, pp., X 2.5. 157
- (105): Sandy packstone (Sandy biomicrite) with bivalves, miliolids and *Borlis sp.*, notice the partial recrystallization of the micrite to microsparite. Shagra Formation, Wadi Essel, SN: 23, pp., X 2.5. 162
- (106): Sandy wackestone with shell fragments, notice the recrystallization of the bioclastics, Shagra Formation, Wadi Abu -Dabbab, SN: 36, CN., X 2.5. 162
- (107): Other part of the same thin section, notice the recrystallization of the bioclastics and onchoids. CN., X 2.5. 163
- (108): Algal sandy wackestone, notice the partial recrystallization of micritic matrix, Shagra Formation, Wadi Um-Gheig,SN:9,pp., X 4 163
- (109): Bufflestone with miliolids, notice the micritization of miliolids and the other bioclasts, Shagra Formation, Wadi Abu-Dabbab, SN: 34, CN., X 2.5 164