

Updated Role of Magnetic Resonance Imaging in Assessment of Endometriosis

Thesis

*Submitted for Fulfillment of the Master Degree
in Radiodiagnosis*

By

Shayan Abd El-Aziz Mohamed

M.B.B.Ch.

Under Supervision of

Prof. Dr. Safaa Kamal Mohamed

Professor of Radiodiagnosis

*Radiodiagnosis and Nuclear Medicine Department
Faculty of Medicine - Ain Shams University*

Dr. Mary Yaftah Tadros

Lecturer of Radiodiagnosis

*Radiodiagnosis and Nuclear Medicine Department
Faculty of Medicine - Ain Shams University*

**Faculty of Medicine
Ain Shams University
2019**



Acknowledgement

*Thanks first and last to **ALLAH** for his guidance, support and care in every step in our lives.*

*I am most thankful to **Prof. Dr. Safaa Kamal Mohamed**, Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for her continuous support and guidance not only in this work, but throughout my career.*

*Moreover, I would like to express my sincere and deepest gratitude to **Dr. Mary Yaftah Tadros**, Lecturer of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for her supervision, help and valuable support and guidance, I am deeply affected by her character, perfection, care and consideration.*

Dr. Shayan Abd El-Aziz

Table of Contents

Subjects	Page
List of Abbreviations	I
List of Tables.....	II
List of Figures	III
Introduction.....	1
Aim of the Work.....	4
Review of Literature	
Chapter (1): Gross Anatomy of the Female Pelvis	5
Chapter (2): Normal MRI Imaging Appearance of the Female Pelvis.....	30
Chapter (3): Pathology of the Endometriosis	42
Chapter (4): Technique of MRI in Endometriosis.....	56
Chapter (5): MRI Characteristics and Locations of Endometriosis.....	72
Patients and Methods.....	108
Results.....	115
Illustrative Cases.....	121
Discussion.....	152
Summary & Conclusion	161
References	164
Arabic Summary.....	-

List of Abbreviations

Abb.	Full term
ADC	Apparent diffusion coefficient
AWE	abdominal wall endometriosis
BMI	Body mass index
CT	Computed Tomography
DBCE	Double barium contrast enema
DCE	Dynamic contrast enhanced
DIE	Deep infiltrating endometriosis
DPE	Deep pelvic endometriosis
DWI	Diffusion-weighted image
E2	Estradiol
EPI	echo planar imaging
FOV	Field of view
ICD	Implanted cardioverter defibrillator
MRI	Magnetic resonance imaging
SPIR	Spectral presaturation with inversion recovery.
SWI	Susceptibility-weighted image
T1WI	T1 weighted image
T2WI	T2 weighted image
TAUS	Transabdominal ultrasound
TE	Echo time
TR	Repetition time
TRA	Transverse
TRUS	Transrectal ultrasound
TSE	Turbo spin echo
TVUS	Transvaginal ultrasound
US	Ultrasound
UTE	Urinary tract endometriosis

List of Tables

No	Table	Page
4.1	Dynamic contrast enhanced MRI protocol.	65
4.2	Conventional MRI protocol.	69
4.3	Contrast-enhanced T1WI protocol.	70
4.4	Dynamic contrast enhanced MRI protocol.	70
4.5	DWI protocol.	70
4.6	SWI protocol.	71
5.1	MRI imaging features of endometriotic lesions.	74
6.1	The number and percentage of different clinical presentation in childbearing period.	109
6.2	The MRI sequences used in the study	112
7.1	Age distribution of the studied participants.	115
7.2	MRI Findings in the study participants.	116
7.3	Laparoscopic Findings in the study participants.	118
7.4	Correlation of MRI results with Laparoscopic results.	119
7.5	Validity of MRI in the diagnosis of endometriosis.	120

List of Figures

No	Figure	Page
1.1	Attachment of uterine tube, round ligament, and ligament of the ovary to superolateral angle of the uterus.	7
1.2	Ligaments, fascia, and spaces of uterus.	9
1.9	Peritoneum of female pelvis	11
1.4	Microstructure of the uterus.	14
1.5	Internal female genital organs.	16
1.6	The broad ligament and blood supply to the uterus and ovaries.	18
1.7	The relations of the vagina.	22
1.8	Microstructure of the ovary	26
1.9	Relationship of bladder to the neighboring organs.	29
2.1	Sagittal T2WI demonstrating the three compartments of the pelvis.	30
2.2	Sagittal and axial T2WI through female pelvis showing anus, bladder, cervix, rectum, pubic symphysis, vagina, and the junctional zone of the uterus.	32
2.3	Axial T2WI shows normal female urethra.	34
2.4	Axial T2WI showing three distinct zones of female urethra.	36
2.5	Axial T2WI in two patients (a, b) show variable thickness in the vagina depending on estrogen levels.	38
2.6	Coronal T2WI showing the uterus and both ovaries.	39
2.7	Sagittal T2WI showing normal female urinary bladder, uterus, and rectum.	41

List of Figures

No	Figure	Page
3.1	Common sites of endometriosis.	43
3.2	Macroscopic appearances of endometriosis according to the activity of the lesion.	47
3.3	Macroscopic appearances of ovarian endometriotic cyst.	48
3.4	Microscopic appearances of endometriosis.	49
3.5	Typical ultrasound appearances of endometriotic cyst.	54
3.6	Axial CT scan of scar showing AWE. the lesion can be seen involving the fascia and pressing on the rectus muscles	55
4.1	Sagittal 2D T2WI performed at 1.5 Tesla showing the benefits of patient preparation on image quality.	59
4.2	Sagittal 2D T2WI performed at 1.5 Tesla following vaginal and rectal opacification with sonographic gel.	60
4.3	Sagittal 2D T2WI in the same patient performed at 1.5 Tesla and 3.0 Tesla provide similar good Imaging quality for the evaluation of pelvic anatomy.	63
5.1	Axial T1WI, T2WI and fat-suppressed T1WI show bilateral ovarian endometrioma in 39-years-old female.	76
5.2	Axial T2WI, and fat-suppressed T1WI show endometriosis ob bilateral uterosacral ligaments in 29-years-old female.	78
5.3	Sagittal T2WI showing female pelvic compartments	79
5.4	Sagittal T2WI shows solid endometriosis of vesicouterine pouch.	80

List of Figures

No	Figure	Page
5.5	Sagittal T2WI shows deep infiltrating endometriosis of the bladder.	81
5.6	Sagittal and coronal T2WI show ureteral endometriosis.	82
5.7	Axial fat-saturated T1WI, axial and sagittal T2WI show severe pelvic endometriosis in 45-years-old female.	84
5.8	Axial T2WI shows deep infiltrating endometriosis of uterine serosa with ovarian endometrioma.	85
5.9	Axial T1WI with and without fat suppression demonstrating haematosalpinx	86
5.10	Axial T2WI and coronal T2WI show deep endometriosis of the right round ligament of the uterus.	87
5.11	Sagittal, oblique coronal, and oblique axial T2WI show endometriosis of the retrocervical region in a 39-year-old female.	89
5.12	Sagittal T2W, T1WI, and sagittal T1W post contrast show endometriosis of rectovaginal in 39-year-old patient.	90
5.13	Sagittal T2WI shows rectal extra luminal mass, solid invasive endometriosis.	91
5.14	Axial T2WI and T1WI show endometriosis of uterosacral ligaments.	92
5.15	Sagittal T2W MRI showing rectovaginal septum endometriosis.	94
5.16	Sagittal, oblique axial, oblique coronal fat-suppressed T1WI, and sagittal, oblique axial and oblique coronal T2WI show small endometriotic implants involving the posterior vaginal fornix and the cervix in a 46-year-old woman.	95

No	Figure	Page
5.17	Sagittal and axial T2WI MRI showing endometriosis invade the peripheral layer of the rectal wall.	96
5.18	Sagittal and axial T2WI show bilateral ovarian endometriotic cysts and posterior cul-de-sac obliteration in 34-year-old woman.	97
5.19	Axial pre-contrast T1WI, axial T2WI, DWI, and ADC map demonstrate bilateral hydrosalpinx, and complex lesion in the right adnexa in 27-year-old woman presenting with infertility.	99
5.20	Fast spin-echo T2WI, fat-saturated spin-echo T1WI, and SWI show abdominal wall endometriosis in 34-year-old woman.	101
5.21	Sagittal, and axial T2WI, sagittal, and axial post-contrast T1WI, DWI, and ADC maps show malignant transformation in endometriosis.	104
5.22	Axial T1WI, T2WI, T1WI post-contrast, and DWI show endometrioid adenocarcinoma associated with endometriotic cyst in 45-year-old woman.	105
7.1	Age distribution of the studied participants.	115
7.2	MRI Findings in the study participants.	117
7.3	Laparoscopic Findings in the study participants.	118
8.1	Axial T1W (a), axial T2W (b), sagittal T1W post contrast (c) and axial DWI (d) show a poorly margined anterior pelvic wall lesion measuring 2.5 x 2.0 cm (arrow) displaying low signal intensity in T1W, heterogenous low and isointense T2 signal, enhancement in post contrast study and restricted diffusion in DWI. The lesion is seen overlying and partially non separable from the left rectus abdominis, feature suggesting abdominal wall endometriosis.	121

No	Figure	Page
8.2	Axial T1WI (a), T2WI (b), coronal T1WI post contrast (c) and axial DWI (d) show Left ovarian cystic lesion measuring 4.7x5.5cm., with thick wall (arrow), shows high intensity in T1W1, intermediate intensity in T2W1 (T2 shading), post contrast enhancing and diffusion restriction in DWI feature of endometrioma. The uterus is seen deviated to the right side.	122
8.3	Axial T1WI (a), axial T2WI (b), sagittal T2WI (c) and axial T2WI SPIR (d) show irregular solid mass (arrow), measuring about 1.9 x 1 cm., seen at the vesicovaginal septum on the left side as well as anterior vaginal wall and anterior uterine wall (c). The mass is hypointense on T2WI, intermediate with hyperintense focus on T1WI, this focus is still hyperintense on fat suppressed images (d) denoting bloody or protineous content. Obliteration of rectovaginal pouch is also noticed these features mostly represent endometriotic implant at the vesicovaginal septum, anterior uterine and vaginal walls.	124
8.4	Axial T1WI (a), T2WI (b), T2WI SPIR (c) and DWI (d) show the uterus with a small mural left sided hemorrhagic cystic lesion (white arrows) showing high on T1WI, T2WI and preserved signal in fat suppressed image (c) with diffusion restriction on DWI. The lesion surrounded by low signal rim, measuring nearly 1 cm in diameter (possibility of cystic adenomyosis). Right ovarian cyst (yellow arrows) is also noticed measuring about 2.6 cm in diameter containing simple fluid with low signal in T1WI and high signal in T2WI.	126

No	Figure	Page
8.5	Axial T1WI (a), sagittal T2WI (b), axial T2WI SPIR (c) and DWI (d), in these images the uterus is seen bulky measuring about 14 x 11 x 10 cm. with arcuate shape. The myometrium is seen diffusely thickened with hypointense signal on T1WI and T2WI, it shows few tiny cystic lesions of hyperintense signal on T1WI and T2WI denoting hemorrhage. There are also innumerable tiny cystic foci embedded within of high signal on T2WI denoting cystic changes., picture is suggestive of diffuse form of adenomyosis with subsequent distorted uterine cavity. There are few subserosal cystic lesions of high signal on T2WI seen mainly at anterior wall most likely represents subserosal endometriotic cyst (arrows). Note preserved signal in fat suppressed image (c) and restricted diffusion in DWI.	128
8.6	Axial T1WI (a) and sagittal T2WI (b&c), axial T2WI SPIR (d) and DWI (e) show two left adnexal cystic masses (arrows). The largest is seen at the Douglas Pouch, measuring 16 x 11.5 x 7.5 cm, it shows multiple thin septae. The smaller measures 5 x 4.8 x 4.3 cm. They show high T1WI SI. The first one is intermediate intensity on T2WI and the smaller one is of low T2WI (T2 shading). They also show no suppression at fat suppressed sequence (d) and diffusion restriction in DWI (e) picture is suggestive of adnexal endometrioma.	130
8.7	Axial T1WI (a), coronal T1WI post contrast (b), coronal T2WI (c&d), axial T2WI SPIR (e) and axial DWI (f) show Large right adnexal cystic lesion (white arrows) measuring 7.1 x7 cm. It exhibits high signal on T1WI and intermediate signal on T2WI (T2 shading). A	132

No	Figure	Page
	similar cystic lesion is seen at the left ovary (yellow arrows) measuring 2.7x3.3cm. It shows blood/fluid level and few floating dark signal lesions on all pulse sequences (dark spot) denoting old blood component. They also show post contrast enhancement in right (white arrows) and left (yellow arrows) ovarian cysts(b) with no suppression of signal on SPIR (e) and diffusion restriction with variable degreesin DWI (f) features suggesting bilateral ovarian endometriomas.	
8.8	<p>(a-d): Axial T1WI (a), coronal T2WI (b), axial T2WI SPIR (c) and axial DWI (d) show multiple follicles in the right ovary most of them show high signal on T1WI and low signal on T2WI (T2 shading) with fluid level (white arrows) They also show preserved signal in fat suppressed image (c) and restricted defusion on DWI (d) features suggesting endometrioma, the endometrial thickness is normal but with ill defined junctional zone shows segmental areas of increased thickness reaching up to 19 mm especially the left wall (star) (b) mostly segmental adenomyosis left ovarian cyst is also noticed with low signal in T1WI and high signal in T2WI (yellow arrows).</p> <p>(e-g): Axial T1WI (e) sagittal T2WI (f) sagittal T1WI post contrast (g) show ill defined lesion (arrow) measuring 3.3 x 2.3 cm seen in the right aspect of the anterior abdominal wall showing low signal on T1WI and intermediate signal on T2WI wih inhomogenous post contrast enhancement, likely at the site of Cescerian scar mostly represents endometriotic implant at scar. It reaches the right rectus abdominus muscle</p>	134

No	Figure	Page
	inseparable from it, features suggesting abdominal wall endometriosis.	
8.9	Axial T1WI (a), axial T2WI (b), sagittal T2WI (c), axial T2WI SPIR (d), sagittal T1WI post contrast (e) and axial DWI (f) show a small lesion (arrow) measuring 3x2 cm is noted in the douglas pouch with slightly high signal In T1WI and low signal in T2WI with central area of high T2 signal. On axial images the lesion appears to be separate from the uterus but on sagittal image (c) there is pedicle connecting it to posterior uterine wall. Lesion show no contrast enhancement on post contrast image (e), increase signal intensity in SPIR (d) and diffusion restriction in DWI (f) denoting endometriotic implant in Douglas pouch.	136
8.10	Axial T1WI (a), axial T2WI (b), coronal T1WI post contrast (c), axial T2WI SPIR (d) and axial DWI (e) show Left multi-locular cystic adnexal lesion (white arrows) measuring 9x5 cm with high signal in T1WI and intermediate signal in T2WI. The Right ovary shows a small cyst (yellow arrow) with intermediate signal in T1WI and low signal in T2WI (features suggesting Left adnexal endometriosis and right chocolate cyst). Right adnexal convoluted tubular cystic lesion (arrow head) having intermediate signal in T1WI and high signal in T2WIs likely representing haemosalpinx. Subserous lesion (star) measuring 3 cm in diameter is noted related to the posterior uterine wall showing isointense signal in T1WIs and hypointense signal in T2WI (possibility of endometriotic implant on serosal surface of uterus). The post contrast image (c) show rim contrast enhancement and enhancing thin septae likely representing hemorrhagic	138

No	Figure	Page
	products in the left cystic adnexal lesion and thick wall enhancement in the right ovarian cyst. They also show preserved signal in fat suppressed images (d) Features suggesting Left adnexal endometrisosis and right chocolate cyst. The DWI show restricted diffusion in the left adnexal lesion (white arrow) and no diffusion restriction in the right ovarian cyst (yellow arrow). Features suggesting Left adnexal endometrisosis and right chocolate cyst.	
8.11	Axial T2WI (a&b), xial T2WI SPIR (c&d) and axial DWI (e&f) show right adnexal cystic lesion (white arrow), measuring 7.2 x 6.5 cm, displaying T2 shading with fluid fluid level another smaller cystic lesion is seen in the right ovary (red arrow), measuring 2.7 x 4.0 cm display high T2 signal intensity with non dependent low intensity component (a). Left ovary shows small 1.5 x 1.6 mm cystic lesion (yellow arrow), display T2 shading (b) denoting Large right adnexal & small left ovarian endometrioma and right ovarian hemorrhagic cyst. The large right adnexal (white arrows) & small left ovarian (yellow arrows) endometriomas and right ovarian (red arrows) hemorrhagic cyst show preserved signal in fat suppressed images (c&d). With diffusion restriction in DWI (e&f).	140
8.12	Axial T1WI (a), axial T2WI (b), axial T2WI SPIR (c) and DWI (d) show Multi-septated left adnexal cystic lesion (arrow) displaying high signal in T1 and intermediate signal in T2 WI, with high signal in fat suppression image (c) and diffusion restriction in DWI (d). It measures about 54 x 31 mm, MRI criteria is suggestive of hemorrhagic endometriotic cyst.	142

No	Figure	Page
8.13	Axial T1WI (a), T2WI (b), T2WI SPIR (c) and DWI (d) the uterus shows asymmetrical thickening of the junctional zone measuring 12.5 mm. in greatest dimension with blurred interface between the junctional zone and the outer myometrium, it also displays hyperintense T2 microcysts (arrows). Multiple low T2 intensity focal lesions seen within the myometrium, with preserved signal intensity in fat suppressed image (c). And restricted diffusion in DWI (d), likely representing adenomyomas.	144
8.14	Coronal T1WI (a) T2WI (b) axial T2WI (c) and axial T2WI SPIR (d) Uterus is seen bulky (arrow) shows blurred junctional zone reaching up to 15 mm posteriorly. With preserved signal intensity in fat suppressed image (d), likely representing adenomyomas.	146
8.15	(a&b): Axial T1WI (a) and sagittal T2WI (b) show anterior & posterior uterine wall focal lesions merging with the junctional zone (white arrows) measuring about 21mm with indistinct margin & with small cysts seen inside likely denoting focal adenomyosis. (c-f): Axial T1WI (c), T2WI (d), T2WI SPIR (e) and DWI (f) show right ovarian large follicle (white arrows) measuring 22x34 mm, with low signal in T1WI, and relatively high signal in T2WI with preserved signal in fat suppressed image (e). Left ovary displays small thin wall cystic lesion (yellow arrows) measuring 27 x 20 mm that shows high T1 signal intensity & T2 shading with diffusion restriction in DWI (f), and preserved signal in fat suppressed image (e) denoting bilateral endometriotic cyst.	148