

سامية محمد مصطفى



شبكة المعلومات الجامعية

بسم الله الرحمن الرحيم



سامية محمد مصطفى



شبكة المعلومات الجامعية



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



سامية محمد مصطفى



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



سامية محمد مصطفى



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



سامية محمد مصطفى



شبكة المعلومات الجامعية



بالرسالة صفحات لم ترد بالأصل




**STUDIES ON ENTERIC PROTOZOA
INFECTING CHICKENS IN PORT-SAID PROVINCE**

A Thesis


**Presented by
Samar William Abd El-Malek Tadros
(B.V.Sc. Suez Canal University 1992)**

**For
The Degree of M.V.Sc.
(Parasitology)**



Prof. Dr. H.M. El-Gwady
Prof. and Head of Parasitology
Dept. Faculty of Vet. Med.
Suez Canal University

Under the Supervision of



Prof. Dr. Abd El-Moneim A. El-Meligy
Prof. of pathology and Vice Dean
Faculty of Vet. Med
Suez Canal University



Dr. M. W. Ghattas
Researcher at Animal Health Institute
(Port-Said Branch)

Department of Parasitology
Faculty of Veterinary Medicine
Suez Canal University
1997

B

1811E

Suez Canal University
Faculty Of Vet . Med .
Department of Parasitology

APPROVAL SHEET

This to approve that the dissertation submitted by **SAMAR WILLIAM ABD EL-MALEK TADROS** to Suez Canal University entitled (Studies on enteric Protozoa infecting chickens) For the M.V.SC. Degree has been approved in / /1997 by the examining committee:-

Prof. 1- **Dr.H.M.EL.GWADY .**

Prof. and Head of Parasitology Dept.Faculty of Vet . Med . ,
Suez Canal University .

Prof. 2- **Dr . A.A.EL.MELIGY**

Prof.of Pathology and Vice Deam of Faculty of Vet.Med.
Suez Canal University.

3- **Dr . ALY A. A. HASSAN**

Prof. of Parasitology and Head of Parasitology Dept .,
Faculty of Vet.Med., Cairo University.

4- **Dr.ISMAIL M.SHALABY .**

Associate Prof. Zoology Dept., Faculty of Science.Suez
Canal University .

Approved by Faculty Council in / /1997

ACKNOWLEDGMENT

I wish to express my deepest cardinal thanks and gratitudes to;

Prof. Dr. H.M.EL-Gwady

Prof. Dr. A.A.EL- Meligy

Dr. M.G. Hassan

Dr. A.A. Abdel Aal

Dr. M.W. Ghattas

Dr. M.M EL-Hamamy

For their help, assistance and guidance in completion of this work.

CONTENTS

	page
1-Introduction	1
2- Review of Literatures ;	3
2-a; <i>Eimeria</i> species;	3
2-a-1; Incidence & seasonal dynamics .	3
2-a-2; Diagnosis of <i>Eimeriasis</i> .	10
2-a-3; Identification of <i>Eimeria</i> species ;	12
2-a-3-1; Oocyst morphology, sporulation time and pathogenicity of <i>Eimeria</i> species ;	12
2-a-3-1-1; <i>E. acervulina</i> .	12
2-a-3-1-2; <i>E. mitis</i> .	16
2-a-3-1-3; <i>E. tenella</i> .	20
2-a-3-1-4; <i>E. necatrix</i> .	22
2-a-3-1-5; <i>E. praecox</i> .	25
2-a-3-1-6; <i>E. maxima</i> .	28
2-a-3-1-7; <i>E. brunetti</i> .	31
2-a-3-1-8; <i>E. mivati</i> .	34
2-a-3-1-9; <i>E. hagani</i> .	37
2-a-3-2; Sporulation of <i>Eimeria</i> species oocysts ;	40
2-a-3-2-1; Effect of physical factor on sporulation .	42
2-a-3-2-2; Effect of chemical factor on sporulation ;	43
1- Formalin	43
2- Sodium chloride .	45
3- Hcl .	46
4- H ₂ SO ₄ .	46
2-b; <i>Cryptosporidium</i> species ;	46
2-b-1; Incidence and seasonal dynamics .	46
2-b-2; Morphology and diagnosis of <i>Cryptosporidiosis</i> species	49
2-b-3; Taxonomy .	57

2-c; Pathological studies;	59
2-c-1; <i>Eimeriasis</i> •	61
2-c-2; <i>Cryptosporidiosis</i> •	75
3- Material and Methods ;	81
3-1; Parasitological studies •	81
3-2; Pathological studies •	84
4-; Results ;	86
4-1; Parasitological studies ;	86
4-1-1; Incidence of <i>Eimeria</i> species •	86
4-1-2; Monthly and seasonal incidence of <i>Eimeria</i> species •	89
4-1-3; Incidence of different <i>Eimeria</i> species •	92
4-1-4; Identification of the detected <i>Eimeria</i> species according to site, morphological features and sporulation time ;	96
4-1-5-a; Morphological status of <i>Eimeria</i> species detected at one site of intestinal tract.	100
4-1-5-b; Morphological status of <i>Eimeria</i> species detected in two sites of intestinal tract.	102
4-1-5-c; Morphological status of <i>Eimeria</i> species detected in four sites of intestinal tract.	104
4-1-6; Incidence of <i>Cryptosporidium</i> species;	107
4-1-7; Monthly and seasonal incidence of <i>Cryptosporidium</i> species •	110
4-1-8; Seasonal incidence of <i>Cryptosporidium</i> species.	114
4-1-9; Effect of physical and chemical factors on sporulation	117
4-1-9-a; Effect of physical factors •	117
4-1-9-b; Effect of chemical factors •	117
4-1-10; Morphological description of <i>Cryptosporidium</i> sp.	120
4-2; Pathological studies	121
Discussion •	125
Summary •	146
References •	160
Arabic Summary •	161

LIST OF TABLES

- Table (1) : Incidence of *Eimeria* species detected among chickens examined at different localities at Port-Said Province .
- // (2) : Monthly and seasonal incidence of *Eimeria* species detected among chickens examined at different localities at Port- Said Province .
- // (3) : Incidence of different *Eimeria* species detected among chickens examined at different localities at Port-Said Province.
- // (4) : Identification of the detected *Eimeria* species according to site, morphological features and sporulation time .
- // (5,a) : Morphological status of *Eimeria* species detected at one site of intestinal tract .
- // (5,b) : Morphological status of *Eimeria* species detected in two sites of intestinal tract .
- // (5,c) : Morphological status of *Eimeria* species detected in four sites of intestinal tract .
- // (6) : Incidence of *Cryptosporidium* species detected among chickens examined at different localities at Port-Said Province
- // (7) : Monthly and seasonal incidence of *Cryptosporidium* species detected among chickens examined at different localities at Port-Said Province .
- // (8) : Seasonal incidence of *Cryptosporidium* among chickens examined at Port -Said Province .
- // (9) To (17)
- : Comparison between the dimensions of the different *Eimeria* spp. detected during this study with those of the previous authors;
- // (9) : *E. acervulina*.
- // (10) : *E. mitis*

Table (11) : *E. tenella*

// (12) : *E. necatrix*

// (13) : *E. praecox*

// (14) : *E. maxima*

// (15) : *E. brunetti*

// (16) : *E. mivati*

// (17) : *E. hagani*

LIST OF FIGURES

Fig. (1) : Bar- chart of the incidence of *Eimeria* species detected among chickens examined at different localities at Port-Said Province .

// (2) : Bar- chart of the monthly and seasonal incidence of *Eimeria* species detected among chickens examined at different localities at Port-Said Province .

// (3) : Bar- chart of the incidence of different *Eimeria* species detected among chickens examined at different localities at Port-Said Province .

// (4) : Bar- chart of the identification of the detected *Eimeria* species according to site , morphological features and sporulation time .

// (5) : Bar- chart of the incidence of *Cryptosporidium* species detected among chickens examined at different localities at Port-Said Province .

// (6) : Bar- chart of the monthly and seasonal incidence of *Cryptosporidium* species detected among chickens examined at different localities at Port-Said Province .

// (7) : Bar- chart of the seasonal incidence of *Cryptosporidium* among chickens examined at Port -Said Province .

// (8) : *Eimeria acervulina*; non sporulated oocyst with micropyle cap from scraping of duodenum X 400.

Fig. (9) : *Eimeria mitis*; non sporulated oocyst without micropyle from scraping of duodenum X 400.

// (10) : *Eimeria mitis*; sporulated oocyst without micropyle duodenum X 400.

// (11) : *Eimeria tenella*; non sporulated oocyst with micropyle cap from scraping of caecum X 400.

// (12): *Eimeria tenella*; sporulated oocyst with micropyle cap from scraping of caecum X 400.

// (13): *Eimeria tenella*; sporulated oocyst without micropyle from scraping of caecum X 1000.

// (14) : *Eimeria necatrix*; non sporulated oocyst without micropyle from scraping of caecum X 1000.

// (15): *Eimeria necatrix*; sporulated oocyst without micropyle micropyle from scraping of caecum X 400.

// (16) : *Eimeria praecox*; unsporulated oocyst with micropyle from scraping of jejunum X 1000.

// (17) : *Eimeria maxima*; sporulated oocyst without micropyle from scraping of ileum X 400.

// (18) : *Eimeria maxima*; sporulated oocyst with micropyle from scraping of caecum X 1000.

// (19) : *Eimeria brunetti*; non sporulated oocyst with micropyle from scraping of jejunum X 400.

// (20): *Eimeria brunetti*; non sporulated oocyst with micropyle cap from scraping of caecum X 1000.

// (21) : *Eimeria mivati*; non sporulated oocyst with micropyle cap from scraping of duodenum X 400.

// (22) : *Eimeria hagani*; non sporulated oocyst without micropyle cap from scraping of ileum X 400.

// (23) : *Eimeria species* (n.sp.); non sporulated oocyst without micropyle from scraping of caecum X 400.

Fig.(24) : *Eimeria species*(n.sp.); non sporulated oocyst with micropyle from scraping of caecum X 400.

// (25) : *Eimeria maxima* and *Eimeria tenella*; sporulated oocysts without micropyle from scraping of caecum X 1000.

// (26) : Caecal coccidiosis; sporulated and non sporulated oocysts in different size and shape from scraping of caecum X 400.

// (27) : *Cryptosporidium baileyi* stained with modified-Zehil Neelson X 1000.

// (28) : *Cryptosporidium baileyi* stained with modified-Zehil Neelson X 1000.

// (29) : Section of Intestine of chicken infected with *E. maxima* and *E. brunetti*. X 250

// (30) : Section of higher magnification of Fig (1) to illustrate the large number of *Eimeria* stages in the villar epithelium. X 100

// (31): Section of intestine of chicken infected with *Eimeria species*. X 250

// (32): Section of intestine of chicken moderately infected with *Eimeria praecox* and *E. mivati*. X 100

// (33) : Section of intestine of chicken infected with *Eimeria maxima* and *E. praecox*. X 250

// (34) : Section of intestine of chicken mildly infected with *Eimeria species*. X 250

// (35) : Section of intestine of chicken infected with *Eimeria maxima* and *E. necatrix*. X 250

// (36): Section of caecum of chicken infected with *Eimeria maxima* and *E. tenella*. X 250

// (37) : Section of caecum, of chicken infected with *Eimeria tenella*. X 100

// (38) : Section of caecum of chicken infected with *Eimeria tenella* and *E. brunetti*. X 100

// (39) : Section of intestine of chicken infected with *Cryptosporidium baileyi*.

Fig.(40) : Section of bursa of fabricius of chicken infected with *Cryptosporidium baileyi*. X 250

// (41) : Higher magnification of Fig. (40) to illustrate the *Cryptosporidium baileyi*. X 1000

// (42) : Section of bursa of fabricius chicken infected with *Cryptosporidium baileyi*. X 250