

# Added Value of Ultrasonography in Diagnosis of Rotator Cuff Injury

#### Thesis

Submitted for Partial Fulfillment of Master Degree in **Radio diagnosis** 

Presented by

Rehab Abdelbaqy Ibrahim Mohammed
(M.B.B.CH)

#### Supervised By

#### Prof. Dr. Mohsen Gomaa Hasan

Professor of Radio diagnosis Faculty of Medicine - Ain Shams University

#### Dr. Tarek Wahby Hemeda

Lecturer of Radio diagnosis Faculty of Medicine - Ain Shams University

Faculty of Medicine - Ain Shams University
2019



سورة البقرة الآية: ٣٢

#### Acknowledgments

First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.

I wish to express my deepest thanks, gratitude and appreciation to **Prof. Dr. Mohsen Gomaa**\*\*Basan\*, Professor of Radio diagnosis, Faculty of Medicine, Ain Shams University, for his meticulous supervision, kind guidance, valuable instructions and generous help.

Special thanks are due to **Dr. Tarek Wahby Themeda**, Lecturer of Radio diagnosis, Faculty of

Medicine, Ain Shams University, for his sincere

efforts, fruitful encouragement.

I would like to express my hearty thanks to all my family for their support till this work was completed.

Rehab Abdelbaqy Ibrahim Mohammed

### Tist of Contents

Title	Page No.
List of Abbreviations	5
List of Tables	7
List of Figures	9
Introduction	1 -
Aim of the Work	22
Review of Literature	
Anatomy of the Shoulder Joint	23
Pathology of Rotator Cuff Tears	65
■ Technique of MRI and Ultrasound of Shoulder	98
High Resolution Ultrasound and MRI Assessment Painful Shoulder Joint	
Patients and Methods	150
Results	154
Case Presentation	173
Discussion	187
Summary	193
Conclusion	195
References	196
Arabic Summary	

### Tist of Abbreviations

Abb.	Full term
AC	A cromion
ACJ	Acromioclavicular joint
AHD	Acromiohumeral distance
AHI	Acromio-humeral interval
AP	Antero-posterior
HAD	Calcium hydroxyapatite deposition
CHL	Coraco-humeral ligament
CL	Clavicle
Co	Coracoid process
Del	$Deltoid\ muscle$
<i>F</i>	Fluid
Fig	Figure
<i>GT</i>	Greater Tuberosity
<i>HH</i>	Humeral Head
<i>IGHL</i>	Inferior gleno-humeral ligament
InfraS	$ In fraspinatus\ muscle$
<i>INT R</i>	Internal rotation
<i>LHB</i>	Long head of biceps
<i>LS</i>	$ Longitudinal\ section$
<i>Lt</i>	Left
<i>LT</i>	Lesser tuberosity
<i>Max</i>	Maximum
Min	Minimum
<i>MGHL</i>	Middle gleno-humeral ligament

### Tist of Abbreviations cont...

Abb.	Full term
MRI	Magnetic Resonance Imaging
PA	Postero-anterior
<i>RC</i>	Rotator Cuff
<i>RT</i>	Right
SGHL	Superior gleno-humeral ligament
<i>SAIS</i>	Subacromial Impingement Syndrome
SST	Supraspinatus muscle
SubS	Subscapularis muscle
<i>Tm</i>	Teres minor muscle
TS	Transverse section
US	Ultrasound
USA	United states of America

### Tist of Tables

Table No.	Title	Page No.
Table (1):	Origins, Insertions, Actions, and Supplies of the Rotator Cuff Muscles.	
<b>Table (2):</b>	Ligamentous lesions according to Rocclassification adapted for MRI	
<b>Table (3):</b>	Age and gender of the studied patients	s154
<b>Table (4):</b>	Side affected with expected cause	155
<b>Table (5):</b>	Relation to other diseases	156
<b>Table (6):</b>	Comparison between MRI & US reg supraspinatous muscle tear	-
<b>Table (7):</b>	Comparison between MRI and US regother rotator cuff muscles	-
<b>Table (8):</b>	Comparison between MRI and US reg AcJ disease	
<b>Table (9):</b>	Comparison between MRI and US reg bursal abnormality	
<b>Table (10):</b>	Relation between results of rotate muscle tendinopathy by US and MR	
<b>Table</b> (11):	Relation between results of partial the (bursal) by US and MR	
<b>Table (12):</b>	Relation between results of partial the (articular) by US and MR	
<b>Table (13):</b>	Relation between results of partial the (interstitial) by US and MR	
<b>Table (14):</b>	Relation between results of full the (incomplete) by US and MR	
<b>Table (15):</b>	Relation between results of full the (complete) by US and MR	

### Tist of Tables cont...

Table No.	Title	Page N	0.
Table (16):	Relation between results of other cuff muscles by US and MR		. 167
Table (17):	Relation between results of ACJ discussion US and MR	•	. 168
Table (18):	Relation between results of bursal and MR	v	. 169
<b>Table (19):</b>	Relation between results of total by MR		. 170
Table (20):	Diagnostic accuracy of US in detection MRI results		. 171

## List of Figures

Fig. No.	Title Page I	No.
Figure (1):	Different Acromion shapes	25
Figure (2):	Shoulder anatomy, anterior view	
Figure (3):	Shoulder anatomy, posterior view	
Figure (4):	Normal labral variants	
Figure (5):	Sublabral foramen	
Figure (6):	Sublabral recess	
Figure (7):	Buford complex	
Figure (8):	Sagittal-oblique MR image of the shoulder	
Figure (9):	Sagittal T2-weighted MR image with fasturation demonstrates prominer anterior band of the IGHL seen attaching	at nt
	high at the level of the superior labrum (A	36
<b>Figure</b> (10):	Coracohumeral ligament and rotate	or
	interval	
Figure (11):	Subcoracoid bursa and superio	
	subscapularis recess	
Figure (12):	(a-b); (a) Demonstrative sagittal view	
	the shoulder joint. (b) T1 sagittal oblique view of the shoulder	
Figure (13):	Shoulder anatomy, lateral view	
Figure (14):	Graphic image of a lateral view of the	
rigure (14).	rotator cuff muscle	
Figure (15):	Quadrilateral Space	
Figure (16):	Normal sonographic anatomy of shoulder join	
Figure (17):	Sonographic anatomy of biceps tendon	
Figure (18):	Sonographic anatomy of subscapular	
riguic (10).	muscle tendon	
<b>Figure (19):</b>	Sonographic anatomy of supraspinate	ıs
	muscle tendon	
<b>Figure (20):</b>	Subacromial subdeltoid bursa	
<b>Figure (21):</b>	Normal sonographic anatomy	
	infraspinatou muscle	
<b>Figure (22):</b>	Sonographic anatomy of gelnoid labrum	51

Fig. No.	Title	Page No.	
<b>Figure (23):</b>	Subacromial-subdeltoid burs	. •	
Figure (24):	view Biceps tendon, transverse vie		
Figure (25):	Rotator interval, transverse vi		
Figure (26):	Subscapularis tendon, longitu		
<b>Figure (27):</b>		Subscapularis tendon and coracoid process,	
<b>g</b> (= v)-	longitudinal view	<u>-</u>	
<b>Figure (28):</b>	Subacromial-subdeltoid burs		
<b>g</b> (,-	view	, 0	
<b>Figure (29):</b>	Infraspinatus tendon, longitu		
<b>Figure (30):</b>	Supraspinatus tendon, trans		
Figure (31):	Acromioclavicular joint, long		
<b>Figure (32):</b>	(a-b): Axial MR anatomy sup		
<b>Figure (33):</b>	(a-b) Coronal oblique conv	•	
	anatomy		
<b>Figure (34):</b>	(a-c): Sagittal conventional		
	(mid-sagittal, lateral and med	dial images) 64	
<b>Figure (35):</b>	Coronal graphic shows	a partial	
	undersurface tear of the ar	ticular surface	
	of the supraspinatus tendon	involving the	
	critical zone	65	
<b>Figure (36):</b>	Coronal graphic shows a l	bursal surface	
	partial tear with reactive bur	rsal changes 66	
<b>Figure (37):</b>	Coronal graphic shows a	an interstitial	
	delaminating partial tear	67	
<b>Figure (38):</b>	Coronal graphic shows a full	thickness tear	
	through the mid substa	ance of the	
	supraspinatus tendon		
<b>Figure (39):</b>	Focal full-thickness suprasp	oinatus tendon	
	tears		
<b>Figure (40):</b>	Partial-thickness tendon tear		
<b>Figure (41):</b>	Thinning of the rotator cuff		
<b>Figure (42):</b>	Tendon tears	74	

Fig. No.	Title	Page No.
Figure (43):	( <b>a-c</b> ). Tendon tears	75
Figure (44):	Medial tendon retraction	
Figure (45):	Factors affecting the medial	
1 1gui C (40).	tendon tear	
Figure (46):	Sagittal extent of rotator cuff	
Figure (47):	Involvement of the rotator inte	
Figure (48):	Disruption of the transverse for	
Figure (49):	(40) Drawing illustrates use of	•
1 19410 (10).	plane (dashed line) relative to	•
	base and acromial spine for	
	volume loss in the supraspir	_
	(41) Sagittal fat-saturated T2-	
	image. (42) Abnormal occupat	•
<b>Figure (50):</b>	Tangent sign	
<b>Figure (51):</b>	Muscle fatty degeneration and	
8 - ( )	in a rotator cuff muscle	
<b>Figure (52):</b>	Supraspinatus tendon, longi	tudinal view.
	The tendon demonstrates a h	
	echogenicity without any	_
	representative of a tear	89
<b>Figure (53):</b>	Subscapularis tendon, longitudi	nal view 90
<b>Figure (54):</b>	Supraspinatus tendon, longitu	ıdinal view 90
<b>Figure (55):</b>	Supraspinatus tendon, longitud	inal view 91
<b>Figure (56):</b>	Subscapularis tendon, longitudi	nal view 91
<b>Figure (57):</b>	Greater tuberosity, transverse	e view 92
<b>Figure (58):</b>	Supraspinatus tendon, transv	erse view 92
<b>Figure (59):</b>	Supraspinatus tendon, longitu	ıdinal view 93
<b>Figure (60):</b>	Supraspinatus tendon, longitud	dinal view 93
<b>Figure (61):</b>	Subscapularis tendon, longitu	dinal view 94
<b>Figure (62):</b>	Supraspinatus tendon, longitu	ıdinal view 94
<b>Figure (63):</b>	Long head of biceps (LHB)	tendon (short
	axis)	101

Fig. No.	Title	Page No.
Figure (64):	Long head of biceps (LHB)	
<b>Figure (65):</b>	a-Subscapularis tendon (lon	
	Subscapularis tendon (long axi shows hyperechoic and fibrillars tendon (arrows)	s). US image subscapularis103
<b>Figure (66):</b>	<ul><li>a): Subscapularis tendon (</li><li>b): US image shows hyper</li></ul>	-
	fibrillar subscapularis tendon (	
<b>Figure (67):</b>	Supraspinatus tendon (short a	xis)104
<b>Figure (68):</b>	a) Acromio-clavicular joint. (b)	•
	shows acromio-clavicular joint	
	characteristic hyperechoic bon	•
T' (00)	the distal clavicle (C) and acro	
<b>Figure (69):</b>	(a) Dynamic assessment for	
	impingement. (a)T ransduce over superolateral aspect of	_
	neutral position. (b) after abd	
	shoulder	
<b>Figure (70):</b>	Dynamic assessment for	
8 . ,	impingement	
<b>Figure (71):</b>	(a) Dynamic assessment for	
	muscle. (b) After adduction of	
	with the hand over the other sl	
Figure (72):	Teres minor muscle	
<b>Figure (73):</b>	MR planes showing how t	
Figure (74):	adequate shoulder MR Imaging planes of MRI shoulde	
Figure (74):	Coronal fat-suppressed T2-w	
1 1gui (10).	image shows distention of	
	subdeltoid bursa without rota	
	(arrow)	

Fig. No.	Title	Page No.
Figure (76):	Sagittal oblique T2WI, type I fla (A), type II concave under surfa type III concave under sur anterior hook (C)	ace (B) and rface with
<b>Figure (77):</b>	Coronal sonography view of le with subacromial impingement.	ft shoulder
Figure (78):	Coronal sonography view of le with shoulder impingement syno	
Figure (79):	1 0	nage of in shows thickening
<b>Figure (80):</b>	Axial T1-weighted image show acromiale	ving the os
<b>Figure (81):</b>	Acromion, transverse view	
<b>Figure (82):</b>	Full-thickness supraspinatus tea	
<b>Figure (83):</b>	Partial-thickness subscapularis	
<b>Figure (84):</b>	Coronal oblique fat-sat PD WI	
<b>Figure (85):</b>	US scans	126
Figure (86):	49-year-old man with shoulder adhesive capsulitis	
<b>Figure (87):</b>	Adhesive capsulitis of right sho	
	year old male (a) Transverse co	olor Doppler
	sonography scan. (b) Transverse	sonography
	scan	129
<b>Figure (88):</b>	MRI of shoulder osteoarthritis	130
Figure (89):	Band like hypointense zones oblique T1-weighted images	
<b>Figure (90):</b>	MRI in shoulder of rheumator	
Figure (91):	Ultrasound image of the left stransverse scan showed hum	shoulder in neral head

Fig. No.	Title	Page No.
Figure (92):	Coronal oblique T1WI through shoulder reveals a large tophaceo of the greater tuberosity (red arro	us erosion ow) in this
Figure (93):	patient with goutUltrasound of the right	
rigure (00)	demonstrating double contour sig	
Figure (94):	Grade III, moderate acromioclavic arthritis (ACJ) (encircled) abn demonstrating subacromial fat et	cular joint ormalities
	joint space narrowing, and irregul	
<b>Figure (95):</b>	Acromioclavicular joint, longitudi	nal view 136
<b>Figure (96):</b>	Coronal PD WI shows complete	disruption
	of acromioclavicular ligaments	
<b>Figure (97):</b>	Acromioclavicular joint, longitudi	nal view 139
<b>Figure (98):</b>	Early MRI changes in distal	
	osteolysis	
<b>Figure (99):</b>	Posttraumatic osteolysis of the cla	
<b>Figure (100):</b>	High signal in the LHBT and	
	tendon on a sagittal oblique im	_
	arrow), tendinopathy	
<b>Figure (101):</b>	LHBT, transverse view	
Figure (102):	An oblique sagittal image	
Figure (103):	LHBT, transverse view	
Figure (104):	Septic arthritis of the right should	
Figure (105):	Posterior longitudinal scan of	
<b>7.</b> (4.00)	joint	146
<b>Figure (106):</b>	MR images of the left shoulder A	
71 (10=)	with fat suppression	
Figure (107):	Hill-Sachs lesion	
Figure (108):	Subscapular fossa, longitudinal vi	
Figure (109):	Gender distribution among the	
	patients	154

Fig. No.	Title	Page No.
Figure (110):	Shows Percent of Supraspinator between MRI & US	
Figure (111):	Shows Percent of other rotator con injury between MRI & US	uff muscles
<b>Figure</b> (112):	Shows Percent of ACJ diseas MRI & US	e between
Figure (113):	Shows percent of bursal a between MRI & US	bnormality 160
Figure (114):	Relation between results of remuscle tendinopathy by US and	MR 161
Figure (115):	Relation between results of thickness (bursal) by US and MR Relation between results of	R 162
Figure (116): Figure (117):	thickness (articular) by US and I Relation between results of	MR 163
Figure (118):	thickness (interstitial) by US and Relation between results of full	d MR 164
Figure (119):	(incomplete) by US and MR Relation between results of full	l thickness
Figure (120):	(complete) by US and MRRelation between results of oth	ner rotator
Figure (121):	cuff muscles by US and MRRelation between results of ACJ US and MR	disease by
Figure (122):	Relation between results of burby US and MR	sal disease
Figure (123): Figure (124):		mild joint in the 2 weighted hal at the braspinatus ted image changes of nt with
	impingement over the supraspinatous tendon	174