

سامية محمد مصطفى



شبكة المعلومات الجامعية

بسم الله الرحمن الرحيم



سامية محمد مصطفى



شبكة المعلومات الجامعية



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



سامية محمد مصطفى



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم


قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



سامية محمد مصطفى



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة




سامية محمد مصطفى



شبكة المعلومات الجامعية



بالرسالة صفحات لم ترد بالأصل



REDO SURGERY FOR LOWER LIMB SALVAGE

Essay

*Submitted for partial fulfilment of
Master Degree (M.Sc.)*

In

General Surgery

Presented by

Yousef Abdalla Yackoub

M.B., B.Ch.

Cairo University

Under supervision of

Professor Dr. Abd El-Kader Koth

Professor of General Surgery

Faculty of Medicine - Cairo University

Assistant Professor Dr. Alaa Abd El-Haleem

Assist professor of General Surgery

Faculty of Medicine - Cairo University

Faculty of Medicine

Cairo University

2004

B

١٤٠٤٢

THE NEW YORK PUBLIC LIBRARY

اجتماع لجنة الحكم على الرسالة المقدمة من
الطبيب / الشيخ محمد الراجحي
توطئة للحصول على درجة الماجستير / الدكتوراه
في العلوم الشرعية

Redo surgery for lower limb salvage

باللغة العربية :
الاجابات المأثورة للإشكالات المطروحة

عن المشرفين

۳. تاریخ / آریزن / سرحدی (مابین زمین و شمس) امتحان خارجی

بكالية الطب - جامعة القاهرة وذلك لمناقشة الطالب فى جلسة علنية فى موضوع الرسالة والنتائج


قرار اللجنة :

10/10/10

.....

W. H. L.

عَصَا



Abstract

Failure arterial grafts constitute a problem for vascular surgery, it is an inevitable consequence of arterial reconstructions.

Graft failure classified as early <30 days or late more than 30 days.

Early failure is generally attributed to technical errors, poor patient selection, persistent underlying disease or surface thrombogenicity of the graft. Other causes include: flow inadequate to sustain patency in the conduit used during a period of increased thrombogenicity, a condition that occurs only with PTFE and other prosthetic grafts but rarely with a vein graft, and hypercoagulable states. Thromboembolic graft occlusions not only occur early but are primarily an intraoperative event.

Key Words:


- Redo Surgery
- Lower Limb Salvage

the following table

Acknowledgement

I am greatly honored to work under the supervision of *Professor Dr. Abdel Kader Kotb* professor of general surgery, Cairo University. He has given me much of his time, his guidance, advice and scientific attitude which helps to make this work come to light. No words can be sufficient to express my deep gratitude to him.

I wish to express my great appreciation and sincere thanks to *Dr. Alaa Abd El Haleem*, Assistant Professor of general surgery, Cairo University, for his valuable suggestions and continuous support which make the accomplishment of this work possible, and for the time he spent for evaluating and discussing this work, his valuable remarks will add to the depth of this work.



Introduction

1

1000000

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1