

بسم الله الرحمن الرحيم





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بعض الوثائق الأصلية تالفة





بالرسالة صفحات
لم ترد بالأصل



**Incidence of intradialytic hypertension in
hemodialysis patients in Beheira governorate
and effect of high flux versus low flux
dialysis on it**

Thesis

Submitted for partial fulfilment of Master degree in
Internal Medicine

By

Ahmed Moustafa Abdalla Elgendy

M.B.B.CH, Faculty of Medicine- Alexandria University

Under Supervision of

Prof. Dr. Gamal El-Sayed Mady

Professor of Internal Medicine and Nephrology
Faculty of Medicine – Ain Shams University

Prof. Dr. Hisham Atef Abou El Leil

Professor of Internal Medicine and Nephrology
Faculty of Medicine – Ain Shams University

Dr. Ahmed Mohamed Tawfik

Lecturer of Internal Medicine and Nephrology
Faculty of Medicine – Ain Shams University

Faculty of Medicine
Ain Shams University

2020



Acknowledgments

*First and foremost, I feel always indebted to **Allah**, the **Most Beneficent** and **Merciful** who gave me the strength to accomplish this work,*

*My deepest gratitude to my supervisor, **Prof. Dr. Gamal El-Sayed Mady**, Professor of Internal Medicine and Nephrology, Faculty of Medicine – Ain Shams University, for his valuable guidance and expert supervision, in addition to his great deal of support and encouragement. I really have the honor to complete this work under his supervision.*

*I would like to express my great and deep appreciation and thanks to **Prof. Dr. Hisham Atef Abou El Leil**, Professor of Internal Medicine and Nephrology, Faculty of Medicine – Ain Shams University, for his meticulous supervision, and his patience in reviewing and correcting this work,*

*I must express my deepest thanks to Dr. **Ahmed Mohamed Tawfik**, Lecturer of Internal Medicine and Nephrology, Faculty of Medicine – Ain Shams University for guiding me throughout this work and for granting me much of his time. I greatly appreciate his efforts.*

*Special thanks to my all members of my Family, specially **Mother** and my **Wife**, for their continuous encouragement, enduring me and standing by me.*

✍ Ahmed Moustafa Abdalla Elgendy

List of Contents

<i>Subject</i>	<i>Page No.</i>
List of Abbreviations.....	i
List of Tables	iv
List of Figures.....	vii
Introduction.....	1
Aim of the Work.....	8
Review of Literature	
Chapter (1): Intradialytic hypertension	9
Chapter (2): Dialysis Membranes	41
Chapter (3): High-Flux Hemodialysis	65
Patients and Methods.....	91
Results.....	96
Discussion	112
Summary and Conclusion.....	119
Recommendations	124
References.....	125
Arabic Summary	—

List of Abbreviations

<i>Abbr.</i>	<i>Full-term</i>
ACEI	: Angiotensin-converting enzyme inhibitor
ARBs	: Angiotensin-receptor blockers
AVF	: Arterio-venous fistula
AVG	: Arterio-venous graft
BMD	: Bone mineral density
BMI	: Body mass index
BP	: Blood pressure
BUN	: Blood urea nitrogen
Ca	: Calcium
CKD	: Chronic kidney disease
CLIMB	: Crit-Line Intradialytic Monitoring Benefit
CRP	: C-reactive protein
CV	: Cardiovascular
CVVHD	: Continuous venovenous hemodialysis
CyC	: Cystatin-C
DRA	: Dialysis-Related Amyloidosis
EBPG	: European Best Practice Guidelines
ECV	: Extracellular volume
EPCs	: Endothelial progenitor cells
EPO	: Erythropoietin
ESRD	: End stage renal disease
ET-1	: Endothelin-1
EVAL	: Ethylene vinyl-alcohol
FGF-23	: Fibroblast growth factor-23
FMD	: Flow-mediated vasodilation
GPIHBP1	: Glycosylpho-sphatidylinositol - anchored binding protein 1

HCO	: High cut-off
HD	: Hemodialysis
HDL	: High-density lipoprotein
HDx	: Expanded HD therapy
HEMO	: Hemodialysis
HF	: High-flux
HFHD	: High flux hemodialysis
HLA	: Human leukocyte antigen
HPM	: High performance membrane
HTN	: Hypertension
IDH	: Intradialytic hypertension
IDWG	: Interdialytic weight gain
IPTH	: Intact parathyroid hormone
JSDT	: Japanese Society of Dialysis Therapy
Kuf	: Ultrafiltration coefficient
LDL	: Low-density lipoprotein
LF	: Low-flux
LFHD	: Low flux hemodialysis
LPL	: Lipoprotein lipase
LVM	: Left ventricular mass
MAP	: Mean arterial blood pressure
MCO	: Medium cut-off membranes
MPO	: Membrane Outcomes Permeability
NO	: Nitric oxide
NT-proBNP	: N-terminal-proBNP
PAN	: Polyacrylonitrile-based
PD	: Peritoneal dialysis
PEPA	: Polyester–Polymer–Alloy
PMMA	: Polymethyl methacrylate
Po4	: Phosphorus

PS	: Phosphatidylserine
PSf	: Polysulfone-based membranes
PSGL-1	: P-selectin glycoprotein ligand-1
PVP	: Polyvinylpyrrolidone
Qb	: Blood flow rate
Qd	: Dialysate flow rate
RAAS	: Renin-angiotensin-aldosterone system
RCT	: Randomized clinical trial
ROS	: Reactive oxygen species
S	: Sieving coefficient
SBP	: Systolic blood pressure
SCE	: Saponified cellulose ester
TNF- α	: Tumor necrosis factor- α
TnT	: Troponin T
UFR	: Ultrafiltration rate
URR	: Urea reduction ratio
β2M	: Beta 2 microglobulin

List of Tables

Table No.	Title	Page No.
Table (1):	Definitions of IDH from clinical studies and reviews Hypertension.....	11
Table (2):	Prevalence of Intradialytic Hypertension Among Hemodialysis Patients.....	13
Table (3):	Prospective Observational Studies Associating Intradialytic Hypertension With Increased Risk of Mortality.....	16
Table (4):	Antihypertensive medications and extent of drug removal with HD	28
Table (5):	Classification of dialysis membranes based on biocompatibility. The different permeabilities are highlighted	53
Table (6):	Classification of solutes molecular weight.....	59
Table (7):	Classification of dialyzers based on dialysis membrane permeability.....	61
Table (8):	Dialyzer classification	63
Table (9):	Comparison between the studied groups according to effect on IDH	96
Table (10):	Comparison between the studied groups according to demographic data.....	97
Table (11):	Comparison between both groups as regards the etiology of ESRD	98
Table (12):	Comparison between the studied groups according to HD duration.....	98

Table (13):	Comparison between both groups according to the type of vascular access	99
Table (14):	Comparison between both groups according to anti-hypertensive drugs used:	99
Table (15):	Comparison between both groups according to the hemoglobin level:	100
Table (16):	Comparison between both groups as regards blood urea nitrogen (BUN) and creatinine levels:	101
Table (17):	Comparison between both groups as regards serum electrolytes levels:	102
Table (18):	Comparison between both groups according to IPTH levels	103
Table (19):	Comparison between the studied groups according to lipid profile	104
Table (20):	Comparison between the studied groups regarding Kt/V	105
Table (21):	Comparison between the studied groups regarding urea reduction ratio	105
Table (22):	Comparison between the studied groups according to interdialytic weight gain	106
Table (23):	Comparison between the studied groups according to ultrafiltration rate	106
Table (24):	Comparison between the studied groups according to body mass index	107
Table (25):	Comparison between the studied groups according to ECHO findings	107

Table (26):	Laboratory markers and adequacy of dialysis at the start and at the end of the study in High flux patients:.....	108
Table (27):	Laboratory markers and adequacy of dialysis at the start and at the end of the study in Low flux group:	110

List of Figures

<i>Figure No.</i>	<i>Title</i>	<i>Page No.</i>
Figure (1):	Blood and dialysis solution flow pathways through a hollow fiber and parallel-plate dialyzer	43
Figure (2):	Cross Section of a Dialyzer	44