

بسم الله الرحمن الرحيم



-Call 4000





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة يعبدا عن الغبار













بالرسالة صفحات لم ترد بالأصل





"Possible Cardioprotective Effect of Olive Oil in Experimentally Induced **Hyperthyroidism in Rats**"

A thesis

Submitted for partial fulfillment of Master Degree in Physiology

By

Eman Mohammed Ali Mousa

M.B., B.Ch. Physiology

Supervisors

Prof. Dr. Nermine Kamal Mohamed Saleh Daoud

Professor of Physiology Head of Physiology Department Faculty of Medicine, Ain Shams University

Dr. Abd El- Hamid A Mohamed

Assistant professor of Physiology Faculty of Medicine, Ain shams University

Dr. Manal Said Abd el Hamid

Assistant professor of Physiology Faculty of Medicine, Ain shams University

Dr. Doaa Ahmed Abou-bakr Darwish

Lecturer of Physiology Faculty of Medicine, Ain shams University Physiology Department Faculty of Medicine- Ain Shams University

2020



سورة البقرة الآية: ٣٢

Acknowledgments

First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.

I wish to express my deepest thanks, gratitude and appreciation to **Prof. Dr. Mermine Kamal Mohamed Saleh Daoud**, Professor of Physiology, Head of Physiology Department, Faculty of Medicine, Ain Shams University, for her meticulous supervision, kind guidance, valuable instructions and generous help.

Special thanks are due to **Dr. Abd El- Hamid A**Mohamed, Assistant Professor of Physiology, Faculty of

Medicine, Ain shams University, for his sincere efforts, fruitful
encouragement.

I am deeply thankful to **Dr. Manal Said Abd el Hamid**, Assistant professor of Physiology, Faculty of Medicine, Ain shams University, for her great help, outstanding support, active participation and guidance.

Really I can hardly find the words to express my gratitude to **Dr. Doaa Ahmed Abou-bake Darwish,** Lecturer of Physiology, Faculty of Medicine, Ain shams University, for her supervision, continuous help, encouragement throughout this work.

Thanks to **Dr. Hanan A. Saleh**, Professor of Histology, Faculty of Medicine, Ain Shams University and **Dr. Nermeen Tayseer**, Professor of Clinical Pathology, Faculty of Medicine, Ain Shams University.

I would like to express my hearty thanks to all my family for their support till this work was completed, special thanks Eng. Mahmoud Saad.

Eman Mohammed Ali Mousa

List of Contents

Title	Page No.
List of Tables	i
List of Figures	xi
List of Abbreviations	xvi
Introduction	1
Aim of the Work	3
Review of Literature	4
Materials and Methods	26
Results	49
Discussion	116
Summary and Conclusion	139
References	146
Appendix	180
Arabic Summary	

List of Tables

Table No	. Title	Page No.
Table (1):	Mean ± SEM of initial body weight (IBV body length (IBL), initial body mass indefinal body weight (FBW), final body length final body mass index (FBMI), body percent change (BW%) and body material percent change (BMI%) in all studied groups.	ex (IBM), th (FBL), weight ss index
Table (2):	Mean±SEM values of absolute and weights of right ventricle (RV, RV/E ventricle (LV, LV/BW) and whole hea WH/BW) in all studied groups	relative BW), left art (WH,
Table (3):	Mean±SEM of electrocardiographic parameters	meters in
Table (4):	all studied groups	maximal changes ting rate
Table (5):	groups Mean ±SEM of peak developed tension Baseline values, maximal respon isoproterenol infusion, delta changes and values of perfused hearts isolated from al	(PT; g), ases to recovery
Table (6):	groups	maximal changes nsion/left
Table (7):	perfused hearts isolated from all studied mean±SEM of baseline values, responses to isoproterenol infusion, delta and recovery values of time to peak tensimsec) of perfused hearts isolated from all	maximal changes ion (TPT; l studied
Table (8):	groups Mean±SEM of baseline values, responses to isoproterenol infusion, delta and recovery values of tension generation time (g/msec) of perfused hearts isolated studied groups	changes per unit from all

Table No	. Title	Page No.
Table (9):	Mean±SEM of baseline values, maximal re isoproterenol infusion, delta changes and values of half relaxation time (HRT;	drecovery
Table (10):	perfused hearts isolated from all studied gr Mean±SEM of baseline values, responses to isoproterenol infusion, delta and recovery values of coronary flow raml/min), of perfused hearts isolated	oups
Table (11):	studied groups	71 maximal a changes rate/ left
Table (12):	ventricular weight (CFR/LVW; ml/min/1 perfused hearts isolated from all studied Mean values ± SEM of heart beating rat baseline values, recovery value reperfusion absolute and percent change	groups
Table (13):	perfused hearts isolated from all studied Mean values ± SEM of peak developed to baseline values, recovery values, and re absolute and percent change values of hearts isolated from all studied groups	ension (g), eperfusion perfused
Table (14):	Mean values ± SEM of peak developed to left ventricular weight (g/100mg), at values, recovery values, and reperfusion and percent change values of perfuse isolated from all studied groups	ension per baseline a absolute ed hearts
Table (15):	Mean values ± SEM of time to peal (msec), at baseline values, recovery vareperfusion absolute and percent change perfused hearts isolated from all studied	k tension lues, and values of
Table (16):	Mean values ± SEM of tension gener unit time (g/msec), at baseline values, values, and reperfusion absolute and change values of perfused hearts isolates studied groups	ration per recovery l percent d from all

Table No	o. Title	Page No.
Table (17):	Mean values ± SEM of half relaxation at baseline values, recovery v reperfusion absolute and percent char perfused hearts isolated from all study	values, and nge values of
Table (18):	Mean values ± SEM of coronary (ml/min), at baseline values, recovery reperfusion absolute and percent charperfused hearts isolated from all study	y flow rate y values, and nge values of
Table (19):	Mean values ± SEM of coronary flow ventricular weight (ml/min/100mg), values, recovery values, reperfusion percent change values of perfused he from all studied groups	at baseline absolute and earts isolated
Table (20):	Mean±SEM of serum tri-iodothyronin iodothyronin (T4), thyroid Stimulati (TSH) and cardiac Hemoxygenase end of all studied groups	ing hormone zyme-I (HO-I)
Table (21):	Initial body weight (IBW), initial (IBL), initial body mass index (IBM weight (FBW), final body length (FBI mass index (FBMI), body weight per (BW%) and body mass index per (BMI%) in control group	body length (), final body (L), final body (reent change) (cent change
Table (22):	Initial body weight (IBW), initial (IBL), initial body mass index (IBM weight (FBW), final body length (FBI mass index (FBMI), body weight per (BW%) and body mass index per (BMI%) in hyperthyroid group	body length (), final body (L), final body (reent change cent change
Table (23):	Initial body weight (IBW), initial (IBL), initial body mass index (IBM) weight (FBW), final body length (FBI mass index (FBMI), body weight per (BW%) and body mass index per (BMI%) in olive oil-treated hyperthyre	body length (), final body (L), final body (reent change cent change

Table No	. Title	Page No.
Table (24):	Absolute and relative weights of rig (RV, RV/BW), left ventricle (LV, I whole heart (WH, WH/BW) in control :	LV/BW) and
Table (25):	Absolute and relative weights of rig (RV, RV/BW), left ventricle (LV, I whole heart (WH, WH/BW) in hyperth	tht ventricle LV/BW) and
Table (26):	Absolute and relative weights of rig (RV, RV/BW), left ventricle (LV, I whole heart (WH, WH/BW) in olive hyperthyroid group	tht ventricle LV/BW) and e oil-treated
Table (27):	Electrocardiographic parameters in co	
Table (28):	Electrocardiographic parameters in h	nyperthyroid
Table (29):	Electrocardiographic parameters in treated hyperthyroid group	olive oil-
Table (30):	Effect of graded Isoproterenol (ISO) spontaneous beating rate (bpm) of per isolated from control group	infusion on fused hearts
Table (31):	Effect of graded Isoproterenol (ISO) spontaneous beating rate (bpm) of per isolated from hyperthyroid group	infusion on fused hearts
Table (32):	Effect of graded Isoproterenol (ISO) spontaneous beating rate (bpm) of per isolated from olive oil-treated hyperthy	infusion on fused hearts
Table (33):	Effect of graded Isoproterenol (ISO) peak developed tension (PT; g) of per	infusion on fused hearts
Table (34):	isolated from control group Effect of graded Isoproterenol (ISO) peak developed tension (PT; g) of per- isolated from hyperthyroid rats	infusion on fused hearts
Table (35):	Effect of graded Isoproterenol (ISO) peak developed tension (PT; g) of perisolated from olive oil-treated hyperthy	infusion on fused hearts

Table No	Title	Page No.
Table (36):	Effect of graded Isoproterenol peak developed tension/left (PT/LVW; g/100mg) of perfus from control group	ventricular weight sed hearts isolated
Table (37):	Effect of graded Isoproterenol peak developed tension/left (PT/LVW; g/100mg) of perfus from hyperthyroid group	l (ISO) infusion on ventricular weight sed hearts isolated
Table (38):	Effect of graded Isoproterenol peak developed tension/left (PT/LVW; g/100mg) of perfus from olive oil-treated hyperthy.	l (ISO) infusion on ventricular weight sed hearts isolated
Table (39):	Effect of graded Isoproterenol time to peak tension (TPT; hearts isolated from control gro	(ISO) infusion on msec) of perfused
Table (40):	Effect of graded Isoproterenol time to peak tension (TPT; hearts isolated from hyperthyr	I (ISO) infusion on msec) of perfused
Table (41):	Effect of graded Isoproterenol time to peak tension (TPT; hearts isolated from olive oil-tr group	I (ISO) infusion on msec) of perfused reated hyperthyroid
Table (42):	Effect of graded Isoproterenol tension generation per unit perfused hearts isolated from c	I (ISO) infusion on time (g/msec) of
Table (43):	Effect of graded Isoproterenol tension generation per unit perfused hearts isolated from h	I (ISO) infusion on time (g/msec) of
Table (44):	Effect of graded Isoproterenol tension generation per unit perfused hearts isolated from	I (ISO) infusion on time (g/msec) of m olive oil-treated
Table (45):	hyperthyroid group Effect of graded Isoproterenol half relaxation time (HRT; hearts isolated from control gro	(ISO) infusion on msec) of perfused

Table No	o. Title Page No.
Table (46):	Effect of graded Isoproterenol (ISO) infusion on half relaxation time (HRT; msec) of perfused hearts isolated from hyperthyroid group
Table (47):	Effect ofgraded Isoproterenol (ISO) infusion on half relaxation time (HRT; msec) of perfused hearts isolated from olive oil-treated hyperthyroid
Table (48):	group
Table (49):	Effect of graded Isoproterenol (ISO) infusion on coronary flow rate (CFR; ml/min) of perfused hearts isolated from hyperthyroid group
Table (50):	Effect of graded Isoproterenol (ISO) infusion on coronary flow rate (CFR; ml/min) of perfused hearts isolated from olive oil-treated hyperthyroid
Table (51):	group
Table (52):	Effect of graded Isoproterenol (ISO) infusion on coronary flow rate/ left ventricular weight (CFR/LVW; ml/min/100mg) of perfused hearts isolated from hyperthyroid group
Table (53):	Effect of graded Isoproterenol (ISO) infusion on coronary flow rate/ left ventricular weight (CFR/LVW; ml/min/100mg) of perfused hearts isolated from olive oil-treated hyperthyroid group 212
Table (54):	Baseline values, recovery values, absolute reperfusion values, percent change from baseline values, and percent change from recovery values of heart beating rate (bpm) of hearts isolated from control group