

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

## بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

## جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY



# Value of Neutrophil to Lymphocyte ratio in Prognosis of Elderly Patients with Community Acquired Pneumonia Compared to CRP Level

A Systematic Review/Meta-Analysis

Submitted for Partial Fulfillment of Master Degree in **Chest Diseases** 

By

Amal Mordi Shehata M.B.B.Ch. Ain-Shams University

Under Supervision of

#### **Prof. Gehan Mohamed Elassal**

Professor of Chest Diseases Faculty of Medicine – Ain Shams University

#### **Prof. Mohamed Ali Elsayed**

Professor of Chest Diseases
Faculty of Medicine – Ain Shams University

Faculty of Medicine - Ain Shams University
2020



سورة البقرة الآية: ٣٢

#### Acknowledgments

First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.

I wish to express my deepest thanks, gratitude and appreciation to **Prof. Gehan Mohamed Elassal**, Professor of Chest Diseases, Faculty of Medicine, Ain Shams University, for her meticulous supervision, kind guidance, valuable instructions and generous help.

Special thanks are due to **Prof. Mohamed**Ali Elsayed, Professor of Chest Diseases, Faculty
of Medicine, Ain Shams University, for his sincere
efforts, fruitful encouragement.

I would like to express my hearty thanks to all my family & my colleagues for their support till this work was completed.

Amal Mordi Shehata

#### Tist of Contents

Title	Page No.
List of Tables	5
List of Figures	6
List of Abbreviations	11
Introduction	1 -
Aim of the Work	4
Review of Literature	
■ Pneumonia	5
■ Inflammatory Biomarkers	53
Patients and Methods	72
Results	79
Discussion	108
Summary and Conclusion	115
Recommendations	117
References	118
Arabic Summary	

#### Tist of Tables

Table No	. Title	Page	No.
Table 1:	Demographic characteristics comorbidities of the studied cases		79
Table 2:	Clinical data and severity by CURB 65 of the studied cases	score	
Table 3:	Laboratory data of the studied case admission	es on	
Table 4:	Laboratory data of the studied cases two weeks from admission:	after	
Table 5:	Outcome of the studied cases		86
Table 6:	Correlation between demographic dat	•	
	morbidities and outcome of the studied		87
Table 7:	Correlation between clinical data, CUI		
	score and outcome among studied cases		89
Table 8:	Correlation between NLR, TLC, and Cl		
<b></b>	admission and outcome of the studied c		91
Table 9:	Correlation between NLR, TLC, and		
	after two weeks from admission		0.4
Table 10	outcome of the studied cases		94
Table 10:	<u> </u>		
	among all cases on admission and afterweeks of admission		Ω7
Tahla 11.	Comparison between NLR, TLC, and		91
Table II;	among cases that improved after		
	hospital stay on admission and after		
	weeks of admission		100
Table 12:	Comparison between NLR, TLC, and		
	among cases with long hospital sta		
	admission and after two weeks of admis	•	103
Table 13:	Comparison between NLR, TLC, and	CRP	
	among died cases on admission and	after	
	two weeks of admission		
Table 14:	Role of NLR in prognosis of CAP cases.		107

### List of Figures

Fig. No.	Title	Page	No.
Figure 1:	Radiographic images in a patient with upper lobe pneumonia.	_	17
Figure 2:	Patient with pneumococcal pneumonia. chest CT shows a left upper lobe opextending to the periphery	acity	18
Figure 3:	Image in a 49-year-old woman pneumococcal pneumonia		25
Figure 4:	Image in a 48-year-old patient Haemophilus influenzae pneumonia		26
Figure 5:	Bilateral, multilobar consolidations community-acquired legionella pneum led to acute respiratory failure	nonia	27
Figure 6:	Chest radiograph illustrating "r		28
Figure 7:	Nosocomial left lower-lobe pneumon polymicrobial etiology developed i patient recovering from total colectomy	in a	31
Figure 8:	Left lower-lobe cavitation in a patient nosocomial Gram-negative (Pseudom pneumonia.	onas)	35
Figure 9:	Non-resolving pneumonia investigate computed tomography scan bronchoscopy turned out to bronchoalveolar carcinoma	d by and be	
Figure 10:	Interpretation of NLR		69
Figure 11:	Distribution of co-morbidities as studied cases	_	80

Fig. No.	Title	Page No.
Figure 12:	Shows history of previous pneumamong studied cases	
Figure 13:	Shows CURB-65 scoring among cases.	82
Figure 14:	Shows radiological findings among st	
Figure 15:	Shows outcome of the studied cases	86
Figure 16:	Shows mean age among studied cases different outcomes	
Figure 17:	Shows no significant correlation be gender and outcome among studied case	
Figure 18:	Show significant correlation be radiological pattern of pneumonia outcome of the studied cases	and
Figure 19:	Shows mean TLC and neutrophil country admission among cases with difficuttomes	ferent
Figure 20:	Shows mean of lymphocytic cour admission among cases	

Fig. No.	Title	Page No.
Figure 21:	Shows a highly significant corbetween NLR and outcome of studie the median value of NLR was significant than among patients who died in than among those who improved aft hospital stay.	d cases, ificantly hospital er short
Figure 22:	Shows mean of TLC and neutroph after 2 weeks of admission amon with different outcomes	g cases
Figure 23:	Shows mean of lymphocytic count weeks of admission among cases	
Figure 24:	Shows highly significant corbetween NLR and outcome of studie NLR after 2 weeks of admission from 1.5 to 9.4 in improved case median of 5.45, ranged from 10 cases with long hospital stay with m 12.1 and ranged from 18 to 63 in die with median of 25.	d cases, ranged es with to16 in edian of ed cases
Figure 25:	Shows highly significant difference neutrophils and lymphocytes on ad and after 2 weeks	mission

Fig. No.	Title	Page No.
Figure 26:	Shows significant difference in C admission and after 2 weeks amo cases, ranged from 24 to 194 with me 70 on admission and from 6 to 35 median of 24 after 2 weeks of admissi	ong all dian of 0 with
Figure 27:	Shows significant difference in N admission and after 2 weeks amore cases, ranged from 3.9 to 60 with me 15.75 on admission and from 1.5 to 6 median of 12.1 after 2 weeks of admission and states.	ong all dian of 33 with
Figure 28:	Shows highly significant difference and neutrophil count on admission after 2 weeks among improved cases.	on and
Figure 29:	Shows significant difference in N admission and after 2 weeks improved cases.	among
Figure 30:	Shows a highly significant differe CRP on admission and after 2 weeks improved cases	among
Figure 31:	Shows highly significant difference is neutrophil and lymphocyte court admission and after 2 weeks among with long hospital stay	nts on g cases
Figure 32:	Shows significant difference in N admission and after 2 weeks among with long hospital stay	g cases
Figure 33:	Shows highly significant difference and neutrophil count on admission after 2 weeks among died	on and

Fig. No.	Title	Page No.
Figure 34:	Shows no improvement in NLR among survivors (no significant difference in on admission and after 2 weeks among cases).	NLR died
Figure 35:	Shows NLR predicted mortality; no do occurred when NLR is less than mortality was 8.3% when NLR is bet 10 and 20 and mortality was 91.7% NLR is greater than 20	10, ween when

#### Tist of Abbreviations

Abb.	Full term
ABG	Arterial blood gas
	Acquired immune deficiency syndrome
	Acute Physiology and Chronic Health
111 110112	Evaluation
ARDS	Acute respiratory distress syndrome
	American thoracic society
	Bilevel positive airway pressure
<i>bpm</i>	2 0 2
_	British Thoracic Society
	Blood urea nitrogen
<i>C</i>	Celsius
CA-MRSA	Community acquired Methicillin-resistant
	Staphylococcus
<i>CAP</i>	Community-acquired pneumonia
	$ Complete\ blood\ count$
<i>CLD</i>	Chronic liver disease
<i>CMV</i>	Cytomegalovirus
<i>CNS</i>	Central nervous system
	Chronic obstructive pulmonary disease
	Continuous positive airway pressure
<i>CRP</i>	C-reactive protein
	Computed tomography
<i>CURB</i>	Confusion, Urea, Respiratory rate, Blood
	pressure
<i>CXR</i>	
	Disturbed conscious level
dl	
<i>DM</i>	
	Deoxyribonucleic acid
	Epstein-Barr virus
	Ethylene diamine tetra acetic acid
	Erythrocyte Sedimentation Rate
ESRD	End stage renal disease

#### Tist of Abbreviations (cont...)

Fc	Abb.	Full term
FiO2 gram  G-CSF. Granulocyte colony stimulating factor  GI. Gastrointestinal  H influenzae Haemophilus influenzae  HAP Hospital-acquired pneumonia  HCAP Healthcare-associated pneumonia  HIV Human immunodeficiency virus  Hrs. Hours  HS-CRP High sensitivity C-reactive protein  HTN Hypertension  ICU Intensive care unit  IDSA Infectious diseases society of America  IHD Ischemic heart disease  IL Interleukin  IQR Inter quartile range  IV Intravenous  Kg. Kilogram  L. Liter  LCP Lymphocyte count percentage  LDC Leukocyte differential count  M. catarrhalis Moraxella catarrhalis  MDR Multi-drug resistant  mg Milligram  ml. Milliliter  mmHg. Millimeter of mercury  MRSA Methicillin-resistant Staphylococcus  NAMCS National Ambulatory Medical Care Survey  NCP Neutrophil to lymphocyte count ratio	Fc	Fragment crystallizable
g gram G-CSF Granulocyte colony stimulating factor GI Gastrointestinal H influenzae Haemophilus influenzae HAP Hospital-acquired pneumonia HCAP Healthcare-associated pneumonia HIV Human immunodeficiency virus Hrs Hours HS Highly significant HS-CRP High sensitivity C-reactive protein HTN Hypertension ICU Intensive care unit IDSA Infectious diseases society of America IHD Ischemic heart disease IL Interleukin IQR Inter quartile range IV Intravenous Kg Kilogram L Liter LCP Lymphocyte count percentage LDC Leukocyte differential count M. catarrhalis Moraxella catarrhalis MDR Multi-drug resistant mg Milligram ml Milligram ml Milliter mmHg Millimeter of mercury MRSA Methicillin-resistant Staphylococcus NAMCS National Ambulatory Medical Care Survey NCP Neutrophil to lymphocyte count ratio		
G-CSF		
GI		
H influenzae		
HAP Healthcare-associated pneumonia HCAP Healthcare-associated pneumonia HIV Human immunodeficiency virus Hrs. Hours HS Highly significant Hs-CRP High sensitivity C-reactive protein HTN Hypertension ICU Intensive care unit IDSA Infectious diseases society of America IHD Ischemic heart disease IL Interleukin IQR Inter quartile range IV Intravenous Kg Kilogram L Liter LCP Lymphocyte count percentage LDC Leukocyte differential count M. catarrhalis Moraxella catarrhalis MDR Multi-drug resistant mg Milligram ml Milliter mmHg Millimeter of mercury MRSA Methicillin-resistant Staphylococcus NAMCS National Ambulatory Medical Care Survey NCP Neutrophil count percentage NLCR Neutrophil to lymphocyte count ratio		
HCAP. Healthcare-associated pneumonia HIV. Human immunodeficiency virus Hrs. Hours HS. Highly significant Hs-CRP. High sensitivity C-reactive protein HTN. Hypertension ICU. Intensive care unit IDSA. Infectious diseases society of America IHD. Ischemic heart disease IL Interleukin IQR. Inter quartile range IV. Intravenous Kg. Kilogram L. Liter LCP. Lymphocyte count percentage LDC. Leukocyte differential count M. catarrhalis Moraxella catarrhalis MDR Multi-drug resistant mg Milligram ml. Milliter mmHg Millimeter of mercury MRSA Methicillin-resistant Staphylococcus NAMCS. National Ambulatory Medical Care Survey NCP. Neutrophil count percentage NLCR. Neutrophil to lymphocyte count ratio		= '
HIV		
Hrs		
HS		, · · · · ·
Hs-CRP. High sensitivity C-reactive protein HTN. Hypertension ICU Intensive care unit IDSA Infectious diseases society of America IHD Ischemic heart disease IL Interleukin IQR Inter quartile range IV Intravenous Kg Kilogram L Liter LCP Lymphocyte count percentage LDC Leukocyte differential count M. catarrhalis Moraxella catarrhalis MDR Multi-drug resistant mg Milligram ml Milliter mmHg Millimeter of mercury MRSA Methicillin-resistant Staphylococcus NAMCS National Ambulatory Medical Care Survey NCP Neutrophil count percentage NLCR Neutrophil to lymphocyte count ratio		
HTN		
ICU		
IDSA		
IHD		
IL		· · · · · · · · · · · · · · · · · · ·
IQR Inter quartile range   IV Intravenous   Kg Kilogram   L Liter   LCP Lymphocyte count percentage   LDC Leukocyte differential count   M. catarrhalis Moraxella catarrhalis   MDR Multi-drug resistant   mg Milligram   ml Milliliter   mmHg Millimeter of mercury   MRSA Methicillin-resistant Staphylococcus   NAMCS National Ambulatory Medical Care Survey   NCP Neutrophil count percentage   NLCR Neutrophil to lymphocyte count ratio		
IVIntravenous KgKilogram LLiter LCPLymphocyte count percentage LDCLeukocyte differential count M. catarrhalisMoraxella catarrhalis MDRMulti-drug resistant mgMilligram mlMilliliter mmHgMillimeter of mercury MRSAMethicillin-resistant Staphylococcus NAMCSNational Ambulatory Medical Care Survey NCPNeutrophil count percentage NLCRNeutrophil to lymphocyte count ratio		
Kg		
LLiter  LCPLymphocyte count percentage  LDCLeukocyte differential count  M. catarrhalisMoraxella catarrhalis  MDRMulti-drug resistant  mgMilligram  mlMilliliter  mmHgMillimeter of mercury  MRSAMethicillin-resistant Staphylococcus  NAMCSNational Ambulatory Medical Care Survey  NCPNeutrophil count percentage  NLCRNeutrophil to lymphocyte count ratio		
LCPLymphocyte count percentage  LDCLeukocyte differential count  M. catarrhalisMoraxella catarrhalis  MDRMulti-drug resistant  mgMilligram  mlMilliliter  mmHgMillimeter of mercury  MRSAMethicillin-resistant Staphylococcus  NAMCSNational Ambulatory Medical Care Survey  NCPNeutrophil count percentage  NLCRNeutrophil to lymphocyte count ratio	•	_
LDCLeukocyte differential count  M. catarrhalisMoraxella catarrhalis  MDRMulti-drug resistant  mgMilligram  mlMilliliter  mmHgMillimeter of mercury  MRSAMethicillin-resistant Staphylococcus  NAMCSNational Ambulatory Medical Care Survey  NCPNeutrophil count percentage  NLCRNeutrophil to lymphocyte count ratio		
M. catarrhalis		
MDRMulti-drug resistantmgMilligrammlMillilitermmHgMillimeter of mercuryMRSAMethicillin-resistant StaphylococcusNAMCSNational Ambulatory Medical Care SurveyNCPNeutrophil count percentageNLCRNeutrophil to lymphocyte count ratio		* **
mg       Milligram         ml       Milliliter         mmHg       Millimeter of mercury         MRSA       Methicillin-resistant Staphylococcus         NAMCS       National Ambulatory Medical Care Survey         NCP       Neutrophil count percentage         NLCR       Neutrophil to lymphocyte count ratio		
mlMilliliter  mmHgMillimeter of mercury  MRSAMethicillin-resistant Staphylococcus  NAMCSNational Ambulatory Medical Care Survey  NCPNeutrophil count percentage  NLCRNeutrophil to lymphocyte count ratio		
mmHg.Millimeter of mercuryMRSA.Methicillin-resistant StaphylococcusNAMCS.National Ambulatory Medical Care SurveyNCP.Neutrophil count percentageNLCR.Neutrophil to lymphocyte count ratio		
MRSAMethicillin-resistant Staphylococcus NAMCSNational Ambulatory Medical Care Survey NCPNeutrophil count percentage NLCRNeutrophil to lymphocyte count ratio		
NAMCSNational Ambulatory Medical Care Survey NCPNeutrophil count percentage NLCRNeutrophil to lymphocyte count ratio		
NCPNeutrophil count percentage NLCRNeutrophil to lymphocyte count ratio		
NLCRNeutrophil to lymphocyte count ratio		
		- · · · · · · · · · · · · · · · · · · ·