

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY



Role of Nail Fold Capillaroscopy as A Method of Early Detection of Atherosclerosis in Systemic Lupus Erythematosus Patients

Thesis

Submitted For Partial Fulfillment of Master Degree in Rheumotology

By **Huda Taha Hussein** *M.B.B.Ch*

Under Supervision of

Prof. Dr. Hanan Mohamed Farouk

Professor of Internal Medicine and Rheumatology Faculty of Medicine - Ain Shams University


Dr. Rasha Mahmoud Mohamed

Lecturer of Internal Medicine and Rheumatology Faculty of medicine-Ain Shams University

Dr. Fatma Mohammad Aboud

Lecturer of Internal Medicine and Rheumatology Faculty of medicine-Ain Shams University

> Faculty of Medicine Ain Shams University 2020



Acknowledgment

In the name of **Allah**, I would like to begin by confessing that all the favor and thanks to Allah who empowered me and gave me strength to complete this work.

I am profoundly grateful to **Prof. Dr. Hanan Farouk,** Professor of Internal medicine and Rheumatology and Immunology at Faculty of Medicine —Ain Shams University for her valuable help and fruitful support step by step till this thesis is finished.

My deep appreciation to **Dr. Rasha Mahmoud**, Lecturer Internal Medicine and Rheumatology-Ain Shams University for her useful advice and continuous guidance throughout the thesis

I would like to express my deep appreciation, sincere and profound thanks to **Dr. Fatma Aboud**, Lecturer of Internal Medicine and Rheumatology – Ain shams university for being my mentor and being always supportive and full of positive energy.

I would like to express my deepest appreciation and gratitude to my mom and dad for their extreme support in my life, Family and for being the greatest mom and dad someone can have.

And finally my biggest thanks to my Husband and my special daughters for always being by my side, giving me strength to be in the best Form and being the best Husband and daughters.

At last, my deep thanks to the subjects accepting participation in the study.

🖎 Dr. Huda Taha

List of Contents

| Title | Page No. |
|---|-------------|
| List of Tables | i |
| List of Figures | iii |
| List of Abbreviations | vii |
| Introduction | 1 |
| Aim of the Work | 4 |
| Review of Literature | |
| Systemic Lupus Erythematosus | 5 |
| Nail Fold Capillaroscopy | 56 |
| Atherosclerosis in Systemic Lupus Eryth | nematosus74 |
| Patients and Methods | 84 |
| Results | 90 |
| Discussion | 117 |
| Summary | 126 |
| Conclusion | |
| Recommendations | 129 |
| References | 130 |
| Arabic Summary | — |

List of Tables

| Table No. | Title | Page No. |
|-------------------|---|-------------|
| Table (1): | Showing the frequency of the presentations of SLE at the disease of at any point of the disease | onset and |
| Table (2): | Representing the ACR nomenclate case definitions for neuropsychiate syndromes | ric lupus |
| Table (3): | Definition of SLE classification criter | ria28 |
| Table (4): | Showing the different components of score | |
| Table (5): | Describing the normal capillaroscopi | c pattern62 |
| Table (6): | Atherosclerotic risk factors in system erythematosus. Traditional an traditional | d non- |
| Table (7): | Demographic characteristics of the patients. | |
| Table (8): | Disease activity among the studied p | atients 90 |
| Table (9): | Laboratory findings among the patients. | |
| Table (10): | Nail fold capillaroscopy findings are studied patients. | _ |
| Table (11): | Carotid artery findings among the cases | |
| Table (12): | Comparison between atherosclerotic atherosclerotic patients a demographic characteristics | regarding |
| Table (13): | Comparison between atherosclerotic atherosclerotic patients regarding th activity. | e disease |

List of Tables cont...

| Table No. | Title | Page No. |
|-------------|--|-------------|
| Table (14): | Comparison between atherosclerotic atherosclerotic patients regarding l findings. | laboratory |
| Table (15): | Comparison between atherosclerotic atherosclerotic patients regarding fold capillaroscopy findings | the nail |
| Table (16): | Correlation between the IM thick capillaroscopic findings& diseases among the studied patients | s activity |
| Table (17): | Relation according to capillaroscopy regarding carotid IM thickness (mm | |
| Table (18): | Diagnostic performance of no capillaroscopy in diagnosing ather in our SLE patients: | rosclerosis |
| Table (19): | Diagnostic characteristics of a capillaroscopy in diagnosing atheros | |

List of Figures

| Fig. No. | Title | Page No. |
|--------------|--|--------------|
| Figure (1): | Showing the different etiologies of S | SLE8 |
| Figure (2): | Showing the pathophysiology of the | |
| Figure (3): | Classification criteria for systemic erythematosus | - |
| Figure (4): | Highlights EULAR 2019 recommen for treating non-renal SLE | |
| Figure (5): | Normal nailfold capillaries | 58 |
| Figure (6): | The different techniques for perfinailfold capillaroscopy | O |
| Figure (7): | Abnormal capillary shapes | 65 |
| Figure (8): | Avascular areas, demonstrating a capillary drop-out" | |
| Figure (9): | Nailfold capillaroscopy showing a pattern | |
| Figure (10): | Nailfold capillaroscopy image progressively worsening damage | |
| Figure (11): | Desribes Foam cell formation. particles (blue) become oxidized vascular wall | in the |
| Figure (12): | Describes the mechanisms lead atherogenesis and cardiovascular in SLE patients | disease |
| Figure (13): | Disease activity among the patients | |
| Figure (14): | Nail fold capillaroscopy abnormalities among the studied ca | - |
| Figure (15): | Nail fold capillaroscopic hemo among the studied cases | _ |

List of Figures Cont...

| Fig. No. | Title | Page No. |
|--------------|---|---------------------|
| Figure (16): | Nail fold capillaroscopy subpartitions among the studied cases | * |
| Figure (17): | Nail fold capillaroscopy distration among the studied cases | |
| Figure (18): | Atherosclerosis among the patients | |
| Figure (19): | Comparison according to atheros regarding SLEDAI score. | |
| Figure (20): | Comparison according to atheros regarding clinical severity. | |
| Figure (21): | Comparison according to atheros regarding LDL | |
| Figure (22): | Comparison according to atheros regarding ESR. | |
| Figure (23): | Comparison according to atheros regarding Capillary density | |
| Figure (24): | Comparison according to atheros regarding length | |
| Figure (25): | Comparison according to atheros regarding arterial limb diameter | |
| Figure (26): | Comparison according to atheros regarding nail fold shape abnormal | |
| Figure (27): | Nail fold capillaroscopy hemoamong the studied cases | _ |
| Figure (28): | Nail fold capillaroscopy subpaplexus among the studied cases | |
| Figure (29): | Comparison according to atheros regarding nail fold capilla disorganized distribution | clerosis roscopy |

List of Figures Cont...

| Fig. No. | Title | Page No. |
|--------------|---|-----------------|
| Figure (30): | Correlation between IM SLEDAI score | |
| Figure (31): | Correlation between IM density | |
| Figure (32): | Correlation between IM length | |
| Figure (33): | Correlation between IM width | |
| Figure (34): | Correlation between IM arterial limb diameter | |
| Figure (35): | Relation between capillar (distribution) regarding thickness | carotid IM |
| Figure (36): | Relation between capillar (hemorrhage) regarding thickness | carotid IM |
| Figure (37): | Relation of capillarose (corkscrew) regarding thickness | carotid IM |
| Figure (38): | Relation of capillarose (meandering) regarding thickness | carotid IM |
| Figure (39): | Relation of capillarose (tortuous) regarding carotic | copy findings |
| Figure (40): | Relation of capillarose (subpapillary plexus) regar thickness | ding carotid IM |
| Figure (41): | ROC curve for nail fold cadiagnosing atherosclerosis. | |

List of Figures Cont...

| Fig. No. | Title | Page No. |
|--------------|---|---|
| Figure (42): | Diagnostic characteristics of Arter diameter ≥17.0 µm in dia atherosclerosis | gnosing |
| Figure (42): | Derived from real cases in this st showing: (a) Normal capilla findings. (b) Meandering and buc capillaries. (c) Microhemo abnormal distribution of capillar Tortuous and meandering capillari | roscopic dding of orrhages ries. (d) |

List of Abbreviations

| Abb. | Full term |
|--------|---|
| ABC | ATP binding cassette transporter |
| | Autoimmune haemolytic anaemia |
| | Antinuclear antibodies |
| | Anti-double-stranded DNA antibodies |
| | Antigen presenting cells |
| | Anti phospholipid antibodies |
| | Antiphospholipid antibodies |
| | Antiphospholipid syndrome |
| | Atherosclerotic cardiovascular disease |
| AZA | Azathioprine |
| | British Isles Lupus Assessment Group |
| | B lymphocyte stimulator |
| CAC | Coronary artery calcium |
| CHD | Coronary heart disease |
| cIMT | Carotid intima media thickness |
| CLE | Cutaneous lupus erythematosus |
| CMV | Cytomegalovirus |
| CNS | Central nervous system |
| CRP | C-reactive protein |
| CTD | Connective tissue |
| CVD | Cardiovascular disease |
| CYC | Cyclophosphamide |
| DLE | Discoid lupus erythematosus |
| ds-DNA | Anti-double stranded DNA |
| EBV | Epstein Barr virus |
| ECLAM | European Consensus Lupus Activity Measure |
| ECs | Endothelial cells |
| GC | Glucocorticoids |
| GI | Gastrointestinal |

List of Abbreviations cont...

| Abb. | Full term |
|----------|--|
| CHIAC | 0 '1 '4' 4 1' |
| | Genome-wide association studies |
| <u> </u> | Hydroxychloroquine |
| | Interstitial Lung disease |
| IFN | |
| IL | |
| | Intima-media thickness |
| | Immunosuppressive |
| | Intrauterine growth restriction |
| | Intravenous immunoglobulin |
| | Lupus Activity Index |
| | Myocardial infarction |
| | Mycophenolate mofetil |
| | Methyl prednisolone |
| MTX | |
| | Nail fold capillaroscopy |
| | National Institute of Clinical Excellence |
| NK | |
| | Novel oral anticoagulants |
| NPSLE | Neuropsychiatric systemic lupus erythromatosus syndromes |
| PBMCs | Peripheral blood mononuclear cells |
| PNS | Peripheral nervous system |
| PWV | Pulse wave velocity |
| RA | Rheumatoid arthritis |
| RCTS | Randomized controls trials |
| RTX | Rituximab |
| SCLE | Subacute cutaneous lupus erythematosus |
| SIS | SLE Index Score |
| SLAM | Systemic Lupus Activity Measure |