



بسم الله الرحمن الرحيم

∞∞∞∞

تم رفع هذه الرسالة بواسطة / سامية زكى يوسف

بقسم التوثيق الإلكتروني بمركز الشبكات وتكنولوجيا المعلومات دون أدنى

مسئولية عن محتوى هذه الرسالة.

ملاحظات: لا يوجد





# **Role of magnetic resonance imaging in evaluating of non-traumatic chronic ankle pain in adult**

*Thesis*

Submitted for partial fulfillment of Master Degree  
in Radiodiagnosis

*Presented by*

**Kyrillos Soliman Kadous Boutros**

(M.B.B.C.H)

Faculty of Medicine Ain Shams University

*Supervised by*

**Prof. Dr. Sherif Hamed Abou Gamrah**

Professor of Radiology

Faculty of Medicine Ain Shams University

**Dr. Suzan Farouk Ibrahim**

Lecturer of Radiology Faculty of Medicine Ain Shams University

**Faculty of Medicine  
Ain Shams University  
2022**



## Acknowledgments

*First and foremost, I feel always indebted to GOD, the Most Kind and Most Merciful.*

*I'd like to express my respectful thanks and gratitude to **Prof. Dr. Sherif Hamed Abou Gamrah**, Professor of Radiodiagnosis, Faculty of Medicine Ain Shams University, for giving me the honor and great advantage of working under his supervision, valuable teaching and the continuous education.*

*My sincere thanks and utmost appreciation to **Dr. Suzan Farouk Ibrahim**, Lecturer of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for his tremendous assistance, close supervision and continuous advice during different stages of this work.*

*Special thanks to my Parents, my **Wife** and all my **Family** members for their continuous encouragement, enduring me and standing by me.*

*✍ **Kyrillos Soliman Kadous Boutros***

## **List of Contents**

| <i><b>Subject</b></i>                                            | <i><b>Page No.</b></i> |
|------------------------------------------------------------------|------------------------|
| <b>List of Abbreviations.....</b>                                | <b>i</b>               |
| <b>List of Tables.....</b>                                       | <b>iii</b>             |
| <b>List of Figures .....</b>                                     | <b>iv</b>              |
| <b>Introduction .....</b>                                        | <b>1</b>               |
| <b>Aim of the study.....</b>                                     | <b>3</b>               |
| <b>Review of Literature</b>                                      |                        |
| Ankle and Hindfoot Overview.....                                 | 4                      |
| Detailed Anatomy of Ankle Joint .....                            | 7                      |
| MRI Technique of the Ankle Joint .....                           | 43                     |
| Pathogenesis of Non-Traumatic Ankle Joint<br>Injuries .....      | 55                     |
| MRI Manifestations of Non-Traumatic Ankle<br>Joint Injuries..... | 77                     |
| <b>Patients and Methods.....</b>                                 | <b>121</b>             |
| <b>Illustrative Cases .....</b>                                  | <b>122</b>             |
| <b>Results.....</b>                                              | <b>128</b>             |
| <b>Discussion .....</b>                                          | <b>134</b>             |
| <b>Conclusion.....</b>                                           | <b>140</b>             |
| <b>References .....</b>                                          | <b>141</b>             |
| <b>Arabic Summary .....</b>                                      | <b>—</b>               |

---

## List of Abbreviations

| <i>Abbr.</i> | <i>Full-term</i>                          |
|--------------|-------------------------------------------|
| <b>AS</b>    | : Accessory soleus                        |
| <b>ATV</b>   | : Anterior tibial vessels                 |
| <b>AVN</b>   | : Avascular necrosis                      |
| <b>CECS</b>  | : Chronic exertional compartment syndrome |
| <b>DA</b>    | : Deep aponeurosis                        |
| <b>DPN</b>   | : Deep peroneal nerves                    |
| <b>DPN</b>   | : Deep peroneal nerve                     |
| <b>EDL</b>   | : Extensor digitorum longus               |
| <b>EHL</b>   | : Extensor hallucis longus                |
| <b>FDL</b>   | : Flexor digitorum longus                 |
| <b>FHL</b>   | : Flexor hallucis longus                  |
| <b>FR</b>    | : Flexor retinaculum                      |
| <b>FSE</b>   | : Fast spin echo                          |
| <b>GSV</b>   | : Greater saphenous vein                  |
| <b>LSV</b>   | : Lesser saphenous vein                   |
| <b>MRI</b>   | : Magnetic resonance imaging              |
| <b>PB</b>    | : Peroneus brevis                         |
| <b>PCI</b>   | : Peroneocalcaneus internus               |
| <b>PD</b>    | : Proton density                          |
| <b>PL</b>    | : Peroneus longus                         |
| <b>PT</b>    | : Peroneus tertius                        |

|             |   |                              |
|-------------|---|------------------------------|
| <b>PTT</b>  | : | Posterior tibialis tendon    |
| <b>PTV</b>  | : | Posterior tibial vessels     |
| <b>SE</b>   | : | Spin echo                    |
| <b>SI</b>   | : | Signal intensity             |
| <b>STIR</b> | : | Short tau inversion recovery |
| <b>STS</b>  | : | Sinus tarsi syndrome         |
| <b>TA</b>   | : | Tibialis anterior tendon     |
| <b>TN</b>   | : | Tibial nerve                 |
| <b>WIs</b>  | : | Weighted images              |

## **List of Tables**

| <b>Table No.</b>  | <b>Title</b>                                                                                                   | <b>Page No.</b> |
|-------------------|----------------------------------------------------------------------------------------------------------------|-----------------|
| <b>Table (1):</b> | Ankle MR Imaging Protocols at 1.5 T.....                                                                       | 54              |
| <b>Table (2):</b> | Different patients with non-traumatic<br>chronic ankle pain were examined<br>showing diseases prevalence. .... | 130             |

## List of Figures

| <i>Figure No.</i>   | <i>Title</i>                                                                                                                                | <i>Page No.</i> |
|---------------------|---------------------------------------------------------------------------------------------------------------------------------------------|-----------------|
| <b>Figure (1):</b>  | AP view bones of the ankle joint.....                                                                                                       | 8               |
| <b>Figure (2):</b>  | Tarsal tunnel and flexor retinaculum.....                                                                                                   | 12              |
| <b>Figure (3):</b>  | Extensor retinacula .....                                                                                                                   | 12              |
| <b>Figure (4):</b>  | A transverse section through the lower part of the left ankle joint, superior aspect.....                                                   | 14              |
| <b>Figure (5):</b>  | Diagram illustrates calcaneal tendon of the ankle.....                                                                                      | 15              |
| <b>Figure (6):</b>  | Lateral aspect ligament of the ankle joint.....                                                                                             | 17              |
| <b>Figure (7):</b>  | Supramalleolar.....                                                                                                                         | 21              |
| <b>Figure (8):</b>  | MRI coronal plane of the ankle .....                                                                                                        | 26              |
| <b>Figure (9):</b>  | MRI sagittal plane of the ankle.....                                                                                                        | 31              |
| <b>Figure (10):</b> | Illustration of the peroneus quadratus, which may insert on the retrotrochlear eminence (1), cuboid (2), or peroneus longus tendon (3)..... | 33              |
| <b>Figure (11):</b> | 3D rendering demonstrating the peronealcalcaneal variant of the peroneus quartus. ....                                                      | 34              |
| <b>Figure (12):</b> | Peroneus quartus (peronealcalcaneal variant).....                                                                                           | 34              |
| <b>Figure (13):</b> | Axial MR image demonstrating peroneus quadratus medial and posterior to the peroneus brevis and longus tendons.....                         | 35              |

- Figure (14):** 3D rendering demonstrating the peroneocalcaneus internus (PCI) courses lateral to the flexor hallucis longus muscle and tendon (FHL) before inserting below the sustentaculum tali. The posterior tibial (PTT) and flexor digitorum longus (FDL) tendons are also labeled..... 36
- Figure (15):** Axial illustration of the ankle demonstrating four medial tendons instead of the usual three tendons: Tibialis posterior (TP), flexor digitorum longus (FDL), flexor hallucis longus (FHL), and peroneocalcaneus internus (PCI) ..... 36
- Figure (16):** Axial T1-weighted image demonstrating peroneocalcaneus internus; lateral to the flexor digitorum longus ..... 37
- Figure (17):** 3D rendering demonstrating the accessory soleus muscle (AS) located superficial to the deep aponeurosis (DA) and flexor retinaculum (FR)..... 37
- Figure (18):** Accessory soleus muscle. Axial T1-weighted images demonstrating an accessory soleus muscle..... 38
- Figure (19):** Accessory soleus muscle. .... 38
- Figure (20):** 3D illustration of the flexor digitorum accessories longus..... 39
- Figure (21):** Within the distal lower leg, a triangular shaped accessory muscle; abuts the adjacent neurovascular

|                     |                                                                                                                                                                                                                                                                              |    |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
|                     | bundle and lies posterior to the flexor<br>halluces longus muscle (FHL). ....                                                                                                                                                                                                | 40 |
| <b>Figure (22):</b> | Axial (A) and sagittal (B) drawings at<br>the ankle demonstrate an accessory<br>flexor digitorum longus tendon<br>coursing within the posterior flexor<br>compartment as the most superficial<br>posterior structure and causing<br>crowding of the flexor compartment ..... | 40 |
| <b>Figure (23):</b> | The FDAL remains fleshy until just<br>prior to exiting the tarsal tunnel. The<br>muscle is seen posterior to the flexor<br>halluces longus tendon (FHL) .....                                                                                                                | 41 |
| <b>Figure (24):</b> | .....                                                                                                                                                                                                                                                                        | 41 |
| <b>Figure (25):</b> | .....                                                                                                                                                                                                                                                                        | 42 |
| <b>Figure (26):</b> | 8-channel foot and ankle coil.....                                                                                                                                                                                                                                           | 45 |
| <b>Figure (27):</b> | Patient positioned for examination of<br>the foot and ankle using flat <b>(A)</b> and<br>circumferential <b>(B)</b> coils. ....                                                                                                                                              | 46 |
| <b>Figure (28):</b> | Localizer images of ankle joint MRI<br>examination.....                                                                                                                                                                                                                      | 47 |
| <b>Figure (29):</b> | Sagittal plane of ankle joint MRI<br>examination.....                                                                                                                                                                                                                        | 51 |
| <b>Figure (30):</b> | Axial plane of ankle joint MRI<br>examination.....                                                                                                                                                                                                                           | 51 |
| <b>Figure (31):</b> | Coronal plane of ankle joint MRI<br>examination.....                                                                                                                                                                                                                         | 52 |

|                     |                                                                                                                                                                 |    |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| <b>Figure (32):</b> | Diagrammatic drawing of osteochondritis dissecans as seen in the coronal plane.....                                                                             | 57 |
| <b>Figure (33):</b> | Posteromedial impingement, with entrapment of fibrotic scar tissue in the posteromedial ankle gutter (circled). ....                                            | 66 |
| <b>Figure (34):</b> | Posterolateral impingement, with the marsupial meniscus extending intra-articularly between the posterior tibiofibular and posterior talofibular ligaments..... | 67 |
| <b>Figure (35):</b> | Bone contusions.....                                                                                                                                            | 78 |
| <b>Figure (36):</b> | Talar stress injury, Sagittal STIR image showing oedema in the talar tubercles, .....                                                                           | 80 |
| <b>Figure (37):</b> | Stress fracture of the distal tibia, the sagittal T1-weighted GRE sequence.....                                                                                 | 81 |
| <b>Figure (38):</b> | Focal subchondral low signal on T1W/PDW and high signal on FS T2W FSE/STIR, with normal overlying cartilage .....                                               | 82 |
| <b>Figure (39):</b> | Coronal STIR MR image showing a lateral osteochondral lesion of the talus with deficient overlying cartilage .....                                              | 83 |
| <b>Figure (40):</b> | Coronal FS T2W FSE MR Image showing a medial osteochondral lesion of the talus with a thin fluid SI line separating it from the underlying bone,.....           | 83 |

- Figure (41):** (A) Axial T2W FSE MR Image showing a loose osteochondral fragment in the anterior joint recess. Sagittal STIR MR images showing. (B) An oedematous OLT with a hypointense margin and mild surrounding marrow oedema ..... 84
- Figure (42):** A): Unstable talar osteochondral lesion. Sagittal STIR MR image reveals osteochondral lesion with fluid-like signal intensity interposed between the fragment and the donor site ..... 85
- Figure (43):** Avascular necrosis of the talus. Sagittal STIR MR image demonstrates serpentine areas of increased signal intensity in the talus ..... 87
- Figure (44):** Normal Achilles tendon ..... 93
- Figure (45):** Paratenonitis axial FS T2W FSE MR Image showing fluid signal intensity around the tendon ..... 94
- Figure (46):** A- Sagittal PD-weighted fat-sat image shows Achilles tendinosis with advanced internal degeneration and a longitudinal tear. B- Axial PD-weighted fat-sat image shows the partial tear entering the tendon from the medial side, with associated peritendinitis ..... 95
- Figure (47):** A- Sagittal STIR and B- Axial FS T2W FSE MR images complete tear..... 96
- Figure (48):** Chronic complete tear, Sagittal T1W SE MR image showing irregular

|                     |                                                                                                                                                                                                                           |     |
|---------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
|                     | expansion of the distal tendon and scar<br>tissue filling a chronic tear.....                                                                                                                                             | 96  |
| <b>Figure (49):</b> | Ankle tendons MRI axial PD.....                                                                                                                                                                                           | 97  |
| <b>Figure (50):</b> | Posterior tibial (PTT) Acute<br>tenosynovitis: .....                                                                                                                                                                      | 98  |
| <b>Figure (51):</b> | Posterior tibial (PTT) Chronic<br>tenosynovitis .....                                                                                                                                                                     | 99  |
| <b>Figure (52)</b>  | A- Type I tear of the posterior tibial<br>tendon.....                                                                                                                                                                     | 100 |
| <b>Figure (53):</b> | Type II tears of the posterior tibial<br>tendon. Axial proton-density-weighted<br>MR image shows an attenuated<br>posterior tibial tendon with a caliber<br>equal to that of the adjacent flexor<br>digitorum tendon..... | 101 |
| <b>Figure (54):</b> | Type III tears of the posterior tibial<br>tendon.....                                                                                                                                                                     | 101 |
| <b>Figure (55):</b> | Dislocation of the posterior tibial<br>tendon.....                                                                                                                                                                        | 103 |
| <b>Figure (56):</b> | Flexor hallucis longus tenosynovitis.....                                                                                                                                                                                 | 104 |
| <b>Figure (57):</b> | Flexor hallucis longus tendon.....                                                                                                                                                                                        | 105 |
| <b>Figure (58):</b> | Peroneal tenosynovitis. ....                                                                                                                                                                                              | 106 |
| <b>Figure (59):</b> | Longitudinal tear of the peroneus<br>brevis tendon.....                                                                                                                                                                   | 107 |
| <b>Figure (60):</b> | Axial STIR-weighted MR image<br>shows complete tear of anterior<br>talofibular ligament.....                                                                                                                              | 111 |

- Figure (61):** Coronal STIR-weighted MR image shows loss of fatty striations of the PDTL..... 113
- Figure (62):** Fluid in the tibiofibular recess measures approximately, 10 mm with torn anterior and posterior inferior tibiofibular/syndesmotoc ligaments (blue and pink arrows, respectively)..... 114
- Figure (63):** (A) On the lateral of the foot radiogram, the os trigonum with a sclerotic signal close to the talus. (B, C) There is no connection between the os trigonum and the talus in the fat saturated T2W sagittal and axial plane images. The ossicle, which is the signal of medullary edema, is seen as a separate bone structure (arrows). In addition to os trigonum edema, talus and soft edema ..... 116
- Figure (64):** A and B, Coronal T1-weighted (A) and sagittal T1-weighted (B) MR images demonstrate disruption of cervical ligament (CL) (arrow). C, Arthroscopic finding demonstrates partial tear of CL..... 117
- Figure (65):** Sagittal T1 (A) and sagittal T2 (B) and sagittal STIR (C) revealing prominent osseous bump at the posterior superior aspect of the calcaneus..... 122
- Figure (66):** Axial T2 (A) and Sagittal STIR (B) revealing Os trigonum with surrounding soft tissue altered signal displaying high

- STIR signal, localized fluid in the posterior joint recesses and mild tenosynovitis involving the flexor hallucis longus (FHL) denoting os trigonum syndrome..... 122
- Figure (67):** Sagittal T2 (A), Sagittal STIR (B) and sagittal T1 (C) revealing sinus tarsi fat replaced by fluid signal eliciting low T1 and bright T2/STIR signals and thickening Of The Inter-Osseous Ligament consistent with sinus tarsi syndrome..... 123
- Figure (68):** **CASE 3:** Sagittal T1 (A) and Sagittal T2 (B) revealing Osteoarthritic changes of tibio-talar and sub-talar articulation in terms of narrowed joint space, irregularity of articular surfaces with subchondral marrow degenerative changes, multiple osteochondral injures at the opposing tibio-talar articular surfaces surrounded by marrow edema eliciting low T1 and bright T2 signals and associated pseudo-cystic changes and geode formation as well as calcaneal plantar spur evidence of degenerative osteoarthritic changes. .... 123
- Figure (69):** Sagittal STIR (A), Sagittal T2 (B) and sagittal T1 (C) revealing Osteochondral defect at the medial aspect of the talar dome with underlying area of marrow signal