



بسم الله الرحمن الرحيم

∞∞∞∞

تم رفع هذه الرسالة بواسطة / سلوي محمود عقل

بقسم التوثيق الإلكتروني بمركز الشبكات وتكنولوجيا المعلومات دون أدنى

مسئولية عن محتوى هذه الرسالة.

ملاحظات: لا يوجد



**Assessment of the Mandibular Foramen
Location in a Sample of Egyptian
Population Using CBCT
"A Retrospective Study"**

A Thesis

Submitted for Partial Fulfillment of the Master Degree of
Science in Oral and Maxillofacial Radiology,
Faculty of Dentistry, Ain-Shams University

By

Aya ElSayd Abou Efetouh Metwaly

B.D.S. Faculty of Dentistry, Mansoura University, Dakihlya(2013)

Email: abouelfetouh.aya@gmail.com

Mobile: 01000495687

Under Supervision of

Dr. Mary Medhat Farid

Professor of Oral and Maxillofacial Radiology

Faculty of Dentistry - Ain Shams University

Dr. Fatma Mostafa Elbadawy

Lecturer of Oral and Maxillofacial Radiology

Faculty of Dentistry - Ain Shams University

*Faculty of Dentistry
Ain Shams University
2022*

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

سبحانك لا علم لنا
إلا ما علمتنا إنك أنت
العليم العليم

صدق الله العظيم

سورة البقرة الآية: ٣٢

Acknowledgments

*First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.*

*I wish to express my deepest thanks, gratitude and appreciation to **Dr. Mary Medhat Farid**, Professor of Oral and Maxillofacial Radiology, Faculty of Dentistry, Ain Shams University, for his meticulous supervision, kind guidance, valuable instructions and generous help.*

*Special thanks are due to **Dr. Fatma Mostafa Elbadawy**, Lecturer of Oral and Maxillofacial Radiology, Faculty of Dentistry, Ain Shams University, for his sincere efforts, fruitful encouragement.*

I would like to express my hearty thanks to all my family for their support till this work was completed.

Last but not least my sincere thanks and appreciation to all patients participated in this study.

Aya ElSaid Abou Efetouh Metwaly

List of Contents

Title	Page No.
List of Tables.....	i
List of Figures.....	v
List of Abbreviations	xvii
Introduction	1
Review of Literature.....	2
Aim of the Study	36
Patients and Methods.....	37
Case Presentation.....	48
Results.....	53
Discussion	107
Summary.....	120
Conclusion and Recommendations	128
References	130
Arabic Summary.....	١

List of Tables

Table No.	Title	Page No.
Table 1:	Mann-Whitney U test comparing the distance from midpoint of the anterior margin of mandibular foramen fossa to the nearest point of the anterior ramus border in males and females.....	56
Table 2:	Mann-Whitney U test comparing the distance from midpoint of the posterior margin of mandibular foramen fossa to the nearest point of the posterior ramus border in males and females.....	58
Table 3:	Mann-Whitney U test comparing the distance from the anterior to posterior ramus border passing by the midpoint of the mandibular foramen fossa in males and females.....	60
Table 4:	Mann-Whitney U test comparing the distance from the lowest point of the mandibular notch to the entering point of mandibular foramen fossa in males and females.....	62
Table 5:	Mann-Whitney U test comparing the distance from the lowest point of mandibular notch to the most convex point of inferior ramus border passing by the entering point of the mandibular foramen fossa in males and females.....	64
Table 6:	Mann-Whitney U test comparing the depth of the mandibular foramen which is the distance from the point beyond the lingual tip on the inner plate surface of mandibular foramen to symmetrical point on the external plate surface in males and females.	66
Table 7:	Mann-Whitney U test comparing the distance from the midpoint of mandibular foramen to occlusal plane of the mandibular first and second molars in males and females.....	68
Table 8:	Mann-Whitney U test comparing the distance from the apex of retromolar trigone to midpoint of MF in males and females.....	70
Table 9:	Mann-Whitney U test comparing the distance from anterior border of mandibular foramen to posterior border of mandibular foramen passing by entering point of mandibular foramen in males and females.....	72

List of Tables cont...

Table No.	Title	Page No.
Table 10:	Mann-Whitney U test comparing the distance from entering point of mandibular foramen fossa to the point of the greatest convexity of the inferior ramus border in males and females.	74
Table 11:	Summary of comparison of measurements between males and females	76
Table 12:	Mann-Whitney U test comparing the distance from midpoint of the anterior margin of the mandibular foramen fossa to the nearest point of the anterior ramus border in different age groups.	77
Table 13:	Mann-Whitney U test comparing the distance from midpoint of the posterior margin of mandibular foramen to the nearest point of the posterior ramus border in different age group.	78
Table 14:	Mann-Whitney U test comparing the distance between anterior and posterior ramus border passing by the midpoint of the MF fossa in different age groups.....	79
Table 15:	Mann-Whitney U test comparing the distance from the lowest point of the mandibular notch to the entering point of mandibular foramen in different age group.	80
Table 16:	Mann-Whitney U test comparing the distance between mandibular notch and inferior ramus border passing by entering point of mandibular foramen in different age group.	81
Table 17:	Mann-Whitney U test comparing the depth of mandibular foramen Fossa which forms mandibular foramen fossa depression from the point beyond the lingual tip on the inner plate surface to symmetrical point on the external plate surface in different age groups.	82
Table 18:	Mann-Whitney U test comparing the distance from the midpoint of mandibular foramen to occlusal plane of the mandibular first and second molars in different age group.....	83
Table 19:	Mann-Whitney U test comparing the distance from the apex of retromolar trigone to midpoint of mandibular foramen in different age groups.....	84

List of Tables cont...

Table No.	Title	Page No.
Table 20:	Mann-Whitney U test comparing the distance from anterior border of mandibular foramen to posterior border of mandibular foramen in different age group.....	85
Table 21:	Mann-Whitney U test comparing the distance from entering point of mandibular foramen fossa to the point of the greatest convexity of the inferior ramus border in different age group.	86
Table 22:	Summary of comparison of measurements different age groups	87
Table 23:	Mann-Whitney U test comparing the distance from midpoint of the anterior margin of the mandibular foramen fossa to the nearest point of the anterior ramus border according to dental status.	88
Table 24:	Mann-Whitney U test comparing the distance from midpoint of the posterior margin of MF fossa to the nearest point of the posterior ramus border according to dental status.....	90
Table 25:	Mann-Whitney U test comparing the distance between anterior and posterior ramus border passing by the midpoint of the mandibular foramen fossa according to dental status.....	92
Table 26:	Mann-Whitney U test comparing the distance from the lowest point of the mandibular notch to the entering point of mandibular foramen fossa according to dental status.	94
Table 27:	Mann-Whitney U test comparing the distance between mandibular notch and inferior ramus border passing by entering point of mandibular foramen fossa according to dental status.	96
Table 28:	Mann-Whitney U comparing the depth of the mandibular foramen Fossa which forms mandibular foramen fossa depression from the point beyond the lingual tip on the inner plate surface to symmetrical point on the external plate surface according to dental status.	98
Table 29:	Mann-Whitney U comparing the distance from the apex of retromolar trigone to midpoint of mandibular foramen to according to dental status.	100

List of Tables cont...

Table No.	Title	Page No.
Table 30:	Mann-Whitney U test comparing the distance from anterior border of mandibular foramen to posterior border of mandibular foramen passing by midpoint of mandibular foramen according to dental status.	102
Table 31:	Mann-Whitney U test comparing the distance from entering point of mandibular foramen fossa to the point of the greatest convexity of the inferior ramus border according to dental status.	104
Table 32:	Summary of comparison of measurements between edntate and dentate population.	106

List of Figures

Fig. No.	Title	Page No.
Figure 1:	The reference points in the mandibular ramus.....	3
Figure 2:	Medial surface of the right mandibular ramus.....	4
Figure 3:	Common location of the mandibular foramen in adults. ¹⁵	5
Figure 4:	Imaginary line extends from sigmoid notch to inferior border of the mandible	8
Figure 5:	Graphical representation to the changes between the edentate/dentate mandibles	9
Figure 6:	Pathway of mandibular nerve. ¹⁵	11
Figure 7:	Transverse section of the right mandibular ramus at the level of the lingula showing anteroposterior arrangement of IAN and inferior alveolar vasculatures	12
Figure 8:	Diagrammatic representation of the normal anatomy of the mandibular nerve in the infratemporal fossa	14
Figure 9:	Patient's face in horizontal position showing needle aligned with plane from lower border of intertragic notch to corners of the mouth. Needle is directed to target area on tragus. ⁴²	15
Figure 10:	Technique for Vazirani–Akinosi nerve block. ⁴³	16
Figure 11:	(11A) Diagrammatic representation of a transverse section of the right mandibular ramus at the level at which an IANB would be given. (11B) Photograph of a skull with simulated maximum opening of the mouth.....	18

List of Figures cont...

Fig. No.	Title	Page No.
Figure 12:	(12A) intraoral photographs of right side of the oral cavity showing the key intraoral landmarks observed and palpated when administering an CIANB. (12B) show site of syringe's insertion is 6-10 mm above occlusal plane of mandibular teeth. (12C) shows the barrel of syringe in the contra lateral side of side of needle's insertion between premolar and molar area, level of insertion 6 to 10 mm above occlusal plane.....	19
Figure 13:	(13A) illustrated the palpation of the anterior border the ramus and retraction of the of tissue mildly by operator's finger. (13B) illustrated the red dot is the imaginary midpoint, blue dot is the first site of insertion (6 to 8 mm above the midpoint and 8 to 10 mm posterior to the anterior border of ramus). (13C) The first site of needle's insertion and the syringe's barrel is placed at contra lateral side between canine and premolar. (13D) The barrel of syringe is adjusted towards midline of mandible to insert the needle freely to the further along medial side of ramus, the needle is inserted 20 to 24 mm to be superior to IAN entry into its MF. (13E) The barrel of syringe is taken back to the contra lateral side in order to bring the needle closer to the bone.....	20
Figure 14:	(A) illustrates short distance from internal oblique ridge to lingula. (B) illustrates the needle is in place in the supralingual notch, touching the bone and above occlusal plane in range from 6 to 10 mm, IOR= internal oblique ridge.....	21

List of Figures cont...

Fig. No.	Title	Page No.
Figure 15:	(A) illustrates the novel approach to IANB (B) illustrates conventional inferior alveolar nerve block (CIANB). ⁵⁰	22
Figure 16:	Illustrates retromolar triangle intraorally. ⁵²	23
Figure 17:	Schematic diagrams show the relationship of the pterygomandibular space and the landmarks for the insertion and the angles of the syringe in the anterior technique and conventional technique	24
Figure 18:	The modification of the lingual cut according to Hunsuck and Epker. Notice the short horizontal cut on the lingual side. ⁶¹	26
Figure 19:	Introduction of the caudal (fourth) osteotomy. As the osteotomy reaches the mandibular angle, it allows the surgeon to split the mandible as if opening a book. ⁶¹	26
Figure 20:	(A) The incision is made well lateral to the attached gingiva in the molar region. (B) The mandibular ramus is exposed in the subperiosteal plane. (C) The lateral ramus, angle, and body are exposed. (D) The inferior border is exposed. ⁵⁷	27
Figure 21:	(A, B) A channel retractor is placed to support the mandible. (C, D) The anterior ramal border and coronoid process are exposed to the insertion of the temporalis muscle. ⁵⁷	27
Figure 22:	(A) A medial ramus retractor is placed to protect the nerve. (B) A horizontal osteotomy (dashed line) is made with (C) a Lindermann burr. (D) Alternately, a reciprocating saw can be used. ⁵⁷	28

List of Figures cont...

Fig. No.	Title	Page No.
Figure 23:	A reciprocating saw is used along the external oblique ridge and continued to the region of the first and second molar where a vertical osteotomy is made	28
Figure 24:	(A) With a series of osteotomes the corticotomy is converted to an osteotomy. (B) A ramal spreader is introduced, and the separation of the medial and lateral cortices is completed with the nerve within the medial component. (C, D). The lateral ramal component should be completely freed and independent of the distal dental arch. ⁵⁷	29
Figure 25:	The mandible is either advanced or set back by placing it within the prefabricated surgical splint and maxillary-mandibular dental fixation. ⁵⁷	29
Figure 26:	The repositioned elements are fixed with positional screws that can be placed via transbuccal.....	30
Figure 27:	Development of a long osteotomy to preserve proximal segment width	31
Figure 28:	(A) The different FOV option sizes from the Vatech CBCT (Vatech America, FortLee, NJ). (B) Different field of views available in CBCT units. ⁴⁴	33
Figure 29:	CBCT image acquisition, the X-ray tube and detector move concomitantly around the rotation axis. ⁶⁵	34
Figure 30:	Corrected axial view on I-CAT Vision.....	41
Figure 31:	Corrected coronal view on I-CAT Vision.....	42
Figure 32:	Corrected Sagittal view obtained.	42

List of Figures cont...

Fig. No.	Title	Page No.
Figure 33:	Corrected sagittal view on I-CAT vision software showed AB-MF in forty-three dentate female.....	43
Figure 34:	Corrected sagittal view on I-CAT vision software showed PB-MF in forty three dentate female.....	43
Figure 35:	Corrected sagittal view on I-CAT vision software showed the CW in forty three dentate female.....	43
Figure 36:	Corrected sagittal view on I-CAT vision software showed MF-MN in a forty-three dentate female.....	44
Figure 37:	Corrected sagittal view on I-CAT vision software showed MF-IB in a forty-three dentate female.....	44
Figure 38:	Corrected sagittal view on I-CAT vision software showed RW in a forty-three years old dentate female.....	44
Figure 39:	Corrected axial view on I-CAT vision software to measure MFD.	45
Figure 40:	Corrected axial view on I-CAT vision software showed MFD in a forty-three years old dentate female.....	45
Figure 41:	Corrected axial view on I-CAT vision software to measure (MF-OP and MF-ARMT)	46
Figure 42:	Corrected coronal view on I-CAT vision software to measure (MF-OP and MF-ARMT).	47

List of Figures cont...

Fig. No.	Title	Page No.
Figure 43:	Corrected sagittal view on I-CAT vision software showed MF-OP in a forty-three dentate female.....	47
Figure 44:	Corrected sagittal view on I-CAT vision software showed MF-ARMT in a forty-three dentate female.....	47
Figure 45:	Corrected sagittal view showed AB-MF = 25.25 mm.....	48
Figure 46:	Corrected sagittal view showed PB-MF = 10.50 mm.....	48
Figure 47:	Corrected sagittal view showed CW = 5 mm.	49
Figure 48:	Corrected sagittal view showed RW = 33.75 mm.....	49
Figure 49:	Corrected sagittal view showed MN-MF = 17.50 mm.....	50
Figure 50:	Corrected sagittal view showed MF-IB = 22.25 mm.....	50
Figure 51:	Axial view showed MFD = 4 mm.....	51
Figure 52:	Corrected sagittal view showed MF-OP = 7 mm.....	51
Figure 53:	Corrected sagittal view showed MF-ARMT = 7.25 mm.....	52
Figure 54:	Gender distribution of the studied CBCT scans	54
Figure 55:	Age distribution of the studied CBCT scans.....	55
Figure 56:	Dental status of the of the studied CBCT scans	55