



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

∞∞∞∞

تم رفع هذه الرسالة بواسطة /صفاء محمود عبد الشافي

بقسم التوثيق الإلكتروني بمركز الشبكات وتكنولوجيا المعلومات دون

أدنى مسئولية عن محتوى هذه الرسالة.

ملاحظات: لا يوجد





Regenerative potential of immature dog teeth treated with bioceramic scaffold and bone morphogenic protein (An Animal study)

Thesis
Submitted to the Faculty of Dentistry,
Ain Shams University

For

Partial Fulfillment of Requirements of the Doctorate Degree
in Endodontics

By

Samar Talaat Mohamed

B.D.S

(Future University-2011)

M.Sc.

(Ain Shams University-2017)

2022

بسم الله الرحمن الرحيم

(قَالُوا سُبْحَانَكَ لَا
عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا
إِنَّكَ أَنْتَ الْعَلِيمُ
الْحَكِيمُ ﴿٣٢﴾)

صدق الله العظيم

Supervisors

Prof. Ahmed Abd El Rahman Hashem

**Professor of Endodontics
Head of Endodontic Department
Faculty of Dentistry, Ain Shams University**

Prof. Ashraf Mohamed Abu-Seida

**Professor of Surgery, Anesthesiology and Radiology
Faculty of Veterinary Medicine, Cairo University**

Dr. Adel AbdelWahed Mahmoud

**Associate professor of Endodontics
Faculty of Oral and Dental Medicine, Future
University in Egypt**

Dr. Tarek Mustafa Abdel Aziz

**Lecturer in Endodontics
Faculty of Dentistry, Ain Shams University**

Dedication

To my dearest and greatest mother

To my caring and lovable father

To the soul of my lovely Uncle

To my supportive husband

To my lovely sister and brother

To my sweet angel Tamara

To my best friends Sara and Reham

Acknowledgement

I would like to express my deep gratitude to *Professor. Ahmed Abd El Rahman Hashem*, Professor of Endodontics, Faculty of Dentistry, Ain Shams University for his kind guidance, sincerity, extraordinary supervision, and unlimited support and help throughout my academic and clinical work.

My sincere thanks to *Professor. Ashraf Mohamed Abu-Seida*, Professor of Surgery, Anesthesiology and Radiology, Faculty of Veterinary Medicine, Cairo University, for his precious advice, wise opinions, and cooperation.

My deep appreciation to *Professor. Adel Abdel Wahed Mahmoud*, Associate professor of Endodontics, Faculty of Oral and Dental Medicine, Future University in Egypt for his excellent advice, valuable stimulating guidance and help during this study.

I would like to acknowledge and give my warmest thanks to my supervisor *Dr. Tarek Moustafa Abdel Aziz*, Lecturer in Endodontics, Faculty of Dentistry, Ain Shams University who made this work possible. His guidance and advice carried me through all the stages of writing my thesis.

I am deeply grateful to *Professor. Mohamed Mokhtar*, Professor of Endodontics, Faculty of Dentistry, Ain Shams University for his cooperation, continuous effort that paved the way for completing this study by giving me of his precious time and for his valuable comments and suggestions.

Last but not least, I would like to thank **all members of Endodontic department**, Faculty of Dentistry, Future University in Egypt especially *Hatem Tarek* and *Sherief Elsaqa* for their valuable help and cooperation.

List of Contents:

Subject	Page
List of figures	ii
List of tables	x
List of abbreviations	xiii
I. Introduction.	1
II. Review of literature.	4
• Management of open apex	4
I. Apexification	4
A. Calcium hydroxide apexification	5
B. Apical plug	5
II. Regenerative endodontic therapy	6
A. Revascularization	7
B. Tissue Engineering	12
III. Aim of the work.	37
IV. Materials and methods.	38
1. Animal models	38
2. Classification of samples	40
3. Experimental procedures	43
4. Methods of Evaluation	55
V. Results	65
VI. Discussion	111
vii. Summary and conclusions	132
viii. References	140
Arabic summary	

List of figures:

Figure	Figure Title	Page
1	Schematic diagram for samples Classification	42
2	Showing administration of general anesthesia	44
3	Showing a sterile file size #35 was used to disrupt the remnant pulp tissue in the canal	46
4	A: Showing injection of calcium hydroxide B: Radiograph after calcium hydroxide placement	47
5	Showing a tooth after placement of MTA plug	49
6	Showing injection of Mesoporous silica Nanoparticle scaffold	50
7	Transmission electron microscopy images of MSNs A(200nm), B (100nm)	52
8	Showing sealing of access cavity using glass ionomer restoration	55
9	Bar chart representing mean and standard deviation values for percentage increase in root length in different groups	68
10	Bar chart representing mean and standard deviation values for percentage increase in root thickness in different groups	71
11	Bar chart representing mean and standard deviation values for percentage increase in	74

	apical closure in different groups	
12	Representative radiographs showing no changes in root length, root thickness and apical diameter (positive control group) preoperatively (A), after one month (B) and three months (C) comparing the postoperative radiograph (right side) to preoperative radiograph (left side).	75
13	A representative sample of the negative control group showing marked changes in root length, root thickness and apical diameter from pre-operative (A), after one month (B) and three months (C) comparing the postoperative radiograph (right side) to its preoperative radiograph (left side).	75
14	A representative sample of the blood clot group showing changes in root length, root thickness and apical diameter from pre-operative (A), after one month (B) and three months (C) comparing the postoperative radiograph (right side) to its preoperative radiograph (left side).	76
15	A representative sample of the mesoporous silica scaffold group showing changes in root length, root thickness and apical diameter from pre-operative (A), after one month (B) and three months (C) comparing the	76

	postoperative radiograph (right side) to its preoperative radiograph (left side).	
16	A representative sample of the mesoporous silica+BMP2 group showing changes in root length, root thickness and apical diameter from pre-operative (A), after one month (B) and three months (C) comparing the postoperative radiograph (right side) to its preoperative radiograph (left side).	77
17	Box plot representing median and range values for inflammatory cell counts in different groups (Circles represent outliers)	82
18	Photomicrograph showing moderate inflammatory (score 2) cell infiltration in subgroup A (blood clot) after one month (400x)	82
19	Photomicrograph showing moderate inflammatory cell (score 2) infiltration in subgroup B (meso-silica) after one month (400x)	83
20	Photomicrograph showing mild inflammatory cell infiltration (score 1) group C (Meso-silica+BMP2) after one months (400x)	83
21	Photomicrograph showing severe inflammatory cell infiltration (score 3) and edematous space in group D (positive control) after one months (400x)	84

22	Photomicrograph showing mild inflammatory cell infiltration (score 1) in group II subgroup A (blood clot) (400x)	84
23	Photomicrograph showing no inflammatory cells infiltration (score 1) in group II subgroup C (meso-silica+BMP2) (400x)	85
24	Photomicrograph mild inflammatory cells infiltration (score 1) in group II subgroup B (meso-silica) (400x)	85
25	Bar chart representing prevalence of bone resorption in different groups	88
26	Photomicrograph showing active apical root resorption as odontoclasts and Howship's lacunae could be detected in group I subgroup A (400x)	89
27	Photomicrograph showing active bone resorption as osteoclasts and Howship's lacunae could be detected in group I subgroup B (400x)	89
28	Photomicrograph showing active bone resorption in some areas in group II subgroup B (100x)	90
29	Photomicrograph showing showed no signs of apical root resorption or bone resorption in group II subgroup C (100x)	90
30	Box plot representing median and range values for vial tissue inside the pulp scores in different groups (Circles and stars represent outliers)	94

31	Photomicrograph showing connective tissue (C.T) in growth inside the pulp cavity reaching the apical third of the root canal in group I subgroup A (200x)	95
32	Photomicrograph showing C.T inside the pulp cavity reaching the middle third of the root canal in group I subgroup B (200x)	95
33	Photomicrograph showing apical two third of the root in group II after three months (100x). Vital tissue infiltration along the pulp space is seen (green arrow). Fibrous tissue formation inside the pulp space with signs of complete hard tissue formation and apical closure (yellow arrow) are observed. Mild inflammatory cell infiltration (blue arrow).	96
34	Photomicrograph At a higher magnification showing tissue resembles the pulp tissue with variable amounts of inflammatory cells infiltration (black arrow), noticeable angiogenic activity (green arrows) and a layer of odontoblast-like cells undergo differentiation opposite to a predentin layer (blue arrow) in group I subgroup C (400x)	96

35	Photomicrograph showing C.T in growth inside the pulp cavity. The tissue resembles periodontal connective tissue (black arrow) with variable amounts of inflammatory cells infiltration (green arrows) and noticeable angiogenic activity (red arrow) in group II subgroup A (400x)	97
36	Photomicrograph showing C.T inside the pulp cavity reaching the coronal third of the root canal (blue arrow). In some specimens areas of a mineralized tissue resembling osteodentin with a layer of odontoblast-like cells (black arrow) could be detected in group II subgroup B (200x)	97
37	Photomicrograph showing tissue resembles the pulp tissue no inflammatory cells infiltration. A definite layer of odontoblast-like cells could be observed opposite to a predentin layer (black arrow) in group II subgroup C (400x)	98
38	Box plot representing median and range values for new hard tissue scores in different groups (Stars represent outliers)	102
39	Photomicrograph showing apical hard tissue formation that resembles cementum-like tissue covered by thin layer of cementoid (blue arrow). Cementocyte-like cells (black arrow) and empty lacunae with degenerated	103

	cementocyte-like cells (green arrow) could be detected in group II subgroup A (400x)	
40	Photomicrograph showing large areas of mineralized tissue that resembles osteodentin in group I subgroup B (200x)	103
41	Photomicrograph showing large areas of mineralized tissue that resembles osteodentin covered with a layer of predentin (red arrow). Odontoblast-like cells could be observed entrapped inside the mineralized tissue (black arrow). Areas of tubular dentin (green arrow), and odontoblast-like cells opposite to the predentin layer (blue arrow) could be detected in group II subgroup B (200x)	104
42	Photomicrograph showing several layers of odontoblast-like cells could be observed opposite to predentin in group II subgroup C (400x)	104
43	Photomicrograph showing formation of single layer of odontoblastic like cells in inner surface of dentin in group I subgroup C (200x)	105
44	Bar chart representing prevalence of apical closure in different groups	107
45	Photomicrograph showing apical closure in group I subgroup A (100x)	108