



بسم الله الرحمن الرحيم

∞∞∞∞

تم رفع هذه الرسالة بواسطة / سامية زكى يوسف

بقسم التوثيق الإلكتروني بمركز الشبكات وتكنولوجيا المعلومات دون أدنى

مسئولية عن محتوى هذه الرسالة.

ملاحظات: لا يوجد





The Effect of Intrauterine Lidocaine versus Warm Saline Distention Medium for Pain Control during Office Hysteroscope

Thesis

*Submitted for Partial Fulfillment of Master Degree
in Obstetric & Gynecology*

Presented by

Dina Salah Mohamed Hassan Elwan

M.B.B.Ch, Faculty of Medicine, Mansoura University, 2011

Supervised by

Prof. Dr. Ahmed Ramy Mohamed Ramy

Professor of Obstetric & Gynecology

Faculty of Medicine, Ain Shams University

Dr. Amr Ahmed Mahmoud Riad

Assistant Professor of Obstetric & Gynecology

Faculty of Medicine, Ain Shams University

Dr. Ahmed Gamal Abdelnasser

Lecturer of Obstetric & Gynecology

Faculty of Medicine, Ain Shams University

**Faculty of Medicine
Ain Shams University**

2022

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

لسبحانك لا علم لنا
إلا ما علمتنا إنك أنت
العليم العظيم

صدق الله العظيم

سورة البقرة الآية: ٣٢



Acknowledgement

First and foremost thanks to *ALLAH*, the Most Merciful.

*I wish to express my deepest thanks, gratitude and appreciation to **Prof. Dr. Ahmed Ramy Mohamed Ramy**, Professor of Obstetric & Gynecology, Faculty of Medicine, Ain Shams University, for his meticulous supervision, kind guidance, valuable instructions and generous help.*

*Special thanks are due to **Dr. Amr Ahmed Mahmoud Riad**, Assistant Professor of Obstetric & Gynecology, Faculty of Medicine, Ain Shams University, for his sincere efforts, fruitful encouragement.*

*I am deeply thankful to **Dr. Ahmed Gamal Abdelnaser**, Lecturer of Obstetric & Gynecology, Faculty of Medicine, Ain Shams University, for his great help, outstanding support, active participation and guidance.*

I would like to express my hearty thanks to all my family for their support till this work was completed.

Dina Salah Mohamed Hassan Elwan

Contents

Title	Page
List of Abbreviations.....	I
List of Table.....	II
List of Figures.....	III
Introduction	1
Aim of the work.....	4
Review of Literature	
Hysteroscopy.....	5
Distension Media in Hysteroscopy	20
Pain.....	35
Nerve Supply to the Pelvic Floor	39
Patients and methods	56
Results	65
Discussion.....	85
Conclusion.....	94
Recommendations	95
Summary.....	96
References	98
الملخص العربي.....	122
•	

List of Abbreviations

AAGL	American Association of Gynecological Laparoscopists
ARDS	Adult respiratory distress syndrome
AUB	Abnormal uterine bleeding
BMI	Body mass index
CDIS	Controlled Distention Irrigation System
CPP	Chronic pelvic pain
DRG	Dorsal root ganglia
HS	Highly significant
HS	Hysteroscopy
IASP	International Association for the Study of Pain
IQR	Inter-quartile range
IUP	Intrauterine pressure
IV	Intravenous
KO	Kelvin
NRS	Numerical Rating Scale
NS	Non significant
OD	Outer diameter
S	Significant
SPSS	Statistical Package for Social Science
VAS	Visual analogue scale

List of Table

Table No	Subjects	Page
Table (1):	Comparison between control group, warm saline group and lidocaine infusion group regarding demographic data and characteristics of the studied patients.....	65
Table (2):	Comparison between control group, warm saline group and lidocaine infusion group regarding cycle phase, history of previous uterine procedure, indications and duration by minutes	68
Table (3):	Comparison between control group, warm saline group and lidocaine infusion group regarding endometrial, cavity and tubes findings.....	72
Table (4):	Comparison between control group, warm saline group and lidocaine infusion group regarding visual analogue scale (VAS) pain score	75
Table (5):	Comparison between control group and warm saline group regarding visual analogue scale (VAS) pain score	77
Table (6):	Comparison between control group and lidocaine infusion group regarding visual analogue scale (VAS) pain score	79
Table (7):	Comparison between warm saline group and lidocaine infusion group regarding visual analogue scale (VAS) pain score	81
Table (8):	Relation between number of parity and VAS score at different times of measurement.....	83

List of Figures

Figure No	Subjects	Page
Figure (1):	Karl Storz flexible hysteroscope	7
Figure (2):	Typical instruments feature two sheaths, one for irrigation and another for suction (A) 4-mm and 5-mm sheaths. (B) 2.9-mm and 2.0-mm rod lens telescope. (C) The oval tip allows atraumatic introduction.	9
Figure (3):	Sympathetic, parasympathetic and pudendal nerve supply to the anorectum	40
Figure (4):	Numeric rating scale.....	53
Figure (5):	FACE pain rating scale	54
Figure (6):	Behavioral pain scale	55
Figure (7):	VAS is a straight horizontal line of fixed length (100 mm).	62
Figure (8):	Comparison between control group, warm saline group and lidocaine infusion group regarding age of the studied patients.....	66
Figure (9):	Comparison between control group, warm saline group and lidocaine infusion group regarding BMI of the studied patients.....	66
Figure (10):	Comparison between control group, warm saline group and lidocaine infusion group regarding parity of the studied patients	67
Figure (11):	Comparison between control group, warm saline group and lidocaine infusion group regarding cycle phase	69
Figure (12):	Comparison between control group, warm saline group and lidocaine infusion group regarding history of previous uterine procedure.....	70
Figure (13):	Comparison between control group, warm saline group and lidocaine infusion group regarding indications of the studied patients.....	71

Figure No	Subjects	Page
Figure (14):	Comparison between control group, warm saline group and lidocaine infusion group regarding duration by minutes.....	71
Figure (15):	Comparison between control group, warm saline group and lidocaine infusion group regarding endometrial findings	73
Figure (16):	Comparison between control group, warm saline group and lidocaine infusion group regarding cavity findings.....	73
Figure (17):	Comparison between control group, warm saline group and lidocaine infusion group regarding tubes findings	74
Figure (18):	Comparison between control group, warm saline group and lidocaine infusion group regarding visual analogue scale (VAS) pain score	76
Figure (19):	Comparison between control group and warm saline group regarding visual analogue scale (VAS) pain score	78
Figure (20):	Comparison between control group and lidocaine infusion group regarding visual analogue scale (VAS) pain score	80
Figure (21):	Comparison between warmsaline group and lidocaine infusion group regarding visual analogue scale (VAS) pain score	82
Figure (22):	Relation between number of parity and VAS score at different times of measurement.....	84