

## بسم الله الرحمن الرحيم

 $\infty\infty\infty$ 

تم رفع هذه الرسالة بواسطة / سامية زكى يوسف

بقسم التوثيق الإلكتروني بمركز الشبكات وتكنولوجيا المعلومات دون أدنى مسئولية عن محتوى هذه الرسالة.

ملاحظات: لا يوجد

AIN SHAMS UNIVERSITY

Since 1992

Propries 1992



## Incidence and Prognostic Value of Single Lead QRS Fragmentation in Successfully Revascularized Inferior STEMI Patients

## Thesis

Submitted in Partial Fulfillment of the M.Sc. Degree in **Cardiology** 

By

### **Sherif Abdelhamid Elbialy Hashish**

MB.BCh, Mansoura University Cardiology Resident at National Heart Institute

Supervised by

## Prof. Dr. Ahmed Abdelrahman Sharafeldin

Professor of Cardiology
Faculty of Medicine – Ain Shams University

#### Dr. Sameh Attia Amin

Lecturer of Cardiology
Faculty of Medicine – Ain Shams University

Faculty of Medicine Ain Shams University 2022



سورة البقرة الآية: ٣٢

## Acknowledgment

First and foremost, I feel always indebted to ALLAH, the
Most Kind and Most Merciful.

I'd like to express my respectful thanks and profound gratitude to **Prof. Dr. Ahmed**Abdelrahman Sharafeldin, Professor of Cardiology, Faculty of Medicine – Ain Shams University for his keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.

I am also delighted to express my deepest gratitude and thanks to **Dr. Sameh Attia Amin,**Lecturer of Cardiology, Faculty of Medicine – Ain Shams University, for his kind care, continuous supervision, valuable instructions, constant help and great assistance throughout this work.

I would like to express my hearty thanks to all my family for their support till this work was completed.

Last but not least my sincere thanks and appreciation to all patients participated in this study.

Sherif Abdelhamid Elbialy Hashish

# Tist of Contents

Title	Page No.
Tist of Tables	i
List of Figures	ii
List of Abbreviations	v
Introduction	1
Aim of the Work	3
Review of Literature	
Myocardial Infarction	4
Fragmented-QRS	22
Patients and Methods	37
Results	48
Discussion	75
Limitations	87
Summary	88
Conclusion	93
Recommendations	94
References	95
Arabic Summary	

# Tist of Tables

Table No.	Title	Page No.
<b>Table (1):</b>	Sociodemographic characteristics and ridistribution among the study population (new distribution)	
<b>Table (2):</b>	Incidence of QRS fragmentation shown included patients (n=150)	
<b>Table (3):</b>	The association between socioder characteristics and associated risk distribution and FQRS among included (n=150)	factor d patients
<b>Table (4):</b>	Post PCI ECG data among included patient	ts (n=150) 56
<b>Table (5):</b>	The difference between study groups c ECG findings (n=150)	
<b>Table (6):</b>	The laboratory data of included patients (	n=150)60
<b>Table (7):</b>	The difference between study groups c lab measures (n=150)	
<b>Table (8):</b>	The operative details of included patients	(n=150)63
<b>Table (9):</b>	The difference between study groups c operative details (n=150)	
<b>Table (10):</b>	ECHO findings among included patients	(n=150)67
<b>Table (11):</b>	The difference between study groups c Echo findings (n=150).	
<b>Table (12):</b>	Major adverse cardiac events (MACE) or patients (n=150).	
<b>Table (13):</b>	The association between MACE and FQI included patients (n=150)	_

# Tist of Figures

Fig. No.	Title	Page No.
Figure (1):	12-lead electrocardiogram (ECG) de evidence of inferior myocardial infarcti	-
Figure (2):	Fragmented QRS in inferior leads	21
Figure (3):	Classification of fragmented QRS (v patterns)	
Figure (4):	Effects of low-pass filter	26
Figure (5):	Different morphologies of the fra wave and a Q-R-borderline-fQRS mother right	orphology on
Figure (6):	Different RsR-morphologies	29
<b>Figure (7):</b>	Different morphologies of the notch left and an example of the R-S-bord on the right	lerline-fQRS
Figure (8):	A, ECG of an 11-year-old boy with sm mild right atrial enlargement showing pattern	g RSR0 -V1
Figure (9):	Leads V1 to V3 of a very lean 15-y without heart disease	ear-old man
<b>Figure</b> (10):	Classification of fragmented QRS (v patterns)	
<b>Figure (11):</b>	65 years old male patient, smoke hypertensive with BMI=28,	
<b>Figure (12):</b>	75 years old male patient, s hypertensive not diabetic with BMI=2	
<b>Figure (13):</b>	Difference between studied groups condistribution.	0 0
<b>Figure (14):</b>	Difference between studied groups gender distribution	

# Tist of Figures (Cont...)

Fig. No.	Title	Page No.
<b>Figure (15):</b>	Difference between studied groups of BMI.	-
<b>Figure (16):</b>	Difference between studied groups smoking status.	_
<b>Figure (17):</b>	Difference between studied groups diabetes mellitus.	_
<b>Figure (18):</b>	Difference between studied groups of Hypertension.	_
<b>Figure (19):</b>	Difference between studied groups of Dyslipidemia.	_
<b>Figure (20):</b>	Difference between studied groups of family history.	_
<b>Figure (21):</b>	Difference between studied groups of heart rate.	_
<b>Figure (22):</b>	Difference between studied groups of QRS width.	_
<b>Figure (23):</b>	Difference between studied groups of STR abnormalities	
<b>Figure (24):</b>	Difference between studied groups of Pathological Q wave	_
<b>Figure (25):</b>	Difference between study groups of preoperative CKMB.	
<b>Figure (26):</b>	Difference between study groups of postoperative CKMB	_
<b>Figure (27):</b>	Difference between studied groups of Culprit artery.	
<b>Figure (28):</b>	Difference between studied groups on number of used stents.	

# Tist of Figures (Cont...)

Fig. No.	Title	Page No.
<b>Figure (29):</b>	Difference between studied groups number of affected vessels.	_
<b>Figure (30):</b>	Difference between studied groups LVEDD	•
<b>Figure (31):</b>	Difference between studied groups LVESD	•
<b>Figure (32):</b>	Difference between studied groups ejection fraction	_
<b>Figure (33):</b>	Difference between studied groups wall motion score index	•
<b>Figure (34):</b>	Major adverse cardiac events amon patients	•
<b>Figure (35):</b>	The difference between study groups Major adverse cardiac events amon patients	g included