

بسم الله الرحمن الرحيم



HOSSAM MAGHRABY



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



HOSSAM MAGHRABY

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
على هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن


تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HOSSAM MAGHRABY



بعض الوثائق الأصلية تالفة



HOSSAM MAGHRABY



بالرسالة صفحات

لم ترد بالأصل



HOSSAM MAGHRABY

**Susceptibility of different pests to
infection of entomogenous nematodes**





A THESIS
Submitted for the
award of the degree of
Doctor of Philosophy
(Entomology)

BY

Rawhia Hassan Ramadan
M. Sc. Entomology, Zagazig University
(1989)

Department of Entomology
Faculty of Science
Zagazig University
(BANHA)

1994



قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا
عَلَّمْتَنَا إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ
سُورَةُ الْبَقَرَةِ - آيَةُ ٢٢٠

APPROVAL SHEET

Title of thesis: "Susceptibility of different
pests to infection of
entomogenous nematodes"

Degree: Ph. D. (Entomology)

Student name: Rawhia Hassan Ramadan

Approved by:

Name	Title	Signature
_____	_____	_____
_____	_____	_____
_____	_____	_____

Date: / / 1994

BIOGRAPHY

Date of B. Sc. graduation: May 1984.

Degrees awarded: B. Sc., Entomology (Cairo
University) in 1984.
M. Sc., Entomology (Zagazig
University) in 1988.

Occupation: Assistant Lecturer,
Department of Entomology,
Faculty of Science, Zagazig
University, Banha branch.

ACKNOWLEDGMENT

The author owes her great indebtedness and deep gratitude to Prof. Dr. Nahed Helmy, Professor of Entomology, head of Dept., Faculty of science, Zagazig University, Benha branch for her kind supervision, valuable advice, constructive criticism and final preparation of the manuscript.

I would like also to express my heartiest gratitude and my sincere thanks to the spirit of the late Dr. Tyseer Ahmed H. El-Kifl, an Associate Professor at Department of Pests and Plant Protection, National Research Center for her serious valuable supervision, constructive criticism, literature material and general help throughout this work.

Deep gratitude is also due to Dr. Muhammad M. Shamseldean, Lecturer of Nematology, Department of Agric. Zoology and Nematology, Faculty of Agriculture, Cairo University for his kind supervision, encouragement during the work and final preparation of the manuscript.

I would like to thank with deep gratitude, Dr. Mahmoud M. E. Saleh, researcher of biological control, Department of Pests and Plant Protection, National Research Center for his help during the work and revising the manuscript.

Sincere thanks are also extended to Prof. Dr. Adel Merdan, Professor of Entomology, Faculty of science, Ain-Shams University for his valuable supervision of this work.

To my dear parents, husband and sisters, Amal, Mervat and Reda, I would like to express my gratitude for their kind help during this work.

Last but not least, I want to express my deep thanks to my colleagues of the Entomology Department, Faculty of Science, Zagazig University, Banha branch.

CONTENTS

	page
1. INTRODUCTION.....	1
2. REVIEW OF LITERATURE.....	5
I. ENTOMOPATHOGENIC NEMATODES OF FAMILY STEINERNEMATIDAE AND HETERORHABDITIDAE AS A BIOLOGICAL CONTROL AGENTS OF SCARABAEID PESTS.....	6
II. MODE OF ENTRY AND ACTION OF ENTOMOPATHOGENIC NEMATODES.....	1
III. DISPERSAL AND INFECTIVITY OF ENTOMOPATHOGENIC NEMATODES IN VARIOUS SOILS.....	15
IV. SURVIVAL OF INFECTIVE JUVENILES OF ENTOMOPATHOGENIC NEMATODES IN DIFFERENT SOIL TYPES.....	22
3. MATERIALS AND METHODS.....	
I. REARING OF ADULT <u>PACHNODA</u> <u>FASCIATA</u> F. (COLEOPTERA: SCARABAEIDAE).....	25
II. REARING OF IMMATURE STAGES OF <u>P. FASCIATA</u> F.	
III. REARING OF HAIRY BEETLE <u>TROPINOTA SQUALIDA</u> SCOP. (COLEOPTERA: SCARABAEIDAE)...	26
IV. COLLECTION OF SUGAR-CANE GRUB <u>PENTODEN BISPINOSUS</u> (COLEOPTERA: SCARABAEIDAE).....	
V. SOURCES OF TESTED NEMATODE SPECIES AND STRAINS.....	27
VI. PROPAGATION OF ENTOMOPATHOGENIC NEMATODES.....	29
VII. SUSCEPTIBILITY OF ADULT BEETLES TO ENTOMOPATHOGENIC NEMATODES.....	33

VIII. SUSCEPTIBILITY TESTS OF <u>P. FASCIATA</u> AND <u>T. SQUALIDA</u> LARVAE TO ENTOMOPATHOGENIC NEMATODE INFECTIONS.....	34
IX. SURVIVAL OF INFECTIVE NEMATODE JUVENILES IN DIFFERENT SOILS.....	35
X. DISPERSAL POTENTIALITY OF THE INFECTIVE JUVENILES OF ENTOMOPATHOGENIC NEMATODES.....	
4. RESULTS AND DISCUSSION.....	36
I. SUSCEPTIBILITY OF SCARABAEID PESTS TO INDIGENOUS AND FOREIGN ENTOMOPATHOGENIC NEMATODES.....	38
II. COMPARATIVE DISPERSAL OF A FOREIGN AND AN EGYPTIAN ENTOMOPATHOGENIC NEMATODE.....	71.
II. SURVIVAL POTENTIAL OF TWO ENTOMOPATHOGENIC NEMATODES....	75
5. SUMMARY.....	79
6. LITERATURE CITED.....	83
THE ARABIC SUMMARY.....	

LIST OF TABLES

9

Table 1. Percent mortality of 1st instar larvae of Pachnoda fasciata infected with different concentrations of indigenous Egyptian isolates of Heterorhabditis spp., EBNU3, EGB1 and EIS7.....41

Table 2. Percent mortality of 1st instar larvae of Pachnoda fasciata infected with different concentrations of Heterorhabditis bacteriophora (H. b.), Steinernema carpocapsae (S. c.) and Steinernema glaseri (S. g.)..... 44

Table 3. Percent mortality of 3rd instar larvae of Pachnoda fasciata infected with different doses of Heterorhabditis spp., ESS1 and Steinernema glaseri (S.g.)..... 46

Table 4. Rate of migration in infective juveniles (IJ's) of Steinernema glaseri emerging from 3rd instar larvae of Pachnoda fasciata within 6 days post-infection..... 48

Table 5. Percent mortality of 3rd instar larvae of Pentoden bispinosus infected with different doses of Heterorhabditis spp., ESS1 and Steinernema glaseri (S.g.)..... 53

Table 6. Percent mortality of 1st instar larvae of Tropinota squalida infected with different concentrations of Heterorhabditis bacteriophora (H. b.), Steinernema glaseri (S. g.) and the Egyptian isolates Heterorhabditis spp. EBNU3 and EGB1..... 55

Table 7. Percent mortality of 3rd instar larvae of Tropinota squalida infected with different concentrations of Heterorhabditis bacteriophora (H. b.), Steinernema glaseri (S. g.) and Steinernema carpocapsae (S. c.).....58

Table 8. Percent mortality of adults Pachnoda fasciata infected with different concentrations of Heterorhabditis bacteriophora (H. b.), Steinernema carpocapsae (S. c.), Steinernema glaseri (S. g.) and the Egyptian heterorhabditid isolates of EBNU3 and EGB1.....90

Table 9. Percent mortality of adults Tropinota squalida infected with different concentrations of Heterorhabditis bacteriophora (H. b.), Steinernema glaseri (S. g.), Steinernema carpocapsae (S. c.) and an Egyptian isolate of Heterorhabditis sp., EBNU3 92

Table 10. Medium lethal dose (LD₅₀) of entomopathogenic nematodes Steinernema glaseri (S.g.), Steinernema carpocapsae (S.c.), Heterorhabditis bacteriophora (H. b.) and the Egyptian isolates of Heterorhabditis spp. (EBNU3 and EGB1) tested against Pachnoda fasciata (P.f.), Tropinota squalida (T.S.) and Pentoden bispinosus (P.b.) larvae (Coleoptera: Scarabaeidae).....94