

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

# بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

# جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY

### Possible Protective Effects of Stem Cells Treatment on Ischemic/Reperfused Hearts in Hypertensive Rats Model

### Thesis

Submitted for partial fulfillment of master degree in **Basic Medical Science (Medical Physiology)** 

By

#### Nahla Mohamed Hussien

*M.B.*, *B.Ch*.

Under supervision of

#### Prof. Dr. Gehane Mahmoud Hamed

Professor of medical Physiology Faculty of Medicine - Ain Shams University

### **Prof. Dr. Hanaa Ahmed Amer**

Professor of Clinical Pathology Faculty of Medicine - Ain Shams University

#### Assis. Prof. Dr. Manal Said Abd el Hamid

Assistant Professor of medical Physiology Faculty of Medicine - Ain Shams University

### Dr. Einas Mohamed Nagib

Lecturer of medical Physiology Faculty of Medicine - Ain Shams University

> Faculty of Medicine Ain Shams University



سورة البقرة الآية: ٣٢

## Acknowledgments

# First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.

I wish to express my deepest thanks, gratitude and appreciation to **Prof. Dr. Gehane Mahmoud Hamed**, Professor of medical Physiology, Faculty of Medicine-Ain Shams University, for her meticulous supervision, kind guidance, valuable instructions and generous help.

Special thanks are due to **Prof. Dr. Hanaa Ahmed Amer**, Professor of Clinical Pathology, Faculty of Medicine-Ain Shams University, for her sincere efforts, fruitful encouragement.

I am deeply thankful to Assis. Prof. Dr. Manal Said Abd el Hamid, Assistant Professor of medical Physiology, Faculty of Medicine-Ain Shams University, for her great help, outstanding support, active participation and guidance.

I would like to express my deepest gratitude and appreciation to **Dr. Einas Mohamed Magib**, Lecturer of medical Physiology, Faculty of Medicine-Ain Shams University, for her precious help and beneficial advices throughout this thesis. It has been a great honor to work under her supervision.

The authors thank **Dr. Sayed Abdel Raheem**, Assistant professor of Histopathology at Al Azhar Faculty of Medicine, for scientific help in the histopathological assessment and **Dr. Laila Rashed**, Professor of Medical Biochemistry and Molecular Biology at Cairo University, faculty of medicine for scientific help in stem cell and biochemical studies.

Nahla Mohamed Hussien

### **List of Contents**

Title	Page No.
List of Tables	i
List of Figures	vi
List of Abbreviations	x
Introduction	1 -
Aim of the Work	4
Review of Literature	
Hypertension and its Cardiovascular Effects	5
Cardiac Ischemia Reperfusion Injury	19
Stem Cells	30
Materials and Methods	45
Results	76
Discussion	189
Summary and Conclusion	214
References	221
Arabic Summary	

### List of Tables

Table 1	Vo.	Title	Page No.
Table (1):		y Weight (IBW; gm), Final Body 'nge in body weights (% change) in	
<b>Table (2):</b>		y Weight (IBW; gm), Final Body 'nge in body weights (% change) in	
<b>Table (3):</b>		y Weight (IBW; gm), Final Body 'nge in body weights (% change)in H	
<b>Table (4):</b>	(FBW; gm)	data Initial Body Weight (IBW; g and percent change in body weight -SCs treated groups	s (% change) in Control,
<b>Table (5):</b>	(RVW/BW;	icular weight (RVW; mg), Right mg/gm), Left ventricular weight weight index (LVW/BW; mg/gm) in c	(LVW; mg) and Left
<b>Table (6):</b>	(RVW/BW; ventricular v	icular weight (RVW; mg), Right mg/gm), Left ventricular weight weight index (LVW /BW; mg/gm) in	(LVW; mg) and Left Ht-u group84
<b>Table (7):</b>	(RVW/BW ventricular	icular weight (RVW; mg), Right v; mg/gm), Left ventricular weight weight index (LVW /BW; mg/g	t (LVW; mg) and Left m) in Ht- SCs treated
<b>Table (8):</b>	weight inde and Left ve	data of Right ventricular weight (RV x (RVW/BW; mg/gm), Left ventricular weight index (LVW/BW; treated groups	cular weight (LVW; gm) mg/gm) in Control, Ht-u
<b>Table (9):</b>	pressure D (mmHg) re	n systolic blood pressure SBP (n DBP (mmHg) and mean arterial corded at the start of the study (ini- control group	blood pressure MAP itial), at 3 <sup>rd</sup> week and at
<b>Table (10):</b>	Changes in pressure D (mmHg) re-	n systolic blood pressure SBP (n DBP (mmHg) and mean arterial corded at the start of the study (Ini Ht-u group	nmHg), diastolic blood blood pressure MAP itial), at 3 <sup>rd</sup> week and at
<b>Table (11):</b>	pressure DI recorded at	n systolic blood pressure SBP (n BP (mmHg) and mean arterial blood the start of the study (initial), at 3 reated group.	d pressureMAP (mmHg)  B <sup>rd</sup> week and at 6 <sup>th</sup> week
<b>Table (12):</b>	Cumulative recorded at	data of changes in systolic blood the start of the study (initial), at 3 <sup>rd</sup> u and Ht- SCs treated groups	pressure SBP (mmHg) week and at 6 <sup>th</sup> week in
<b>Table (13):</b>	Cumulative recorded at	data of changes in diastolic blood the start of the study (initial), at 3 <sup>rd</sup> u and Ht- SCs treated groups	pressure (DBP, mmHg) week and at 6 <sup>th</sup> week in

Table N	No. Title	Page No.
<b>Table (14):</b>	Cumulative data of changes in mean blood recorded at the start of the study (initial), at control, Ht-u and Ht-SCs treated groups	3 <sup>rd</sup> week and at 6 <sup>th</sup> week in
Table (15):	ECG records: Heart Rate (HR; bpm), PR in QRS complex duration (QRS; msec), R volt observed duration (QTo; msec) and QT intervents in control group.	age (R; μvolt), QT interval val corrected duration (QTc;
<b>Table (16):</b>	ECG records: Heart Rate (HR; bpm), PR into QRS complex duration (QRS; msec), R volt observed duration (QTo; msec) and QT i (QTc; msec) in Ht-u group	age (R; µvolt), QT interval nterval corrected duration
<b>Table (17):</b>	ECG records: Heart Rate (HR; bpm), PR into QRS complex duration (QRS; msec), R volt observed duration (QTo; msec) and QT i (QTc; msec) inHt- SCs treated group	age (R; µvolt), QT interval nterval corrected duration
Table (18):	Cumulative data of ECG changes of Heart R duration (PR; msec), QRS complex duration µvolt), QT interval observed duration (QT corrected duration (QTc; msec) incontrol, groups.	(QRS; msec), R voltage (R; o; msec) and QT interval Ht-u and Ht- SCs treated
<b>Table</b> (19):	Chronotropic activity (heart rate per minute of heart rate (% change) at pre ischemia, at reperfusion calculated from isolated hearts in	t 5, at 15 and at 30 min of
<b>Table (20):</b>	Chronotropic activity (heart rate per minute of heart rate (% change) at pre ischemia, at reperfusion calculated from isolated hearts in	t 5, at 15 and at 30 min of
<b>Table (21):</b>	Chronotropic activity (heart rate per minute of heart rate (% change) at pre ischemia, a reperfusion calculated from isolated hearts in	t 5, at 15 and at30 min of
Table (22):	Cumulative data of chronotropic activity (h and percent change of heart rate (% change) and at 30 min of reperfusion calculated from Ht-u and Ht-SCs treated groups	at pre ischemia, at 5, at15 n isolated hearts in control,
<b>Table (23):</b>	Peak Tension (PT; gm) and percent change ischemia, at 5, at 15 and at 30 min of re isolated hearts in control group.	eperfusion calculated from
Table (24):	Peak Tension/Left Ventricular Weight (P percent change of (PT/LVW; (% change) hearts at pre ischemia, at 5min, at 15min at in control group.	) calculated from isolated nd at 30min of reperfusion

Table 1	Vo. Title	Page No.
Table (25):	Peak Tension (PT; gm) and Percent change of ischemia, at 5, at 15 and at 30 min of rep isolated hearts in Ht-u group	erfusion calculated from
<b>Table (26):</b>	Peak Tension/Left Ventricular Weight (PT percent change of (PT/LVW; % change) hearts at pre ischemia, at 5min, at 15min and in Ht-u group.	7/LVW; gm/100mg) and calculated from isolated at 30 min of reperfusion
<b>Table (27):</b>	Peak Tension (PT; gm) and Percent change of ischemia, at 5, at 15 and at 30 min of rep isolated hearts in Ht-SCs treated group	erfusion calculated from
<b>Table (28):</b>	Peak Tension/Left Ventricular Weight (PT Percent change of (PT/LVW; % change) calcusated pre ischemia, at 5, at 15min and at 30 min treated group	ulated from isolated heart of reperfusion in Ht-SCs
Table (29):	Cumulative data of Peak Tension (PT; gm) (PT;% change) calculated from isolated heart 15 and at 30 min of reperfusion in control, I groups.	sat pre ischemia, at 5, at Ht-u and Ht- SCs treated
<b>Table (30):</b>	Cumulative data of Peak Tension/Left Ventr gm/100mg)and percent change of (PT/LVW; 9 isolated heart sat pre ischemia, at 5, at 15 and a control, Ht-u and Ht-SCs treated groups	% change) calculated from t 30 min of reperfusion in
<b>Table (31):</b>	Time to Peak Tension (TPT; msec) and per change) at pre ischemia, at 5, at 15 and a calculated from isolated hearts in control grou	at 30 min of reperfusion
<b>Table (32):</b>	Half Relaxation Time (HRT; msec) and perchange) at pre ischemia, at 5, at 15 and a calculated from isolated hearts in Control grounds.	cent change of (HRT; % at 30 min of reperfusion
<b>Table (33):</b>	Time to Peak Tension (TPT; msec) and per change) at pre ischemia, at 5, at 15 and a calculated from isolated heart study in Ht-u gr	at 30 min of reperfusion
<b>Table (34):</b>	Half Relaxation Time (HRT; msec) and perchange) at pre ischemia, at 5, at 15 and at 30 mi from isolated heart study in Ht-u group	n of reperfusion calculated
<b>Table (35):</b>	Time to Peak Tension (TPT; msec) and per change) at pre ischemia, at 5, at 15 and a calculated from isolated heart study in Ht-SC:	at 30 min of reperfusion
<b>Table (36):</b>	Half Relaxation Time (HRT; msec) and per change) at pre ischemia, at 5, at 15 and a calculated from isolated heart study in Ht-SC:	at 30 min of reperfusion

Table N	Jo. Title	Page No.
F. 11 (25)		
Table (37):	Cumulative data of Time to Peak Tension (Tl change of (TPT; % change) at pre ischemia, at 5 reperfusion calculated from isolated heart stud Ht- SCs treated groups.	y in control. Ht-u and
Table (38):	Cumulative data of Half Relaxation Time (HRT change of (HRT; % change) at pre ischemia, at 5 reperfusion calculated from isolated heart study i SCs treated groups.	, at 15 and at 30 min of in control, Ht-u and Ht-
<b>Table (39):</b>	Myocardial Flow Rate (MFR; ml/min) and perce change) at pre ischemia, at 5, at 15 and at a calculated from isolated heart study in control gr	30 min of reperfusion roup 140
<b>Table (40):</b>	Myocardial Flow Rate/ Left Ventricular ml/min/100mg) and percent change of (MFRLV ischemia, at 5, at 15 and at 30 min of reperfusion heart study in control group.	VW; % change) at pre calculated from isolated
<b>Table (41):</b>	Myocardial Flow Rate (MFR; ml/min) and perce change) at pre ischemia, at 5, at 15 and at 30 min of from isolated heart study in Ht-u group	of reperfusion calculated
<b>Table (42):</b>	Myocardial Flow Rate/ Left Ventricular ml/min/100mg) and percent change of (MFRLV ischemia, at 5, at 15 and at 30 min of reperfusion heart study in Ht-u group	VW; % change) at pre calculated from isolated
<b>Table (43):</b>	Myocardial Flow Rate (MFR; ml/min) and perce change) at pre ischemia, at 5, at 15 and at a calculated from isolated heart study in Ht-SCs tr	30 min of reperfusion
<b>Table (44):</b>	Myocardial Flow Rate/ Left Ventricular Weight (MF and percent change of (MFR/LVW; % change) at proat 30 min of reperfusion calculated from isolated hea group.	e ischemia, at 5, at 15 and art study in Ht-SCs treated
Table (45):	Cumulative data of myocardial Flow Rate (MFI change of (MFR;% change)at pre ischemia, reperfusion calculated from isolated heart stud Ht-SCs treated groups	R; ml/min) and percent 5, 15 and 30 min of ly in control, Ht-u and
<b>Table (46):</b>	Cumulative data of myocardial Flow Rate/ Le (MFR/LVW; ml/min/100mg) and percent chan change) at pre ischemia, at 5, at 15 and at 30 min of from isolated heart study in control, Ht-u and Ht-Se	ge of (MFR/LVW; % of reperfusion calculated

Table N	lo. Title	Page No.
Table (47):	Lipid profile illustrating the plasma levels triglycerides TG (mg/dl), HDL (mg/dl) and l group.	LDL (mg/dl) in control
<b>Table (48):</b>	Lipid profile illustrating the plasma levels triglycerides TG (mg/dl), HDL (mg/dl) and group.	of cholesterol (mg/dl), LDL (mg/dl) in Ht-u
<b>Table (49):</b>	Lipid profile illustrating the plasma levels triglycerides TG (mg/dl), HDL (mg/dl) and L treated group	DL (mg/dl) in Ht- SCs
Table (50): Table (51):	Atherogenic Index (AI) in control, Ht-u and Ht- Cumulative data of lipid profile illustrating cholesterol (mg/dl), triglycerides TGC (mg/d	g the plasma level of
Table (52):	(mg/dl) and atherogenic Index (AI) in contitreated groups	rol, Ht-u and Ht- SCs155
Table (52):	groups	
<b>Table (53):</b>	Plasma levels of Malondialdehyde (MDA, nr and Ht- SCs treated groups	
Table (54):	Plasma levels of Super oxide dismutase (SOE and Ht- SCs treated groups	• •
Table (55):	Cardiac tissue levels of Malondialdehyde (Mincontrol, Ht-u and Ht- SCs treated groups	
Table (56):	Cardiac tissue levels of Superoxide dismutase incontrol, Ht-u and Ht-SCstreated groups	
Table (57):	Cumulative data of Plasma levels of Nitrites (µr (MDA, nmol/L), Super oxide dismutase (SOD, µ of Malondialdehyde (MDA, nmol/gm tissue) and Superoxide dismutase (SOD, nmol/gm tissue) in treated groups.	u/L), cardiac tissue levels d cardiac tissue levels of control, Ht-u and Ht-SCs
Table (58):	Significant correlation coefficient (p) between arterial pressure (MAP) and other parameters f Ht-SCs treated group and their matching control	n 6 <sup>th</sup> week (final) mean From pooled Ht-u group,
Table (59):	Significant correlation coefficient (p) plasma indicator of NO bioavailability) and other parar group, Ht-SCs treated group and their matching	levels of Nitrites (as an neters from pooled Ht-u
Table (60):	Significant correlations between plasma leve cardiac level of MDA and SOD and other parar group, Ht-SCs treated group and their matching	els of MDA and SOD, meters from pooled Ht-u

## List of Figures

Fig.	No.	Title Page No	).
Figure	(1):	Molecular mechanisms involved in the different types of monogenic hypertension (HTN).	
Figure	(2):	Factors increasing sympathetic activity in hypertension	9
Figure	e (3):	Ang II type 1 (AT <sub>1</sub> R) and type 2 (AT <sub>2</sub> R) receptors	11
Figure	e ( <b>4</b> ):	The actions of Ang II	11
Figure	(5):	A) Angiotensin II and inflammation increases ROS production by NADPH oxidases. B) Overproduction of mitochondrial superoxide and reduced SOD2 activity leads to mitochondrial oxidative stress	14
Figure	e ( <b>6</b> ):	Cross-talk between NADPH oxidase and mitochondria in endothelial dysfunction and hypertension	14
Figure	e ( <b>7</b> ):	Possible role of hypercholesterolemia in the pathogenesis of hypertension	16
Figure	2 (8):	Interactions between endothelial cell (EC) glycocalyx, endothelial Na+ channel (EnNaC), and eNOS (endothelial nitric oxide synthase) contribute to increases in endothelial stiffness	18
Figure	e ( <b>9</b> ):	Central role of calcium in the electro-mechanical dissociation of cardiomyocyte after myocardial reperfusion	21
Figure	e (10):	Inside cardiomyocyte, there is intracellular production of reactive oxygen species through NADPH oxidase, eNOS uncoupled, xanthine oxidase and mitochondrion	24
Figure	(11):	Sources of ROS in ischemia-reperfusion injury.	
Figure	(12):	Role of reactive oxygen species and iron mobilization in myocardial reperfusion injury and its clinical implications	
Figure	(13)	Cell death during ischemia reperfusion	
Figure	, ,	Autophagosome accumulation and cell death	
Figure		Classification of stem cells	
Figure	, ,	Paracrine immunomodulation in the infarcted myocardium	
Figure	, ,	Mechanisms of mesenchymal stem cells (MSCs) for cardiac regeneration	
Figure	(18):	Extracellular vesicles; exosomes and microvesicular bodies	
Figure	, ,	Vesicles from MSCs contain a variety of bioactive components	
Figure	, ,		
Figure	, ,	Isometric force transducer and the recorder	
Figure		Graph showing body weights (gms) all over the six weeks in all studied groups.	
Figure	(22 b):	Graph showing % changes of body weights in all studied groups	
Figure		Graph showing right ventricular weight (RVW, mgs), left ventricular weight (LVW, mgs) in all studied groups	

## List of Figures (Cont...)

Fig. No.	Title	Page No.
Figure (23 b):	Graph showing Right ventricular weight index (RVW/Fleft ventricular weight index (LVW/BW; mg/gm) in all st	
<b>Figure (24):</b>	Graph showing systolic blood pressure (SBP, mmhg) six weeks in all studied groups	
<b>Figure (25):</b>	Graph showing diastolic blood pressure (DBP, mmHg) a weeks in all studied groups	
<b>Figure (26):</b>	Graph showing mean arterial pressure (MAP, mmHg) alweeks in all studied groups	
<b>Figure (27a):</b>	Graph showing heart rate HR; bpm in all studied group	os 103
<b>Figure (27b):</b>	Graph showing R voltage (R; μvolt	
Figure (28):	Graph showing PR interval duration (PR; msec), Q duration (QRS; msec) QT interval observed duration and QT interval corrected duration (QTc; msec) ir groups	RS complex (QTo; msec) a all studied
<b>Figure (29):</b>	Graph showing isolated heart rate per minute (rpm) at baseline value and responses following ischemia and 3 reperfusion of hearts isolated from all studied groups	0 minutes of
<b>Figure (30):</b>	Graph showing Peak Tension (PT; gm) at preische value and responses following ischemia and 30 reperfusion of hearts isolated from all studied groups	mia baseline minutes of
<b>Figure (31):</b>	Graph showing Peak Tension per left ventricular weight (F at pre ischemia baseline value and responses following isc minutes of reperfusion of hearts isolated from all studied gro	hemia and 30
Figure (32):	Graph showing time to peak tension (TPT; msec) at baseline value and responses following ischemia and 3 reperfusion of hearts isolated from all studied groups	0 minutes of
Figure (33):	Graph showing Half Relaxation Time (HRT; msec) a baseline value and responses following ischemia and 3 reperfusion of hearts isolated from all studied groups	0 minutes of
<b>Figure (34):</b>	Graph showing myocardial flow rate (ml/min) at paseline value and responses following ischemia and 3 reperfusion of hearts isolated from all studied groups	0 minutes of
Figure (35):	Graph showing myocardial flow rate per left ventri (MFR/LVW ml/min/100mg) at pre ischemia baselin responses following ischemia and 30 minutes of reperfuisolated from all studied group	e value and sion of hearts
<b>Figure (36):</b>	Graph showing plasma levels of total choles Triglycerides (TG) and Low Density lipoprotein (LD all studied groups.	L) values in
<b>Figure (37):</b>	Graph showing plasma levels of high Density lipopr values in all studied groups.	

## List of Figures (Cont...)

Fig. No.	Title	Page No.
Figure (38):	Graph showing calculated Atherogenic Indestudied groups.	
<b>Figure (39):</b>	Graph showing plasma levels of nitrites in all s	studied groups 163
<b>Figure (40):</b>	Graph showing plasma levels of MDA values in	all studied groups 164
<b>Figure (41):</b>	Graph showing plasma levels of SOD values in a	ıll studied groups 164
<b>Figure (42):</b>	Graph showing Cardiac levels of MDA values in	all studied groups 164
<b>Figure (43):</b>	Graph showing Cardiac levels of SOD values in all	studied groups164
<b>Figure (44):</b>	Graph showing significant correlation coefficient (final) mean arterial pressure (MAP) and other pooled Ht-u group, Ht-SCs treated group controls	her parameters from and their matching
Figure (45):	Graph showing significant correlation coef plasma level of Nitrites and other parameter group, Ht-SCs treated group and their matching	ficient (p) between rs from pooled Ht-urg controls168
<b>Figure (46):</b>	Graph showing significant correlation coefficient level of Nitrites and other parameters from poole treated group and their matching controls	ed Ht-u group, Ht-SCs
<b>Figure (47):</b>	Graph showing significant correlation coef plasma level of Nitrites and other parameter group, Ht-SCs treated group and their matching	rs from pooled Ht-u
<b>Figure (48):</b>	Graph showing significant correlation coeplasma level of MDA and other parameters group, Ht-SCs treated group and their matching	s from pooled Ht-u
<b>Figure (49):</b>	Heart showing average pericardium, average c and average blood vessels in control group	
<b>Figure (50):</b>	Another view showing average muscle fiber borders, and average blood vessel in control gr	
<b>Figure (51):</b>	Heart showing markedly dilated congested s vessel, and scattered muscle fibers with cytoplasm in Ht-u group	bright eosinophilic
<b>Figure (52):</b>	Another view showing markedly dilated coblood vessel, and scattered muscle fibers with cytoplasm and no nuclei, and intra-cytoplasm group.	h bright eosinophilic nic vacuoles in Ht-u
<b>Figure (53):</b>	High power view showing average myocardiaverage muscle fibers with distinct cell borde in Ht-SCs treated group	rs and central nuclei
<b>Figure (54):</b>	Aortic wall showing average intima, average adventitia in control group.	