

بسم الله الرحمن الرحيم





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات




يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بعض الوثائق الأصلية تالفة





بالرسالة صفحات
لم ترد بالأصل





Ain Shams University
Faculty of Medicine
Anesthesiology, Intensive Care and Pain Management

A Comparative Study between Dexamethasone versus Paracetamol Added to Xylocaine in Local Intravenous Anesthesia in Upper Limb Minor Surgeries

A Thesis

Submitted for the Partial Fulfillment of Master's Degree

In Anesthesiology, Intensive Care and Pain Management

BY

Mostafa Mohamed Mohamed Ibrahim Sharaf

M.B.B.CH Alexandria University

Supervisors

Prof. Reem Hamdy Elkabarity

Professor of Anesthesiology, Intensive Care and Pain Management
Faculty of Medicine – Ain Shams University

Dr. Heba Abdel Azim Labib

Assistant Professor of Anesthesiology, Intensive Care and Pain Management
Faculty of Medicine – Ain Shams University

Dr. Marwa Mamdouh Elfar

Lecturer in Anesthesiology, Intensive Care and Pain Management
Faculty of Medicine – Ain Shams University

**Ain Shams University
Faculty of Medicine**

2020

CONTENTS

Title	Page
List of Tables	i
List of Figures	ii
List of Abbreviations	iii
Introduction	1
Aim of the Work	3
Review of Literature	
- Anatomy of upper limb	4
- Intravenous regional anesthesia	16
- Pharmacology of drugs	24
Patients and Methods	43
Results	50
Discussion	71
Summary	76
References	78
Arabic Summary	85

LIST OF TABLES

Table No.	Title	Page
Table (1)	Modified Bromage Scale for Assessment of Motor Block	46
Table (2)	Demographic data in the studied groups, ASA, Surgery time ,type of surgery	50
Table (3)	Comparison between the three studied groups according to Mean arterial blood pressure	51
Table (4)	Comparison between the three studied groups according to heart rate	53
Table (5)	Comparison between the three studied groups according to oxygen saturation	55
Table (6)	Comparison between the three studied groups according to onset and recovery of sensory block	57
Table (7)	Comparison between the three studied groups according to onset and recovery of motor block	59
Table (8)	Intensity Of motor Block according to Modified Bromage scale	61
Table (9)	Assessment of post-operative pain visual analogue scale (VAS score)	63
Table (10)	Time for first rescue dose of analgesia	65
Table (11)	Comparison between the three studied groups according to patient satisfaction	67
Table (12)	Comparison between the three studied groups according to surgeon`s opinion	69

LIST OF FIGURES

Fig. No.		Page
Fig. (1)	Muscles of the upper limb	6
Fig. (2)	Arterial supply of the upper limb	8
Fig. (3)	Superficial (Lt) and Deep (Rt) veins of upper limb	11
Fig. (4)	Nerve supply of upper limb	15
Fig. (5)	IVRA Technique.	20
Fig. (6)	Chemical structure of xylocaine	24
Fig. (7)	Chemical structure of Paracetamol	30
Fig. (8)	Prostaglandin inhibition pathway	32
Fig. (9)	Conversion of Paracetamol to AM404, endocannabinoid reuptake inhibitor	34
Fig. (10)	Chemical structure of Dexamethasone	38
Fig(11)	Visual Analogue scale	47
Fig. (12)	Comparison between the three studied groups according to mean arterial blood pressure	52
Fig. (13)	Comparison between the three studied groups according to heart rate	54
Fig. (14)	Comparison between the three studied groups according to SpO ₂	56
Fig. (15)	Comparison between the three studied groups according to sensory block	58
Fig. (16)	Comparison between the three studied groups according to motor block	60
Fig. (17)	Comparison between the three studied groups according to Intensity Of motor Block: According to Modified Bromage scale	62

Fig. (18)	Comparison between the three studied groups according to post-operative pain score (VAS score)	64
Fig. (19)	Comparison between the three studied groups according to time for first opioid dose	66
Fig. (20)	Comparison between the three studied groups according to Patient satisfaction	68
Fig. (21)	Comparison between the three studied groups according to surgeon`s opinion	70

