

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY

Assessment of Mothers Care towards their Children Suffering from Branched Chain Keto-Aciduria (BCKA)

Thesis

Submitted for Partial Fulfillment of the Requirements for Master Degree in Pediatric Nursing

By Haniem Mahmoud Said

(B.Sc. Nursing Sciences 2005)

Educator in Education Unit

Pediatric Hospital
Ain Shams University

Faculty of Nursing
Ain Shams University
2020

Assessment of Mothers Care towards their Children Suffering from Branched Chain Keto-Aciduria (BCKA)

Thesis

Submitted for Partial Fulfillment of the Requirements for Master Degree in Pediatric Nursing

Supervised by

Prof. Dr/ Sabah Saad Al-Sharkawi

Professor of Pediatric Nursing
Faculty of Nursing- Ain Shams University

Prof. Dr/ Asmaa Nasr-Eldin Mosbeh

Professor of Pediatric Nursing
Faculty of Nursing- Ain Shams University

Prof. Dr/ Osama Kamal Zaki

Professor and Medical Genetics
Faculty of Medicine- Ain Shams University

Faculty of Nursing
Ain Shams University
2020

Acknowledgments

First and forever, thanks to **ALLAH**, Almighty for giving the strength and faith to complete my thesis.

I am deeply grateful to **Prof. Dr. Sabah** Saad Al-Sharkawi, Professor of Pediatric Nursing, Faculty of Nursing; Ain shams University for her constructive criticism, unlimited help and giving me the privilege to work under her supervision.

I would like to express my sincere gratitude to **Prof. Dr. Asmaa Nasr-Eldin Mosbeh,** Professor of Pediatric Nursing, Faculty of Nursing, Ain shams University for her supervision, efforts and time she has devoted to accomplish this work.

I would like to express my deep appreciation and gratitude to **Prof. Dr. Osama Kamal Zaki,** Professor of Clinical Genetics, Genetics Unit, Pediatrics Department, and Faculty of Medicine-Ain Shams University for his supervision, guidance and support throughout this study.

I would like also to express my gratitude to all mothers and their children who participated as study subjects for their cooperation.

Last but not least, I can't forget to thank all my Family for their kind care, help and encouragement.

≥ Haniem Mahmoud Said

List of Contents

Subject Page No.
AbstractI
List of AbbreviationsIII
List of TablesVII
List of FiguresVIII
Introduction and Aim of the Study1
Review of Literature
Overview about the Branched Chain Keto Aciduria 5
Nutritional Requirements for children suffering from Branched Chain Keto Aciduria
Needs for children suffering from Branched Chain Keto Aciduria
Pediatric Nursing Role and Interventions 61
Subjects and Methods75
Results 83
Discussion
Conclusion
Recommendations
Summary
References
Appendices
Arabic Summary

Abstract

Branched-chain Keto-aciduria (BCKA) known as Maple Syrup Urine Disease (MSUD) is an aminoacidpathy due to an enzyme defect in the catabolic pathway of the branched-chain amino acids leucine, isoleucine, and valine. This defect leads to particularly toxic to the central nervous system. Aim: This study aims to assess mothers care for children having BCKA, through assess the mother's knowledge and self-reported practices toward caring for their children with BCKA. **Design:** A descriptive study. sample which included A purposive 100 accompanying their children with BCKA. Setting: This study was conducted at Specialized Genetic Outpatient Clinic at Children's Hospital affiliated to Ain Shams University Hospital. Tools: Tool I: A structure interview questionnaire format to assess characteristics of the study children and their mothers, medical history of children with BCKA, physical and psychological children's health needs and problems, mother's knowledge about BCKA. Tool II: Mothers self-reported practices related to care of their children with BCKA. Tool III: Medical records. Results: This study revealed that the mean age of studied children's was $3.82 \pm$ 2.97 years and the mean age of studied mothers was 33.8 ± 1.35 years. Meanwhile about three quarters of studied mothers had unsatisfactory knowledge about BCKA, as well more than two thirds of them had poor total reported practice. The highest total knowledge and total self reported practices mean score are for mother whose young age, high education and living in urban area. Conclusion: This result concluded that majority of mothers had unsatisfactory total knowledge and poor total self-reported practices about branched-chain keto-aciduria. There were highly significant relation between the mother's characteristics and their care for children suffering from BCKA. **Recommendation:** Teaching program for families having children with BCKA is important to raise their knowledge about the disease, its management and prevention of complications. Future research should be focused on measuring the prevalence of BCKA in Egypt and investigating the risk health problems among mothers of children with BCKA.

Keywords: Branched Chain Keto-Aciduria, Maple syrup urine disease, Problems, Mothers, Children, Care, Amino acids.

List of Abbreviations

Abbreviations Full terms,

AA : Amino Acid

ADHD : Attention-Deficit/Hyperactivity Disorder

αkIC : Alpha-Ketoisocaproic

ASD : Autism Spectrum Disorder
 ASL : Argininosuccinate Synthetase
 BCAAs : Branched Chain Amino-Acids
 BCHA : Branched-Chain Hydroxy Acids
 BCKA : Branched Chain Keto-Aciduria

BCKAD : Branched-Chain α-Ketoacid Dehydrogenase
 BCKD : Branched-Chain α-Ketoacid Dehydrogenase
 BCKDC : Branched-Chain α-Ketoacid Dehydrogenase

Complex

BCKDHA : Branched-Chain α-Ketoacid Dehydrogenase

E1 Alpha polypeptide

BCKDHB: Branched-Chain α-Ketoacid Dehydrogenase

E1 Beta polypeptide

BMI : Body Mass Index

BDI : Beck Depression Inventory Scale

CoA : Co-enzyme Aminoacid

CPS: Carbamyl Phosphate Synthetase

CSF : Cerebrospinal Fluid

DBT : Dihydrolipoamide Branched Chain

Transacylase

DLDD : Dihydrolipoamidedehydrogenase deficiency

DNPH : Dinitrophenylhydrazine
 DRI : Daily Requirements Intake
 EEGs : Electroencephalograms

ESMSUD: European Society of Maple Syrup Urine

Disease

GC-MS : Gas Chromatography-Mass Spectrometry

GLU : Glutamate

GMP : Glycomacropeptide

GTP : Guanosine Triphosphate

HCL: Hydrochloric acid

Hrs: Hours

IEM : Inborn Error of MetabolismIMDs : Inherited Metabolic Disorders

IDSA : Infectious Diseases Society of America

ILE : Isoleucine ISO-LEU : Isoleucine

Intelligence quotient
 α-Ketoisocaproic acid
 α-Ketoisovaleric acid

KMV : α-Keto-β-methylvaleric acid

LEU : Leucine

LNAAMCQMultiple Choice QuestionsMMAMethyl Malonic Acidemia

MMAA : Methyl Malonic Aciduria type A poroteinMMAB : Methyl Malonic Aciduria type B porotein

MMADHC: Methyl Malonic Aciduria and

Homocyctinuria type D porotein

MCEE : Methylmalonyl CoA mutase

Mmol : Millimoles

MS : Mass Spectrometry

MUT : Mutations

MSUD : Maple Syrup Urine Disease
 MRI : Magnetic Resonance Imaging
 NAGS : N-acetyl-Glutamate Synthetase

NBS : Newborn Screening

NIH : National Institute of Health

NORD : National Organization for Rare Disorders

OTC : Ornithine Transcarbamylase

PAA : Plasma Amino Acids

PRO: Protein

TPN : Total Parental NutritionSD : Standard Deviation

STDs : Sexually Transmitted Disease

SPSS : Statistical Package for Social Science

TPN: Total parental nutrition

Tyr : Tyrosine VAL : Valine

U-MOL : Unit Moles

UOAUrine Organic AcidsUSAUnited State of AmericaWHOWorld Health Organization

QOL : Quality of Life

List of Tables

Table T	Vo.	Title	Page No.
Tables i	in Revie	<u>w</u>	
(1):	The cha	nce of BCKA occ	urrence11
(2):		Phenotypes of ciduria	Branched-Chain21
(3):	patients	nended daily nut with Branch a	
(4):	branche	nended intake ed-chain amino a calories	1 '

Tables in Results

Table No.	Title	Page	No.
(1):	Distribution of the studied chi according to their characteristics 100).	(N=	84
(2):	Distribution of the studied mode according to their characteristics 100).	(N=	85
(3):	Number and percentage distribution the studied children as regards history of illness (N=100).	their	86
(4):	Number and percentage distribution the studied children according to family history of the disease ($N = 1$)	their	88
(5):	Number and percentage distribution the studied children according to physical assessment (N=100)	their	89
(6):	Number and percentage distribution the studied children according to health problems (N=100)	their	90
(7):	Number and percentage distribution the studied children according to health needs (N=100)	their	91
(8):	Number and percentage distribution the studied children regarding laboratory investigation (N=100).	thier	92

(9):	Distribution of the studied mothers as regards their knowledge about branched chain keto-aciduria (N= 100).	93
(10):	Number and percentage distribution of the mothers as regards their knowledge about care for their children with branched chain keto-aciduria (N=100)	94
(11):	Distribution of the studied mothers' in relation to their knowledge regarding proper feeding for their children with branched chain keto-aciduria (N= 100)	95
(12):	Number and percentage distribution of the studied mothers' self reported practices about measuring diet at home for their children with branched chain keto-aciduria (N=100)	97
(13):	Number and percentage distribution of the studied mothers' self reported practices about dental care for their children with branched chain keto-aciduria (N=55)	98
(14):	Number and percentage distribution of the studied mothers' self reported practices about nasogastric tube feeding for their children with branched chain	
	keto-aciduria (N=30)	99

(15):	Number and percentage distribution of the studied mothers' self reported practices about measuring weight for their children with branched chain ketoaciduria (N=100)
(16):	Number and percentage distribution of the studied mothers' self reported practices regarding to physical exercise for their children with branched chain ketoaciduria (N=100)
(17):	Number and percentage distribution of the mothers' self reported practices about measuring acetone in the urine for their children with branched chain ketoaciduria (N=100)
(18):	Number and percentage distribution of the studied mothers about their self reported practices about daily observation of childrens' symptoms (N=100)
(19):	Relation between characteristics of studied mothers and their total knowledge regarding to branched chain ketoaciduria
(20):	Relation between the studied mothers' characteristics and their total self reported practices level
(21):	Relation between the mothers' total self reported practices and their total knowledge