



شبكة المعلومات الجامعية  
التوثيق الإلكتروني والميكرو فيلم

# بسم الله الرحمن الرحيم



**HANAA ALY**



شبكة المعلومات الجامعية  
التوثيق الإلكتروني والميكروفيلم



# شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلم



**HANAA ALY**



شبكة المعلومات الجامعية  
التوثيق الإلكتروني والميكروفيلم

# جامعة عين شمس

## التوثيق الإلكتروني والميكروفيلم

### قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها  
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



### يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



**HANAA ALY**

# **MANAGEMENT OF FERTIGATION UNDER ARID CONDITION**

**By**

**SHERIN AHMED MAHMOUD BAKR**

B.Sc. (Agric. Eng.), Fac. Agric., Cairo University, (1999)

M.Sc. (Agric. Sci.), Environmental Science, Ain Shams University, (2010)

**A Thesis Submitted in a Partial Fulfillment  
Of  
the Requirements for Degree of**

**DOCTOR OF PHILOSOPHY  
in  
Agricultural Sciences  
(Bio Engineering Systems in Arid Area)**

**Department of Arid Lands  
Faculty of Agriculture  
Ain Shams University**

**2021**

# **MANAGEMENT OF FERTIGATION UNDER ARID CONDITION**

By

**SHERIN AHMED MAHMOUD BAKR**

B.Sc. (Agric. Eng.), Fac. Agric., Cairo University, (1999)

M.Sc. (Agric. Sci.), Environmental Science, Ain Shams University, (2010)

**Under the supervision of:**

**Dr. Abd El- Ghany Mohamed El- Gindy**

Professor of Emeritus Agricultural Engineering, Agricultural Engineering Department, Faculty of Agricultural, Ain Shams University. (Principal Supervisor).

**Dr. Khled Faran Taher El- Bagoury**

Professor of Agricultural Engineering, Agricultural Engineering Department, Faculty of Agriculture, Ain Shams University.

**Dr. Wael Mahmoud Mokhtar Sultan**

Head of Res. in Agricultural Engineering, Agricultural Engineering Research Institute, ARC.

**Approval Sheet**

**MANAGEMENT OF FERTIGATION UNDER ARID  
CONDITION**

By

**SHERIN AHMED MAHMOUD BAKR**

B.Sc. (Agric. Eng.), Fac. Agric., Cairo University, (1999)

M.Sc. (Agric. Sci.), Environmental Science, Ain Shams University, (2010)

**This thesis for a Ph.D. degree has been approved by:**

**Dr. Asaad Abdelkader Derbala** .....

Prof. of Agric. Eng., Faculty of Agriculture, Tanta University.

**Dr. Usama Ahmed Ali El-Behairy** .....

Prof. of Horticulture, Faculty of Agriculture, Ain Shams University.

**Dr. Khaled Faran Taher El- Bagoury** .....

Prof. of Agric. Eng., Faculty of Agriculture, Ain Shams University.

**Dr. Abd El- Ghany Mohamed El- Gindy** .....

Prof. Emeritus of Agric. Eng. Faculty of Agriculture, Ain Shams  
University.

**Date of examination:**    /    / 2020

## ABSTRACT

**Sherin Ahmed Mahmoud Bakr. Management of Fertigation under Arid Condition. Unpublished Doctor of Philosophy dissertation. Ain Shams University, Faculty of Agriculture, Department of Arid Lands, 2021.**

Excessive use of chemical fertilizers leads to soil destruction, water and environmental pollution, reduced productivity, low product quality, water distribution and fertilizer efficiency. Good irrigation and fertilization systems management methods is very important to prevent soil and plant conservation and achieving highest efficiency of water and fertilizer use ,to produce high quality and quantity crops for increasing local production and export. Onion and cowpea are important crops in Egyptian cultivators. Onion is strategy crops for export, and cowpea is called a poor men's meat. The current study was conducted through two open field experiments on onion and cowpea crops and seasons, to manage suitable fertigation programs through fertilizer type, form, and time vs. emitters type and discharges and to investigate their impacts on some plant attributes, yield and irrigation water productivity.

The first Experiment was carried out from open field in private farm for vegetables and fruits production, located in Hosen sector at Alexandria Cairo desert road km 70 from Cairo, El-Behiara governorate for Onion seeds, design for the first experiment was a split plot design with four replicates and two factors. Main factor : effect of fertilizer application method “F<sub>m</sub>” : Cattle manure (F<sub>1</sub>), mixing of cattle manure and chemical fertilizer “fertigation” with rate of 30 :70 (F<sub>2</sub>) and chemical fertilizer “fertigation” (F<sub>3</sub>). Second factor: emitters type: OT<sub>4</sub> ”I<sub>1</sub>”, emitters built in drip line ( GR 16) “I<sub>2</sub>” and emitters, anti roots “I<sub>3</sub>” and each sub-plot is was 3.5 m long and 3 m., wide (10.5m<sup>2</sup> ie. 1/400 fed. ) and include 6 rows 50 cm wide, each 3.5 m long and the distance between sub-plots is left as 1 m to create a buffer zone

Meanwhile, the second experiment was carried out from open field in Agricultural Engineering Research Instute, Ministry of

Agriculture (AEnRI), Giza governorate, For cowpea dry seeds, the experiment included eight treatments were Emitter discharge “q” {12 L.h<sup>-1</sup> (E<sub>1</sub>), 8 L.h<sup>-1</sup> (E<sub>2</sub>)}, Fertigation form “F<sub>F</sub>” { Liquid (L<sub>F</sub>), Powder (P<sub>F</sub>)} and Fertigation time “F<sub>t</sub>” { Before irrigation (B<sub>I</sub>), After irrigation (A<sub>I</sub>)}. At the first experiment results elucidated that applying mixing of cattle manure and fertigation through anti roots emitter (F<sub>2</sub>I<sub>3</sub>) treatment, obtained highest values of bulb diameter “d<sub>1</sub>” (13.88cm), bulb diameter “d<sub>2</sub>” (25.10cm), bulb mass “B<sub>m</sub>” (195.9g), bulb actual volume “B<sub>v</sub>” (118cm<sup>3</sup>), moisture content “M<sub>C</sub>” (99.52 %), bulb yield “B<sub>y</sub>” (15.01Mg. fed<sup>-1</sup>), and irrigation water productivity “IWP” (6.72 kg.m<sup>-3</sup>), respectively. Meanwhile, by applying cattle manure only through OT<sub>4</sub> emitter (F<sub>1</sub>I<sub>1</sub>) obtained the lowest values of d<sub>1</sub> (6.24cm), d<sub>2</sub> (5.08cm), B<sub>m</sub> (95.6g), B<sub>v</sub> (79cm<sup>3</sup>), M<sub>C</sub> (70.13 %), B<sub>y</sub> (7.80Mg. fed<sup>-1</sup>), and IWP (3.49 kg.m<sup>-3</sup>), respectively.

At the second experiment results elucidated that applying powder fertilizer (after irrigation) (P<sub>F</sub> A<sub>I</sub>) through emitter with 12 Lh<sup>-1</sup> discharge (E<sub>1</sub>) treatment, obtained the lowest values of number of pods / plant “P<sub>n</sub>” (5.6 Pods. plant<sup>-1</sup>), number of seeds / pod “S<sub>n</sub>” (6.4 seeds. pod<sup>-1</sup>), seed mass /pod “S<sub>m</sub>” (0.97 g), number of seed index “S<sub>I</sub>” (15.1 g), seed yield “Y<sub>s</sub>” (0.79 Mg. fed<sup>-1</sup>), irrigation water productivity “IWP” (3.27kg.m<sup>-3</sup>). On the other hand, by applying liquid fertilizer (before irrigation) (L<sub>F</sub> B<sub>I</sub>) treatment through emitter with 8 Lh<sup>-1</sup> discharge (E<sub>2</sub>), obtained the highest values of P<sub>n</sub> (5.3 Pods. plant<sup>-1</sup>), S<sub>n</sub> (6.9 seeds. pod<sup>-1</sup>), S<sub>m</sub> (0.97 g), S<sub>I</sub> (14.1 g), Y<sub>s</sub> (0.75 Mg. fed<sup>-1</sup>), and “IWP” (4.66kg.m<sup>-3</sup>).

**Keywords:** fertigaion - drip irrigation - cowpea - onion -irrigation water productivity - yield.



## CONTENTS

	<b>Page</b>
<b>LIST OF TABLES</b>	<b>IV</b>
<b>LIST OF FIGURES</b>	<b>V</b>
<b>LIST OF PHOTOS</b>	<b>VII</b>
<b>LIST OF ABBRIVATION</b>	<b>VIII</b>
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 REVIEW OF LITERATURE</b>	<b>3</b>
2.1 Management of drip irrigation system.	3
2.2 Water use efficiency and irrigation water productivity.	5
2.3 Effect of drip irrigation systems on water and fertilizer use efficiency.	5
2.4 Studies of fertigation management technology.	7
2.5 Soil moisture distribution.	11
2.6 Management of Nutrient uptake.	12
2.7 Management of crop growth, development, and yield.	15
2.7.1 Onion.	15
2.7.2 Cowpea.	16
<b>3 MATERIALS AND METHODS</b>	<b>21</b>
3.1 Materials	21
3.1.1 Irrigation and fertigation systems.	21
3.2 Nutrient management.	23
3.2.1 Compost.	23
3.2.2 Cattle manure.	24
3.2.3 NPK fertilizer application.	24
3.3 Soil analysis.	25
3.4 Planting procedures.	26
3.5 Experimental treatments and design.	28
3.6 Emitters Evaluation.	32
3.6.1 Hydraulic system.	32
3.6.2 Water temperature and filtration.	35
3.6.3 Supporting frame.	35
3.6.4 Catch can water from emitters.	35
3.6.5 Test procedures.	35
3.6.5.1 Emitter discharge.	36
3.6.5.2 Emitter manufacture's coefficient of variations.	36

3.6.5.3	Emission uniformity.	37
3.6.5.4	Statistical uniformity coefficient.	37
3.6.5.5	Discharge variation.	37
3.7	Studied attributes.	38
3.8	Instruments.	42
3.9	Scheduling of irrigation and fertigation.	42
3.9.1	Water requirement	42
3.9.2	Irrigation interval	44
3.9.3	Scheduling of irrigation	44
3.9.3.1	Irrigation timing	44
3.9.3.2	Calculation of irrigation timing according to the evapo-transpiration (ET) for the experimental area.	44
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	<b>46</b>
4.1	Effect of fertilizer methods and emitters type on onion bulb parameters.	46
4.1.1	Bulb Diameters.	46
4.1.2	Bulb mas.	48
4.1.3	Bulb actual volume.	49
4.1.4	Moisture Content.	50
4.1.5	Bulb yield.	51
4.1.6	Irrigation water productivity.	51
4.1.7	Correlation between onion parameters.	55
4.2	Effect of emitter discharge, fertigation form and fertigation timing on growth vegetative of cowpea plant.	55
4.2.1	Effect of fertigation form and fertigation timing on growth vegetative of cowpea plant at emitter discharge ( $12 \text{ l h}^{-1}$ ).	55
4.2.1.1	Number of Pods.	55
4.2.1.2	Number of Seeds.	57
4.2.1.3	Seed Mass.	57
4.2.1.4	Seed Index.	59
4.2.1.5	Seed yield.	60
4.2.1.6	Irrigation Water Productivity	60
4.2.1.7	Correlation between cowpea parameters at emitter discharge of ( $12 \text{ L.h}^{-1}$ ).	61
4.2.2	The relationship between fertigation form and fertigation timing on growth vegetative of cowpea plant at emitter discharge ( $8 \text{ l h}^{-1}$ ):	61
4.2.2.1	Number of Pods	61
4.2.2.2	Number of Seeds	63

4.2.2.3	Seed Mass	64
4.2.2.4	Seed Index	65
4.2.2.5	Seed yield	66
4.2.2.6	Irrigation Water productivity	66
4.2.2.7	Correlation between cowpea parameters at emitter discharge of (8 L.h <sup>-1</sup> ).	67
4.3	Statistical analysis.	67
4.3.1	Variables definition.	68
4.3.1.1	Effect of fertilizer methods on onion bulb parameters, as affected by three emitters type.	68
4.3.1.2	Effect of fertigation form and fertigation timing on growth vegetative of cowpea plant at emitter discharge (12 L.h <sup>-1</sup> ) and (8 L.h <sup>-1</sup> ).	69
4.3.2	Statistical analysis for onion bulb parameters.	69
<b>5</b>	<b>SUMMARY AND CONCLUSION</b>	<b>73</b>
<b>6</b>	<b>REFERENCES</b>	<b>78</b>
<b>7</b>	<b>APPENDIX</b>	<b>93</b>
<b>8</b>	<b>ARABIC SUMMARY</b>	

## LIST OF TABLES

	<b>Page</b>
3.1 Chemical composition of well water used for irrigation in the first experiment.	21
3.2 Chemical analysis of feculent used for irrigation in the second experiment.	23
3.3 Chemical analysis for compost.	24
3.4 Chemical analysis for cattle manure.	24
3.5 Physical-chemical properties of two experimental soil.	26
3.6 The meteorological measurements monthly during 2016, 2017 and 2018 growing season.	26
3.7 Discharge “q” (nominal, mean and standard deviation “s”, manufacture’s coefficient of variation “CV” (%) and some hydraulic characteristics,i.e., emission uniformity “Eu” (%), statistical uniformity “Us”(%), and discharge variation “q <sub>var</sub> ” (%) for tested emitters at 100 kpa.	41
3.8 Data for calculate total applied water "TAW" Reference evapotranspiration, onion crop.	43
3.9 Data for calculate total applied water "TAW" Reference evapotranspiration, cowpea crop.	44
4.1 Fertilizer methods and bulb diameters “d <sub>1</sub> and d <sub>2</sub> ”, as affected by three emitters.	46
4.2 Relations between fertilizer methods and bulb mass, bulb actual volume and moisture content as affected by three emitters type.	49
4.3 Relations between fertilizer methods, bulb yield and irrigation water productivity as affected by three emitters type.	52
4.4 Relations between fertigation form and fertigation timing on growth vegetative Index on cowpea plant at emitter discharge of (12 L.h <sup>-1</sup> ).	58
4.5 Relations between fertigation form and fertigation time on growth vegetative on cowpea plant at emitter discharge of (8 L.h <sup>-1</sup> ).	62
4.6 Variables values of onion bulb parameters at fertilizer	71

methods and emitters type.

- 4.7 Variables values of growth vegetative of cowpea plant at fertigation form and fertigation time through emitter discharge ( $12 \text{ L.h}^{-1}$ ) and ( $8 \text{ L.h}^{-1}$ ). 72

## LIST OF FIGURES

	<b>Page</b>
3.1 The first experiment layout.	30
3.2 Irrigation system layout for Exp.1.	31
3.3 The second experiment layout	33
3.4 Hydraulic test bench components	34
3.5 Cross section of emitters used in present study.	39
3.6 Cross section of emitters used in present study.	40
4.1 Relations between fertilizer methods and bulb diameters ( $d_1$ and $d_2$ ) as affected by three emitters type.	47
4.2 Relations between fertilizer methods and Bulb mass , Bulb actual volume and moisture content as affected by three emitters type.	47
4.3 Relations between fertilizer methods and bulb yield and irrigation water productivity as affected by three emitters type.	53
4.4 Correlation between bulb diameter " $d_1$ " and bulb diameter " $d_2$ ", bulb yield and irrigation water productivity.	56
4.5 Correlation between bulb diameter " $d_1$ " and bulb mass, bulb actual volume and moisture content.	56
4.6 The relations between fertigation form and fertigation timing on seed index and its effect at emitter discharge of $12 \text{ L.h}^{-1}$ ).	58
4.7 The relations between fertigation form and fertigation timing on number of pods / plant, irrigation water productivity and number of seeds / pod and its effect at emitter discharge of ( $12 \text{ L.h}^{-1}$ ).	59
4.8 The relations between fertigation form and fertigation timing on seed mass /pod and seed yield and its effect at emitter discharge of ( $12 \text{ L.h}^{-1}$ ). Correlation between seed index and both seed yield and irrigation water productivity and at emitter	60

	discharge of (12 L.h <sup>-1</sup> ).	
4.9	The relations between fertigation form and fertigation timing on seed index and its effect at emitter discharge of (8 L.h <sup>-1</sup> ).	62
4.10	The relations between fertigation form and fertigation timing on seed mass /pod and seed yield and its effect at emitter discharge of (8 L.h <sup>-1</sup> ).	63
4.11	The relations between fertigation form and fertigation timing on on number of pods / plant,irrigation water roductivity and number of seeds / pod and its effect at emitter discharge of (8 L.h <sup>-1</sup> ).	64
4.12	Correlation between seed index and both seed yield and irrigation water productivity at emitter discharge of (8 L.h <sup>-1</sup> ).	65
4.13	Correlation between seed index and both seed yield and irrigation water productivity at emitter discharge of (8 L.h <sup>-1</sup> ).	68

## **LIST OF PHOTOS**

	<b>Page</b>
3.1     Onion field experiment	29
3.2     Cowpea field experiment	29