

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY

MICROPROPAGATION OF Loropetalum chinense Plant

By

AHMED FARGHALY HASSANEAN

B.Sc. Agric. Sci. (Horticulture), Fac. Agric., Alazhar Univ., 2001M.S.C. Agric. Sci. (Ornamental Horticulture), Fac. Agric., Cairo Univ., 2015

THESIS

Submitted in Partial Fulfillment of the Requirements for the Degree of

DOCTOR OF PHILOSOPHY

In

Agricultural Sciences (Ornamental Horticulture)

Department of Ornamental Horticulture Faculty of Agriculture Cairo University EGYPT

2020

Format reviewer

Vice Dean for graduate studies

APPROVAL SHEET

MICROPROPAGATION OF Loropetalum chinense Plant.

Ph.D. Thesis
In
Agric. Sci. (Ornamental Horticulture)

By

AHMED FARGHALY HASSANEAN

B.Sc. Agric. Sci. (Horticulture), Fac. Agric., Alazhar Univ., 2001 M.S.C. Agric. Sci. (Ornamental Horticulture), Fac. Agric., Cairo Univ., 2015

APPROVAL COMMITTEE

SUPERVISION SHEET

MICROPROPAGATION OF Loropetalum chinense Plant.

Ph.D. Thesis

In Agricultural Sci. (Ornamental Horticulture)

By

AHMED FARGHALY HASSANEAN

B.Sc. Agric. Sci.(Horticulture), Fac. Agric., Alazhar Univ., 2001 M.S.c. Agric. Sci. (Ornamental Horticulture), Fac. Agric., Cairo Univ., 2015

SUPERVISION COMMITTEE

Dr. HAMDY MOHAMMED ALI EL-BAGOURY(Late)

Professor of Ornamental Horticulture, Fac. Agric., Cairo University

Dr. WEAAM RAMADAN ARAFA SAKR

professor of Ornamental Horticulture, Medicinal and Aromatic Plants Fac. Agric., Cairo University.

Dr. MAMDOUH AHMED EL SHAMY

Head Researcher of Botanical Gardens Research, Agriculture Research Center

Name of Candidate: Ahmed Farghaly Hassanean Degree: Ph.D.

Title of Thesis: Micropropagation of *Loropetalum chinense* Plant.

Supervisors: Dr. Hamdi Mohammed Ali El-Bagoury (Late)

Dr. Weaam Ramadan Arafa Sakr

Dr. Mamdouh Ahmed Ebrahim EL Shamy

Department: Ornamental Horticulture

Date: / /

ABSTRACT

This study was carried out in the Plant Tissue Culture Laboratory, Horticulture Research Institute, Agriculture Research Center, Ministry of Agriculture and land reclamation, Giza, Egypt, during the period from 2016 to 2019. An efficient direct shoot regeneration micropropagation protocol has been established successfully for commercial in vitro propagation of Loropetalum chinense, an ornamental plant with medicinal values. The optimized sterilization conditions for the Terminal node stem cuttings with its two axillary buds explants were exposure to sodium hypochlorite at 0.25 % for 20 min and 0.5 % for 15 or 20 min gave the highest value of survived explants (100 %). In establishment stage, MS medium supplemented with 0.5 mg/l IBA gave (3.0 shoots) the highest number of shoots. The multiplication stage should be cultured on MS medium supplemented with 2.0 mg/l BA which recorded 7.67 shoots at the end of the third subculture. For rooting of the regenerated shoots, using WPM medium supplemented with NAA at 1.0 mg/l resulted in significantly highest number of roots (4.0) and highest rooting percentage (100%) than that recorded with other treatments. Growing medium consists of peatmoss and vermiculite at the ratio of 1:1 (v/v) was the best for acclimatization. Inter simple sequence repeats (ISSR) analysis detected similarity to the mother plant in the in vitro propagated plants, that the protocol will be useful for Loropetalum chinense production. This plant protocol could be used for large scale regeneration of *Loropetalum chinense*.

Key words: Loropetalum chinense, IBA, BA, ISSR analysis ,Tissue culture.

AKNOWLEDGEMENT

Tribute and praise are due, first of all, to Allah All-Merciful The Compationate GOD.

I would like to express my deepest and faithful gratitude to **Dr. Hamdi Mohammed Ali El-Bagoury,** Professor of Ornamental HorticultUre, Faculty of Agriculture, Cairo University, for his supervision of this study

I wish to express my appreciation to **Dr. Weaam Ramadan**Arafa Sakr, Professor of Ornamental, Medicinal and Aromatic Plants, Faculty of Agriculture, Cairo University, for her supervision, valuable advice and guidance during the course of the investigation and during the preparation of this thesis.

My sincerest gratitude to **Dr. Mamdouh Ahmed El - Shamy**, Head Researcher of Botanical Garden Research, Agriculture Research Center, for his supervision, encouragement during the practical study and preparation of this thesis.

Many thanks are also due to my brothers for their unlimited help and encouragement.

LIST OF ABBREVIATIONS

AC: Activated charcoal.

BA: 6- Benzyl adenine.

BAP: Benzyl amino purine.

B5 Medium: Gamborg B5 medium.

Clorox: 5.25 % soudium hypochlorite (commercial bleach).

2,4-D: (2,4-Dichlorophenoxy acetic acid).

GA₃: Gibberellic acid, (Gibberellin A₃).

IAA: Indole-3- acetic acid.

IBA: Indole -3- butyric acid.

Kin: 6-Furfuryl- amino purine (Kinetin).

MS: Murashige& Skoog medium.

NAA: Naphthalene acetic acid.

PVP: Polyvinyl pyrrolidone.

TDZ: Thidiazuron.

Tween 20: Polyoxyethlene sorbitan monolaurate.

V/V: Volume by volume.

WPM: Woody plants medium.

Gl: Glutamine

AdS: Adenine sulphate.

PGR: Plant growth regulators

PPFD: Photosynthetic photon flux density

PG: Phloroglucinol

MSBs: Multiple shoot buds

TIS: Temporary immersion system

DNA: Deoxyribonucleic acid

ISSR: Inter simple sequence repeats

ISSR-PCR: Inter simple sequence repeat - Polymerase chain reaction

SSRs: Simple sequence repeats

PCR: Polymerase chain reaction

McCown's: McCown's woody plant medium.

CK: Cytokinin

RAPD: random amplified polymorphic DNA

UPGMA: unweighted pair group method with arithmetic means

(DAMD): directed amplification of minisatellite DNA

LIST OF TABLES

| No | Title | Page |
|----|--|------|
| 1 | List of used ISSR primers name and their nucleotide sequences. | 52 |
| 2 | Effect of different concentrations of silver nitrate (AgNO3) and soaking period on surface sterilization of <i>Loropetalum chinense</i> | 55 |
| 3 | Effect of different concentrations of Sodium hypochlorite (NaOCl) and soaking period on surface sterilization of <i>Loropetalum chinense</i> | 58 |
| 4 | Effect of different concentrations of mercuric chloride (HgCl ₂) and soaking periods on surface sterilization of <i>Loropetalum chinense</i> . | 60 |
| 5 | Effect of different concentrations of NAA, IBA and 2,4 – D and different media on shoot length during establishment stage of <i>Loropetalum chinense</i> | 62 |
| 6 | Effect of different concentrations of NAA, IBA and 2,4 – D and different media on number of shoots during establishment stage of <i>Loropetalum chinense</i> . | 64 |
| 7 | Effect of different concentrations of NAA, IBA and 2,4 -D and different media on number of leaves during establishment stage of <i>Loropetalum chinense</i> | 65 |
| 8 | Effect of MS medium and different concentrations of BA, Kin and 2iP on shoot length during multiplication stage of <i>Loropetalum chinense</i> | 67 |

| 9 | Effect of MS medium and different concentrations of BA, Kin and 2iP on number of shoots during multiplication stage of <i>Loropetalum chinense</i> | 69 |
|----|--|----|
| 10 | Effect of MS medium and different concentrations of BA, Kin and 2iP on number of leaves during multiplication stage of <i>Loropetalum chinense</i> | 70 |
| 11 | Effect of WPM medium and different concentrations of BA, Kin and 2iP on shoot length during multiplication stage of <i>Loropetalum chinense</i> | 72 |
| 12 | Effect of WPM medium and different concentrations of BA, Kin and 2iP on number of shoots during multiplication stage of <i>Loropetalum chinense</i> | 74 |
| 13 | Effect of WPM medium and different concentrations of BA, Kin and 2iP on number of leaves during multiplication stage of <i>Loropetalum chinense</i> | 75 |
| 14 | Effect of MS,WPM medium supplemented with different concentrations of NAA, IBA on plantlet length during rooting stage of <i>Loropetalum chinense</i> | 77 |
| 15 | Effect of MS,WPM medium supplemented with different concentrations of NAA, IBA on number of leaves during rooting stage of <i>Loropetalum chinense</i> | 80 |
| 16 | Effect of MS,WPM medium supplemented with different concentrations of NAA, IBA on number of roots during rooting stage of <i>Loropetalum chinense</i> | 82 |

| 17 | Effect of MS,WPM medium supplemented with different concentrations of NAA, IBA on root length during rooting stage of <i>Loropetalum chinense</i> | 85 |
|----|---|----|
| 18 | Effect of MS,WPM medium supplemented with different concentrations of NAA, IBA on rooting percentage during rooting stage of <i>Loropetalum chinense</i> | 87 |
| 19 | Effect of different growing media on plantlet length, number of leaves/ plantlet, number of shoots/plantlet and survival percentage during acclimatization stage of <i>Loropetalum chinense</i> | 88 |
| 20 | Type and number of amplified DNA bands generated by five ISSR primers used for the identification of the <i>Loropetalum chinense</i> | 92 |

LIST OF FIGURES

| No | Title | Page |
|----|--|------|
| 1 | Loropetalum chinense | 3 |
| 2 | Effect of IBA at 0.0, 0.5, 1.0, 2.0, 4.0 mg/l in MS medium during establishment stage of <i>Loropetalum</i> | |
| | chinense | 62 |
| 3 | Effect of BA at (0.0, 1.0, 2.0,3.0, 4.0, 5.0 mg/l) in MS medium during multiplication stage of <i>Loropetalum</i> | |
| | chinense | 67 |
| 4 | Effect of BA at (0.0, 1.0, 2.0,3.0, 4.0, 5.0 mg/l) in WPM medium during multiplication stage of <i>Loropetalum</i> | |
| | chinense | 72 |
| 5 | Effect of NAA at (0.0, 1.0, 2.0, 4.0mg/l) in WPM medium during rooting stage of <i>Loropetalum</i> | |
| | chinense | 78 |
| 6 | Effect of growing media on acclimatization stage of | 90 |
| | Loropetalum chinense | 89 |
| 7 | Gel electrophoresis of ISSR fragments profiles generated by using 5 primers: 844-A, 844-B, 17898 A, 17898-B and | |
| | HB-14) | 93 |

CONTENTS

| | Pag |
|--|-----|
| INTRODUCTION | 1 |
| REVIEW OF LITERATURE | 4 |
| 1. Surface sterilization | 4 |
| 2. Establishment stage | 9 |
| 3.Multiplication stage | 14 |
| 4. Rooting stage | 27 |
| 5. Acclimization stage | 39 |
| 6. Molecular genetic identification | 44 |
| MATERIALS AND METHODS | 47 |
| RESULTS | 53 |
| a. Surface sterilization with Silver nitrate | 53 |
| 1. Contamination percentage | 53 |
| Survival percentage | 53 |
| 3. Mortality percentage | 54 |
| b. Surface sterilization Sodium Hypochlorite | 55 |
| 1. Contamination percentage | 55 |
| 2. Survival percentage | 56 |
| 3 Mortality percentage | 57 |
| c. Surface sterilization with mercuric chloride | 58 |
| 1. Contamination percentage | 58 |
| 2. Survival percentage | 59 |
| 3. Mortality percentage | 59 |
| 2. Establishment stage | 61 |
| a. Shoot length (cm) | 61 |
| b. Number of shoots | 63 |
| c. Number of leaves | 64 |
| 3. Multiplication Stage | 65 |
| a. Effect of MS medium supplemented with different | |
| concentrations of BA, Kin and 2iP | 65 |
| , | UJ |
| (1) Number of shoots | 65 |
| (2) b. Shoot length | 68 |