

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

بسم الله الرحمن الرحيم





HANAA ALY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكرونيله



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



HANAA ALY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HANAA ALY





Salivary level of interleukin-17 in patients having atrophic and erosive oral lichen planus before and after treatment with topical steroids

(Controlled clinical trial)

Thesis Submitted to Oral Medicine, Periodontology, Oral Diagnosis and Radiology Department, Faculty of Dentistry,
Ain Shams University

In Partial Fulfillment of the Requirements for Masters' Degree in Oral Medicine

By

Mai Talaat Mohamed Mohamed

B.D.S 2014

Faculty of Dentistry, Future University in Egypt

Supervisors

Ass. Prof. Ahmed Abdel Aziz Hassan

Assistant Professor of Oral Medicine, Periodontology and Oral Diagnosis, Faculty of Dentistry, Ain Shams University

Prof. Olfat Gamil Shaker

Professor of Medical Biochemistry Faculty of Medicine, Cairo University

Dr. Radwa Ragheb Hussein

Lecturer of Oral Medicine, Periodontology and Oral Diagnosis Faculty of Dentistry, Ain Shams University

Acknowledgment

Before everything and after all things I would like to express prayerful thanks to Allah the Almighty for everything.

I would like to express my deepest appreciation as words are not enough to thank my supervisor **Dr. Ahmed Abdel Aziz**, Assistant Professor of Oral Medicine, Periodontology and Oral Diagnosis, Faculty of Dentistry, Ain Shams University, for his enthusiastic encouragement, guidance, suggestions, instructions and support – without delay- in each stage of the work.

My great thanks and appreciation go to the **Prof. Dr. Olfat Shaker**, Professor of Medical Biochemistry and Molecular Biology, Faculty of Medicine, Cairo University for her help and supervision during this study.

A special thanks to **Dr. Radwa Ragheb**, lecturer of Oral Medicine, Periodontology and Oral Diagnosis, Faculty of Dentistry, Ain Shams University, for her encouragement, advice and support in each stage of the work.

I am always thankful to my **colleagues** and **friends** for their continuous unlimited support.

Last but not least, my true love goes to all my **family**, without whom I would never been able to accomplish this work.

Dedication

To my light when all was dark,

My Mother & Father...

To my hope when there was no hope,

My beloved Husband....

Contents

Subject	Page No.
List of Abbreviations	i
List of Tables	iv
List of Figures	vi
Review of literature	1
Aim of the study	52
Subjects and Methods	53
Results	70
Case Presentation	92
Discussion	96
Summary	108
Conclusions	112
Recommendations	113
References	114
Appendix	I
Arabic Summary	

List of Abbreviations

Abbrev. Full term

AD : Atopic dermatitis

ASA : American Society of Anesthesiologists

AV : Aloe Vera

BD : Behcet disease

BM : Basement membrane

BSA : Bovine serum albumin

CCR : Chemokines cell receptors

CHS : Contact hypersenstivity

CL : Chloride

CLP : Cutaneous lichen planus

CNS : Central nervous system

COX : Cyclooxygenase

CS : Clinical score

DIF : Direct immunofluorescence

DM : Diabetes Mellitus

ELISA : Enzyme linked immune sorbent assay

EM : Electron microscope

GVHD : Graft versus host disease

HCV: Hepatitis C virus

HLA: Human leukocyte antigen

HRP : Horseradish peroxidase

HSP: Heat shock protiens

IBD : Inflammatory bowel disease

ICAM : Intercellular adhesion molecule

ICSCI : Intralesional corticosteroids injection

IFN- γ : Interferon gamma

IIF : Indirect immunofluorescence

IL : Interleukin

KCs : Keratinocytes

KP : Koebner phenomenon

KT: Potassium

LCs : Langerhans cells

LOX : Lipooxygenase

LP : Lichen planus

LR : Lichenoid reaction

MB-PDT: Methylene blue-Photo dynamic Therapy

MHC : Major histocompatability

MMP : Matrix Metalo-protinase

MS : Multiple sclerosis

Na : Sodium

NOS : Nitric oxide synthase

OLDR : Oral licheniod drug reaction

OLL : Oral lichenoid lesion

OLP : Oral lichen planus

OLR : Oral lichenoid reaction

OSCC : Oral squamous cell carcinoma

PBMCs: Peripheral blood mononuclear cells

PBS : Phosphate buffered saline

PDT: Photo dynamic therapy

PMDs : Potentially malignant disorder

PUVA : Psoralen and Ultra violet A

PV : Pemphigus vulgaris

PVAS : Pain visual analogue scale

RA : Rheumatoid arthritis

RANTES: Regulated upon activation normal T-cell

expressed and secreted

RORyt : Retinoic acid receptor related orphan nuclear receptor

SAA : Salivary alpha amylase

SD : Standard deviation

SLE : Systemic lupus erythematosis

TGF-B1: Transforming growth factor B1

TH: T Helper

TMB: Tetra-methyl-benzidine

TNF-\alpha: Tumor necrosis factor alpha

UV : Ultra violet

UWS : Unstimulates whole saliva

VAS : Visual analogue scale

VCAM : Vascular cell adhesion molecule

VHT : Vascular hypertension

WHO: World health organization

List of Tables

Table No.	Title	Page No.
Table (1):	Summary of statistics of demogradata	_
Table (2):	Mean and standard deviation values for clinical score (CS different groups) in
Table (3):	Mean and standard deviation (SD) v for clinical score (CS) in diff follow-up intervals	ferent
Table (4):	Mean and standard deviation (SD) v for (CS) percentage change in diff follow-up intervals	ferent
Table (5):	Mean and standard deviation values for pain visual analogue (P-VAS) in different groups	scale
Table (6):	Mean and standard deviation values for pain visual analogue (P-VAS) in different follo intervals	scale w-up
Table (7):	Mean and standard deviation values for pain visual analogue (P-VAS) percentage change in diff follow-up intervals	scale
Table (8):	The mean and standard deviation values for IL-17 salivary concentre (pg/ml) in the three groups	ration

Table (9):	Mean and standard deviation (SD) values for IL-17 salivary concentration (pg/ml) in group (I) before and after treatment	88
Table (10):	Mean and standard deviation (SD) values for IL-17 salivary concentration percentage change in group (I) before and after treatment	89
Table (11):	Correlation between clinical score and IL-17 salivary concentration	90
Table (12):	Correlation between pain visual analogue scale and IL-17 salivary concentration in group I	91

List of Figures

Figure No.	Title	Page No.
Figure (1):	Pain Visual Analog Scale (P-V score sheet.	
Figure (2):	Coated micro well	63
Figure (3):	First incubation	63
Figure (4):	Second incubation	63
Figure (5):	Third incubation	63
Figure (6):	Reacted substrate	63
Figure (7):	Bar chart showing gender distrib in the three study groups	
Figure (8):	Bar chart showing mean and star deviation of the age (years) of participants in the three study group	f the
Figure (9):	Bar chart showing mean and star deviation values of clinical score in the three study groups	(CS)
Figure (10):	Line chart showing mean and star deviation values of clinical score in group (I) before and after treatme	(CS)
Figure (11):	Bar chart showing mean and star deviation values for pain vanalogue scale (P-VAS) in the study groups	risual three
Figure (12):	Line chart showing mean and star deviation values for pain values analogue scale in group (I) (P-VA	isual

Figure (13):	Bar chart showing the mean and standard deviation values of IL-17 salivary concentration (pg/ml) in the three groups
Figure (14):	Line chart showing mean and standard deviation values of IL-17 salivary concentration (pg/ml) in group (I) before and after treatment
Figure (15):	Scatter plot showing the correlation between clinical score and IL-17 salivary concentration in group I90
Figure (16):	Scatter plot showing the correlation between pain visual analogue scale (P-VAS) score and IL-17 salivary concentration