

بسم الله الرحمن الرحيم





HOSSAM MAGHRABY





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



HOSSAM MAGHRABY



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغيار



HOSSAM MAGHRABY



Value of Adding a T2 Mapping Sequence to a Routine MR Imaging Protocol in Evaluation of Articular Cartilage of the Knee Joint

Thesis

Submitted for Partial Fulfillment of M.D Degree in **Radiodiagnosis**

Presented by

Samar Shehata Mohamed El Sayed Hasan Noreldine M.B., B. Ch

M.Sc of Radiodiagnosis
Faculty of Medicine- Ain Shams University

Supervised by

Prof. Dr. Mohamed Amin Nassef

Professor of Radiodiagnosis
Faculty of Medicine - Ain Shams University

Dr. Mennatallah Hatem Shalaby

Associate Professor of Radiodiagnosis Faculty of Medicine - Ain Shams University

Dr. Ali Haggag Ali

Lecturer of Radiodiagnosis
Faculty of Medicine - Ain Shams University

Faculty of Medicine - Ain Shams University 2021



سورة البقرة الآية: ٣٢

Acknowledgments

First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.

I am extremely grateful to Prof. Dr. Mohamed Amin Nassef, Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, without his patience and understanding; this work would not have been possible.

I am deeply indebted to Dr. Mennatallah Hatem Shalaby, Associate Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for her valuable suggestions, guidance, advice and kind instructions through this work.

I would also like to show my deepest gratitude to Dr. Ali Haggag Ali, Lecturer of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for her remarkable ideas and remarks suggestions, guidance, advice and kind instructions through this work.

I would like to express my hearty thanks to all my family for their support till this work was completed.

Samar Shehata Noreldine

List of Contents

Title	Page No.
List of Tables	i
List of Figures	ii
List of Abbreviations	ix
Introduction	1
Aim of the Work	5
Review of Literature	
Normal Articular Cartilage	6
• Morphological MR Imaging of Articular Cartilage	19
• Compositional MR Imaging of Articular Cartilage	36
• Cartilage Degeneration and Osteoarthritis	55
• T2 Mapping Evaluation of Cartilage Degeneration	70
Patients and Methods	78
Results	86
Illustrative Cases	99
Discussion	122
Summary and Conclusion	128
References	130
Arabic Summary	

List of Tables

Table No	. Title Page No.	<u>. </u>
Table 1:	Comparison of different sequences of compositional imaging of cartilage	54
Table 2:	Classification of Osteoarthritis	57
Table 3:	ICRS grading system	63
Table 4:	Modified outerbridge classification	65
Table 5:	MRI sequences used in the study	82
Table 6:	Age distribution of the patients participating in the study.	86
Table 7:	Distribution of cases according to gender	87
Table 8:	Distribution of cases according to Clinical Presentation	88
Table 9:	Distribution of patients according to pain site	89
Table 10:	Conventional MRI findings	91
Table 11:	Conventional MRI diagnostic indices	91
Table 12:	Distribution of patellar degeneration according to ICRS degeneration grades	93
Table 13:	Distribution of trochlear degeneration according to ICRS degeneration grades	94
Table 14:	Distribution of medial femoral degeneration according to ICRS degeneration grades	95
Table 15:	Distribution of medial tibial degeneration according to ICRS degeneration grades	96
Table 16:	Distribution of lateral femoral degeneration according to ICRS degeneration grades	97
Table 17:	Distribution of lateral tibial degeneration according to ICRS degeneration grades	98

List of Figures

Fig. No.	Title	Page No.
Figure 1:	Schematic representation of articular composition and morphology in hea early-stage osteoarthritic (B), and a stage osteoarthritic cartilage (C)	lthy (A) , dvanced-
Figure 2:	(A) Negative ions attract positive ions and water molecules. (B) This of the proteoglycans is constrained surrounding collagen meshwork	counter swelling l by the
Figure 3:	Schematic, cross-sectional diagrament diagrament sectional diagrament sectional diagrament diagrame	ram of
Figure 4:	(A) PD FS MR image in 32 year with arthroscopically normal retroarticular cartilage. (B) A graphic deplayers of articular cartilage on a MR as identified on histology	old man opatellar oiction of RI image
Figure 5:	Axial FS T2W images of the targetted for cartilage imaging (patella A) large
Figure 6:	FOV (B) Small FOV Coronal T1W image of knee joint	
Figure 7:	Shows axial 2D T2-weighted fast S provides excellent depiction of a foca of deep cartilaginous delamination with high contrast between the	E image al region (arrows),
	surfaces and synovial fluid	
Figure 8:	Sagittal PD-weighted non-FS (A) an images of knee reveal efficacy of	
Figure 9:	sequence to depict articular cartilage (a) Coronal 3D intermediate-weight SE source image. Axial (b) and say images obtained with reformatting coronal source image dataset, Corintermediate-weighted fast SE source shows no cartilage abnormalities	ated fast gittal (c) g of the conal 3D ce image
	shows no carmage abhormannes	

Fig. No.	Title Page	No.
Figure 10:	Coronal 3D SPGR image with lipid suppression shows intact high-signal-intensity cartilaginous surfaces in the medial and lateral tibiofemoral compartments	
Figure 11:	Sagittal 3D SPGR of knee provides excellent contrast between cartilage and subchondral bone along signal-intensity cartilaginous surfaces in tibiofemoral compartments	
Figure 12:	(a) T2-weighted fat-suppressed image shows large areas of complete loss of cartilage in the medial tibiofemoral compartment with associated edematous degenerative lesions in subchondral bone marrow (arrowheads). (b) 3D FLASH image accurately depicts the areas of complete loss of cartilage in the medial compartment as seen in a, but the associated	
Figure 13:	bone marrow lesions are barely visible	
Figure 14:	Balanced steady-state free precession (bSSFP) images of knee of healthy 32-year-old male volunteer acquired using iterative decomposition of water and fat	
Figure 15:	(a) Sagittal source and (b) coronal reformatted 3D VIPR-SSFP images exhibit a high cartilage SNR and high contrast among cartilage, synovial fluid, and other joint structures	۰
Figure 16:	(A) Axial T1 image of the retropatellar cartilage obtained in a volunteer reveals morphologically normal cartilage. (B) Corresponding T2 maps for the same volunteer and (C) late unloading	

Fig. No.	Title	Page No.
Figure 17:	Bar diagram, normal knee car	_
Figure 18:	Color map of T1p relaxation tim cartilage	e in tibial
Figure 19:	A)Delayed gadolinium-enhanced resonance imaging (MRI) of (dGEMRIC), B) T2 mapping, mapping, and D) Sodium differentiate between healthy a stage knee osteoarthritis (OA)	magnetic cartilage C) T1rho imaging and early-
Figure 20:	(A) A proton density image of car T1WI after An ionic contrast a DTPA2-) was given time to pen tissue. (C)T1WI when the same was given a nonionic contrast a HPDO3A)	rtilage. (B) gent (Gd- etrate the individual agent (Gd-
Figure 21:	Sagittal ADC map obtained with weighted MR imaging in the knee.	n diffusion
Figure 22:	47 year old male volunteer with Sagittal balanced steady-staprecession (bSSFP) image of kneemm isotropic resolution and scanning time) shows superficial thickness cartilage thinning (arrow	knee pain. ate free e joint (0.3 8 minute al partial-
Figure 23:	(A) A dual-tuned 23 NA/1H coil all measurement of GAG content (dimensional cones acquisition of MRI in healthy 28-year-old female at 3 T	lows direct (B) Three- of sodium e volunteer
Figure 24:	Conceptual model for the risk osteoarthritis (OA)	factors of

Fig. No.	Title	Page No.
Figure 25:	Multiplanar FS PD-weighted images year old woman who previously un- arthroscopic partial medial meniscect	derwent
Figure 26:	Axial image acquired with multi- data image combination sequence grade 1 cartilage lesion	ple-echo shows
Figure 27:	Axial 2D fat-saturated PD-weight spin-echo image of knee joint in a 47 man shows arthroscopically co superficial cartilage fibrillation	ed fast year old onfirmed
Figure 28:	categorized as grade 2 cartilage lesion Coronal FS PD weighted image of t reveals a grade 2 chondromalacia weight-bearing surface of the medial condyle (arrows)	he knee of the femoral
Figure 29:	Axial fast spin-echo PD FS image in old man reveals a wide area of full-the chondromalacia (arrow) involving median ridge and the medial patell categorized as grade 3 lesion (fissucartilage to level of subchondral bone	39 year nickness ng the ar facet uring of
Figure 30:	(A) Axial FS PD-weighted sequence is knee joint shows grade 4 chondro (yellow arrow) of articular cartimedian patellar ridge and lateral facet. (B) Coronal FS PD weighted in an adult patient	image of omalacia lage in patellar
Figure 31:	Sagittal T2 map obtained from a volunteer subject (32 years old volunteer with grade 0 knees) show relaxation times compared to OA pate	healthy d male vs lower

Fig. No.	Title	Page No.
Figure 32:	Sagittal PD MR Image (left) shows articular cartilage (ICRS grade 0) a posterior horn of the meniscus	bove the
Figure 33:	Sagittal PD MR image (left) shows articular cartilage (ICRS grade 2) a posterior horn of the meniscus	s injured above the
Figure 34:	Patient with surgically confirmed fibrillation on the lateral tibial plate deep partial-thickness cartilage lesion	cartilage au and a
Figure 35:	Patient with surgically confirmed softening on the femoral trochlea are partial thickness cartilage lesion	cartilage ıd a deep
Figure 36:	The axial images served as a loca prescribing the coronal and sagittal	alizer for
Figure 37:	The coronal images served as a local prescribing the sagittal sections	alizer for
Figure 38:	Distribution of cases according to ge	
Figure 39:	Distribution of cases according to	
J	presentation	
Figure 40:	Distribution of patients according	to pain
	site.	
Figure 41:	Conventional MRI findings	
Figure 42:	T2 mapping Findings	
Figure 43:	Distribution of patellar dege	
Figure 44:	according to ICRS degeneration grad Distribution of trochlear dege	
rigure 44.	according to ICRS degeneration grad	
Figure 45:	Distribution of medial femoral dege	
118010 10°	according to ICRS degeneration grad	
Figure 46:	Distribution of medial tibial dege	
G : 3 -3	according to ICRS degeneration grad	

Fig. No.	Title	Page No.
Figure 47:	Distribution of lateral femoral degen	oration
riguic 41	according to ICRS degeneration grade	
Figure 48:	Distribution of lateral tibial degen	
8	according to ICRS degeneration grade	
Figure 49	Case (1)	
Figure 50	Case (1)	
Figure 51	Case (1)	100
Figure 52	Case (1)	
Figure 53	Case (1)	101
Figure 54	Case (1)	
Figure 55	Case (2)	102
Figure 56	Case (2)	
Figure 57	Case (2)	103
Figure 58	Case (2)	103
Figure 59	Case (3)	104
Figure 60	Case (3)	104
Figure 61	Case (3)	105
Figure 62	Case (3)	
Figure 63	Case (4)	
Figure 64	Case (4)	
Figure 65	Case (4)	
Figure 66	Case (4)	
Figure 67	Case (5)	
Figure 68	Case (5)	
Figure 69	Case (5)	
Figure 70	Case (5)	
Figure 71	Case (6)	
Figure 72	Case (6)	
Figure 73	Case (6)	
Figure 74	Case (6)	
Figure 75	Case (7)	
Figure 76	Case (7)	
Figure 77	Case (7)	115

Fig. No.	Title	Page No.
Figure 78	Case (7)	115
Figure 79	Case (8)	
Figure 80	Case (8)	117
Figure 81	Case (8)	118
Figure 82	Case (8)	
Figure 83	Case (9)	119
Figure 84	Case (9)	120
Figure 85	Case (9)	
Figure 86	Case (9)	121