

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY



Flow Cytometric Quantitation of Procoagulant Microparticles and Erythrocyte Phosphatidylserine Exposure in B-Thalassemia Patients Thesis

Submitted for Partial Fulfillment of M.D. Degree in Clinical and Chemical Pathology

By

Marwa Mostafa Mohammed Mohammed M.B., B.Ch. M.Sc. Ain Shams University

Supervised by

Professor / Manal Hashim Fayek

Professor of Clinical and Chemical Pathology Faculty of Medicine, Ain Shams University

Professor / Deena Mohamed Habashy

Professor of Clinical and Chemical Pathology Faculty of Medicine- Ain Shams University

Doctor / Botheina Ahmed Thabet Farweez

Assistant Professor of Clinical and Chemical Pathology Faculty of Medicine - Ain shams University

Doctor / Gehan Mostafa Hamed

Assistant Professor of Clinical and Chemical Pathology Faculty of Medicine - Ain shams University

Faculty of Medicine – Ain Shams University 2020



سورة البقرة الآية: ٣٢

Acknowledgment

Thanks first and last to **God** as we owe him for his great support and guidance in this step, and in every step in our life.

I would like to express my sincere gratitude to **Prof. Dr.**Manal Hashim Fayek, Professor of Clinical and Chemical pathology, Faculty of Medicine- Ain Shams University, for her precious supervision and everlasting support. Her valuable advice and immense knowledge were the cornerstone in the progression of this work.

I am extremely grateful to **Dr. Dina Mohamed Habashy,** Professor of Clinical and Chemical Pathology, Faculty of Medicine- Ain Shams University, for her wise suggestions, great assistance and remarkable guidance.

I would like to deeply thank **Dr. Botheina Ahmed Thabet,** Assistant Professor of Clinical and Chemical Pathology,
Faculty of Medicine- Ain Shams University, for her great efforts,
advice and patience. Her close observation, constructive remarks
and continuous encouragement have enlightened my way to finish
this work.

I am very thankful for **Dr. Gehan Mostafa Hamed,** Assistant Professor of Clinical and Chemical Pathology, Faculty of Medicine- Ain Shams University, for providing a valuable guidance and support and being very flexible and patient.

My sincere thanks go to all my colleagues who helped me through my journey to finish this research, without their support this work would have never been accomplished.

Last but not least, I would like to thank my whole family. My deepest gratitude goes to my father, who is my role model and always gives me the strength and motivation.

I would like to dedicate my thesis to my uncle who always appreciated and encouraged the research work.

List of Contents

| Title | Page No. |
|------------------------|----------|
| | |
| List of Tables | i |
| List of Figures | ii |
| List of Abbreviations | iiv |
| Introduction | 1 |
| Aim of the Study | 4 |
| Review of Literature | |
| Thalassemia | 5 |
| Microparticles | 63 |
| Subjects and Methods | 116 |
| Results | 128 |
| Discussion | 156 |
| Summary and Conclusion | |
| Recommendations | |
| References | |
| Arabic Summary | |

List of Tables

| Table No | • | Title | 2 | F | Page N | 0. |
|-------------|----------------|---------------|---------------|---------------------------------------|---|------|
| | - " | | | | | |
| Table (1): | | | | of production | | |
| Table (2): | | | | | | |
| Table (3): | Genotypes as | sociated w | ith β-thalas | semia interme | dia | 42 |
| Table (4): | Origin and an | tigens of M | icroparticle | es | · · · · · · · · · · · · · · · · · · · | 65 |
| Table (5): | Characteristic | s of differer | nt types of s | ecreted vesicle | es | 71 |
| Table (6): | Demographic | data of pa | tient and co | ontrol groups | | .140 |
| Table (7): | Clinical data | of patient g | roups | | · - ······· | 141 |
| Table (8): | Laboratory do | ıta of patieı | nt groups | | | .142 |
| Table (9): | Microparticles | s assay by f | low cytome | etry | · · · · · · · · · · · · · · · · · · · | .143 |
| Table (10): | Comparison of | of demogra | phic data | between patie | ent group | |
| | and control gr | oup | | ··••· | | 144 |
| Table (11): | | | | a between the | | |
| | | | | · · · · · · · · · · · · · · · · · · · | | |
| Table (12): | Comparison of | of the patier | nt subgroup | s clinical data | | .145 |
| Table (13): | Comparison of | of the patier | nt subgroup | s laboratory d | ata | .146 |
| Table (14): | | | | y between po | | |
| | control groups | S | ····· | | ····· | .147 |
| Table (15): | | | | ry between th | | |
| | | | - | roups | | |
| Table (16): | | | | roparticles (/μ | | |
| Table (17): | | | | Microparticle | | |
| | | | | equency | | |
| Table (18): | | | | Microparticle | | |
| | | | | eters | | |
| Table (19): | | | | Microparticle | - | |
| | parameters a | nd Fe chela | tion therap | y | • | 152 |

List of Figures

| Fig. No. | Title | Page No. |
|--------------|---|-----------------|
| Figure (1): | Schematic diagram of the basic structure haemoglobin molecule | • |
| Figure (2): | ${ m O_2	ext{-}Hb}$ dissociation curve in a theortically hea | althy subject 9 |
| Figure (3): | α-and β-globin gene clusters and Hl during development | • |
| Figure (4): | Pathophysiology of β-thalassemia | 22 |
| Figure (5): | Cause and sequelae of ineffective erythropoi | esis25 |
| Figure (6): | Factors contributing to the hypercoagulab | |
| Figure (7): | β-Thalassemia major | 45 |
| Figure (8): | β-Thalassemia minor | 49 |
| Figure (9): | HbA ₂ determination by HPLC | 50 |
| Figure (10): | HbH disease | 60 |
| Figure (11): | Model of formation of microparticles | 67 |
| Figure (12): | Release of microvesicles and exosomes | 69 |
| Figure (13): | Formation of microparticles and apoptotic k | oodies70 |
| Figure (14): | Multifaceted role of MPs in coagulation pro | cesses77 |
| Figure (15): | Mixed MPs in SCD children act at the interhemolysis and clotting activation | |
| Figure (16): | Schematic representation of a flow cytomet | er97 |
| Figure (17): | Pre-analytics as a major source of vari | |
| Figure (18): | Low detection threshold for standard flow c | sytometers 101 |
| Figure (19): | Technical approaches for MP investigation | 111 |
| Figure (20): | Characterization and gating of MPs | 123 |

List of Figures (Cont...)

| Fig. No. | Title | Page No. |
|--------------|--|----------------------|
| Figure (21): | Characterization and gating of MPs control tube | |
| Figure (22): | Characterization and gating of MPs in groups | |
| Figure (23): | Characterization and gating of intact R annexin V | |
| Figure (24): | ROC curve showing diagnostic performan | nce of total MPs 149 |
| Figure (25): | Diagrammatic representation of total Mic among control and patient groups | |
| Figure (26): | Diagrammatic representation of ErMPs, Parents PS ⁺ PMPs, PS ⁺ RBCs among control and expressed as percentage | patient groups |
| Figure (27): | Diagrammatic representation of ErMPs, Pa PS ⁺ PMPs among control and patient gra as absolute number/ µ1 | oups expressed |
| Figure (28): | Diagrammatic representation of ErMPs, Pa PS ⁺ PMPs, PS ⁺ RBCs among thalassen expressed as percentage | nic subgroups, |
| Figure (29): | Diagrammatic representation of ErMPs, Pa PS ⁺ PMPs among the thalassemic subgra as an absolute number/ μ1 | oups, expressed |

List of Abbreviations

| Abb. | Full term |
|--------------------------|--|
| μl: | Microliter |
| | Micro-nuclear magnetic resonance |
| • | 2, 3- biphosphoglycerate |
| | Acute coronary syndrome |
| | Atomic force microscopy |
| | Alanine transferase enzyme |
| | Antiphospholipid syndrome |
| | Aniiphospholipia synaloine Anomalous small-angle X-ray scattering |
| | Anomaious sinair angle x ray scarrening Aspartate aminotransferase enzyme |
| | Aspanaie animonansierase enzyme Area under curve |
| | |
| | Below costal margin |
| | Bone marrow transplantation |
| | Catabolite activator protein |
| | Complete blood count |
| | Clathrin-coated vesicles |
| | Cluster of differentiation |
| CD40L: | • |
| CEACAM: | Carcinoembryonic antigen related cell adhesion |
| -11- | molecule |
| | Cyclic guanosine monophosphate |
| | Coronary heart disease |
| <i>Cm:</i> | |
| | Central nervous system |
| <i>CO</i> ₂ : | Carbon dioxide |
| COPD: | Chronic obstructive pulmonary disease |
| Cryo EM: | Cryo electron microscopy |
| <i>DIC:</i> | Disseminated intravascular coagulation |

| Abb. | Full term |
|-------------------|---|
| dl: | . Deciliter |
| DLS: | . Dynamic light scattering |
| DPG: | . Diphosphoglycerate |
| ELAM-1: | . Endothelial leukocyte adhesion molecule-1 |
| <i>ELISA:</i> | . Enzyme-linked immunosorbent assay |
| <i>EMH:</i> | . Extramedullary hematopoeisis |
| <i>EMPs:</i> | . Endothelial MPs |
| <i>EPO:</i> | . Erythropoietin |
| <i>ER:</i> | . Endoplasmic reticulum |
| ERFE: | . Erythroferrone |
| <i>ErMPs:</i> | . Erythrocyte MPs |
| F VII: | . Factor VII |
| F VIIa: | . Active factor VII |
| <i>FCM:</i> | . Flow cytometry |
| Fe ^{2+:} | . Iron in the ferrous state |
| <i>FFP:</i> | . Fresh frozen plasma |
| <i>FITC:</i> | . Fluorescein isothiocyanate |
| fl: | . Femtoliter |
| <i>FSC:</i> | . Forward scatter |
| GC: | . Guanylate cyclase |
| GLA: | .γ-carboxyglutamic acid |
| gm: | . Gram |
| <i>GP:</i> | . Glycoprotein |
| GTP: | . Guanosine triphosphate |
| <i>GWAS:</i> | . Genome - wide association studies |
| HAs: | . Hemolytic anemias |
| <i>Hb:</i> | . Hemoglobin |

| Abb. | Full term |
|---------------------|---|
| HbBart's: | . Hemoglobin Bart's |
| HbF: | . Fetal Hb |
| HbH: | . Hemoglobin H |
| HbS: | . Hemoglobin S |
| HCO3 ⁻ : | . Bicarbonate ions |
| HCV: | . Hepatitis C virus |
| hHPFH: | . Heterocellular HPFH |
| HIT: | . Heparin-induced thrombocytopenia |
| HPFH: | . Hereditary persistence of fetal Hb |
| HS: | . Highly significant |
| hs-FCM: | . High-sensitivity flow cytometer |
| ICAM: | . Intercellular adhesion molecule |
| ICAM-1: | . Intercellular adhesion molecule-1 |
| ID1: | . Inhibitor of DNA binding 1 protein |
| IDA: | . Iron-deficiency anemia |
| <i>IE:</i> | . Ineffective erythropoiesis |
| /L: | . Interleukin |
| IQR: | . Inter-quartile range |
| <i>ISTH:</i> | . International Society on Thrombosis and |
| | Haemostasis |
| ITP: | . Immune thrombocytopenia |
| KCC: | .K-Cl cotransport |
| Kg: | . Kilogram |
| <i>LCR:</i> | . Locus control region |
| LDH: | . Lactate dehydrogenase enzyme |
| <i>LMP:</i> | . Leukocyte microparticles |
| MCH: | . Mean corpuscular Hb |

| Abb. | Full term |
|-------------------------|---|
| MCV: | . Mean corpuscular volume |
| MelCAM: | . Melanoma cell adhesion molecule |
| mg: | . Milligram |
| miRNA: | . Micro RNA |
| MPN: | . Myeloproliferative neoplasms |
| MPs: | . Microparticles |
| mRNA: | . Messenger RNA |
| MVEs: | . Multi-vesicular endosomes |
| NADPH oxidase: | . Nicotinamide adenine dinucleotide phosphate oxidase |
| NMD: | . Nonsense-mediated decay |
| <i>NO:</i> | . Nitric oxide |
| No.: | . Number |
| <i>NPV:</i> | . Negative predictive value |
| <i>NS:</i> | . Non significant |
| NTA: | . Nanoparticle tracking analysis |
| NTDT: | . Non-transfusion dependent thalassaemia |
| <i>O</i> ₂ : | . Oxygen |
| ODC: | . O_2 -Hb dissociation curve |
| PBS: | . Phosphate Buffer Saline |
| PC: | . Phosphatidylcholine |
| PCO ₂ : | . CO ₂ partial pressure |
| PEA: | . Phosphatidylethanolamine |
| <i>PE:</i> | . Phycoerythrin |
| PECAM: | . Platelet endothelial cell adhesion molecule |
| PF4: | . Platelet factor 4 |
| <i>PFP:</i> | . Platelet free plasma |

| Abb. | Full term |
|------------------------|--|
| pg: | . Picogram |
| pJak2: | . Phosphorylated Jak2 |
| PLTs: | . Platelets |
| PMPGA: | . PMP generation assay |
| PMPs: | . Platelet MPs |
| PNH: | . Paroxysmal nocturnal hemoglobinuria |
| <i>PO₂:</i> | . O_2 partial pressure |
| <i>PPP:</i> | . Platelet poor plasma |
| <i>PPV:</i> | . Positive predictive value |
| <i>PS:</i> | . Phosphatidylserine |
| PTC's: | . Premature termination codons |
| QTL: | . Quantitative trait loci |
| <i>RA:</i> | . Rheumatoid arthritis |
| RANTES: | . Regulated on activation, normal T cells expressed and secreted |
| <i>RBC:</i> | . Red blood cell |
| <i>RDW:</i> | . Red cell distribution width |
| <i>RES:</i> | . Reticuloendothelial system |
| <i>ROC:</i> | . Receiver operating characteristic curve |
| ROS: | . Reactive oxygen species |
| <i>RPS:</i> | . Resistive pulse sensing |
| S. ferritin: | . Serum ferritin |
| S: | . Significant |
| <i>SAXS:</i> | . Small-angle X-ray scattering |
| SCA: | . Sickle cell anemia |
| SCD: | . Sickle cell disease |
| <i>SCF:</i> | . Stem cell factor |