



شبكة المعلومات الجامعية  
التوثيق الإلكتروني والميكروفيلم

# بسم الله الرحمن الرحيم



**HANAA ALY**



شبكة المعلومات الجامعية  
التوثيق الإلكتروني والميكروفيلم



# شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلم



**HANAA ALY**



شبكة المعلومات الجامعية  
التوثيق الإلكتروني والميكروفيلم

# جامعة عين شمس التوثيق الإلكتروني والميكروفيلم

## قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها  
علي هذه الأقراص المدمجة قد أعدت دون أية تغييرات



## يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



**HANAA ALY**

**Ain Shams University**  
**Faculty of Science**  
**Entomology Department**



**Biodiversity and distributional Ecology of ticks  
(Acari: Ixodidae) infesting domestic animals  
in Egypt**

**A Thesis**

**Submitted for the degree of Doctor of Philosophy in Science  
"Entomology Department"**

**By**

**Mohamed Okely Bayoumi**  
(M.Sc. Entomology 2018)

**Under the Supervision of**

**Rabia Abdel Wahab Enan**  
Professor of Entomology, Entomology Department, Faculty of  
Science, Ain Shams University

**Sohair Mohammad Mahmoud GadAllah**  
Professor of Entomology, Entomology Department, Faculty of  
Science, Ain Shams University

**Abdallah M. Samy**  
Lecturer of Entomology, Entomology Department, Faculty of  
Science, Ain Shams University

**(2021)**

## Biography

**Name:** Mohammed Okely Bayoumi

**Degree Award:** M.Sc. (Entomology)

**Department:** Entomology

**Faculty:** Science

**University:** Ain Shams University

**Date of Graduation:** 2015

**Date of Appointment:** June 2016

**Occupation:** Assistant lecturer, Department  
of  
Science,  
Entomology, Faculty of  
Ain-shams University

**Date of Registration  
for Ph.D. degree:** 12/11/2018

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَمَا مِنْ دَابَّةٍ فِي الْأَرْضِ  
إِلَّا عَلَى اللَّهِ رِزْقُهَا وَيَعْلَمُ  
مُسْتَقَرَّهَا وَمُسْتَوْدَعَهَا كُلٌّ  
فِي كِتَابٍ مُبِينٍ

سورة هود آية: ٦

## *ACKNOWLEDGEMENTS*

*“I wish to express my deep thanks to ALLAH who fulfilled my hopes, offered every possible aid for any one in need to it”.*

*I wish to express my greatest gratitude, deepest thanks to Prof. Dr. Rabia Abdel Wahab Enan, Professor of Entomology, Faculty of Science, Ain Shams University for suggesting the research work, kind supervision, her faithful encouragement, valuable advice and guidance during the progress of this study until the preparation and writing of this manuscript.*

*I am particularly grateful to Prof. Dr. Sohair Mohammad Mahmoud GadAllah, Professor of Entomology, Faculty of Science, Ain Shams University for giving me the chance to be one of her students, her faithful encouragement, her generous advice and valuable discussions during the progress of this study.*

*I am deeply indebted to Dr. Abdallah Samy, Lecturer of Entomology, Entomology Department, Faculty of Science, Ain Shams University, for his supervision, valuable advice and guidance during the progress of this study.*

*My thanks are also passed to the staff members and colleagues of Entomology Department for the various help they offered me throughout this*

*study.*

*I am grateful for all the great friends I have. I hope they knew how much I appreciate them.*

*Sincere thanks are also to my family who suffered a lot during the preparation of this work. Finally, I am indebted forever to my mother for her help, support and continuous encouragement.*

## DEDICATION

*This thesis is dedicated to my dear family; especially my father's spirit and my mother, for love, constant prayers, encouragement, and continuous help throughout my whole life. Also, to uncle's and grandfather's spirits for continuous help, support, and encouragement.*

*To my beloved sisters, Salma Okely and Norhan Okely for their help and encouragement.*

# Contents

Chapter I. Introduction and Literature review .....	1
I.1. Introduction .....	1
I.2. Aim of the study.....	4
I.3. Literature review.....	6
I.3.1. Hard ticks of Egypt.....	7
I.3.1.1. Genus <i>Amblyomma</i> Koch, 1844.....	8
I.3.1.2. Genus <i>Haemaphysalis</i> Koch, 1844.....	13
I.3.1.3. Genus <i>Hyalomma</i> Koch, 1844.....	15
I.3.1.4. Genus <i>Ixodes</i> Latreille, 1795.....	31
I.3.1.5. Genus <i>Rhipicephalus</i> Koch, 1844.....	32
I.3.2. Tick-borne pathogens (TBPs).....	45
I.3.3. Analysis and inference.....	51
I.3.4. Ecological niche modeling.....	55
Chapter II: Hard ticks (Acari: Ixodidae) infesting domestic animals in Egypt: diagnostic characters and a taxonomic key to the collected species.....	60
II.1. Introduction.....	60
II.2. Materials and Methods.....	61
II.2.1. Study area and sampling.....	61
II.2.2. Ticks identification .....	62
II.2.3. Imaging and measurements .....	62
II.2.4. Terminology and morphological abbreviations .....	63
II.2.5. Data presentation .....	64
II.2.6. Statistical analysis.....	64
II.3. Results.....	65
II.4. Discussion .....	102

Chapter III. Mapping the environmental suitability of etiological agent and tick vectors of Crimean-Congo Hemorrhagic Fever..	109
III.1. Introduction .....	109
III.2. Materials and Methods .....	112
III.2.1. Occurrence records.....	112
III.2.2. Covariate variables.....	114
III.2.3. Ecological niche modeling.....	117
III.2.4. Niche overlap of CCHF virus and its vectors .....	118
III.3. Results .....	119
III.4. Discussion .....	127
Chapter IV. Mapping the potential distribution of two tick vectors <i>Hyalomma rufipes</i> and <i>H. truncatum</i> .....	133
IV.1. Introduction .....	133
IV.2. Materials and Methods.....	135
IV.2.1. Occurrence data.....	135
IV.2.2. Environmental data .....	136
IV.2.3. Model calibration and evaluation.....	139
IV.3. Results .....	141
IV.4. Discussion .....	152
Conclusion .....	155
Recommendations.....	157
English Summary.....	158
References.....	163
Appendices.....	202
Arabic summary	

## List of Tables

<b>Table Number</b>	<b>Caption</b>	<b>Page Number</b>
<b>1.1</b>	<b>Tick-borne pathogens isolated from hard tick species that infested domestic animals in Egypt with the geographical distribution and host associations.</b>	<b>46</b>
<b>2.1</b>	<b>Hard ticks collected from different governorates in Egypt (2018-2019).</b>	<b>67</b>
<b>2.2</b>	<b>Number of hard ticks infesting diverse animals in Egypt.</b>	<b>68</b>
<b>2.3</b>	<b>Numbers of males, females, and nymphs of four different species of hard ticks.</b>	<b>68</b>
<b>3.1</b>	<b>List of original environmental variables used for the principal component analysis.</b>	<b>116</b>
<b>3.2</b>	<b>Number of occurrence records of seven tick vectors associated with CCHFV after diverse cleaning steps of the original datasets collected from different sources.</b>	<b>119</b>
<b>3.3</b>	<b>Results of Partial area under the curve (AUC) ratios summarizing evaluations of ecological niche models of CCHFV and seven tick vectors.</b>	<b>125</b>
<b>4.1</b>	<b>Summary of general circulation models (GCMs) used in the ecological niche modeling analyses of <i>Hyalomma rufipes</i> and <i>H. truncatum</i> under future climatic conditions</b>	<b>138</b>
<b>4.2</b>	<b>Partial area under the curve (AUC) ratios summarizing the results of ecological niche model evaluations of the two tick vectors <i>Hyalomma rufipes</i> and <i>H. truncatum</i>.</b>	<b>144</b>

## List of Figures

Fig Number	Caption	Page Number
1.1	Number of hard tick species infesting domestic animals in each ecological zone in Egypt.	53
1.2	Number of hard tick species collected from different domestic animals in Egypt.	53
1.3	Number of hard tick species for each genus collected from domestic animals in Egypt.	54
2.1	(a–c) Map of Egypt showing collection sites of hard tick species collected from cattle in Beni-Suef, Giza, and Sharkia governorates.	69
2.2	(a–b) Map of Egypt showing collection sites of hard tick species collected from camels in Alexandria, Aswan, and Giza governorates, Egypt.	70
2.3	Mouthparts of tick genera: <b>a</b> <i>Hyalomma</i> . <b>b</b> <i>Amblyomma</i> . <b>c</b> <i>Rhipicephalus</i> .	71
2.4	<i>Amblyomma</i> species: (a–c) <i>Amblyomma gemma</i> ; <b>a</b> dorsal view. <b>b</b> spiracle. <b>c</b> genital aperture. (d – f) <i>Amblyomma lepidum</i> ; <b>d</b> dorsal view. <b>e</b> spiracle. <b>f</b> genital aperture. (g – i) <i>Amblyomma variegatum</i> ; <b>g</b> dorsal view. <b>h</b> spiracle. <b>i</b> genital aperture.	74
2.5	<i>Hyalomma dromedarii</i> : Unfed male: <b>a</b> dorsal view. <b>b</b> posterior region of the conscutum. <b>c</b> anal shields. <b>d</b> spiracle. <b>e</b> genital aperture.	77
2.6	<i>Hyalomma dromedarii</i> : Fed male: <b>a</b> dorsal view. <b>b</b> posterior region of the conscutum. <b>c</b> anal shields. <b>d</b> spiracle. <b>e</b> genital aperture.	78
2.7	<i>Hyalomma dromedarii</i> : Female: <b>a</b> dorsal view. <b>b</b> spiracle. (c – k) morphological variations of genital aperture.	79

2.8	<i>Hyalomma excavatum</i> : Male: <b>a</b> dorsal view. <b>b</b> posterior region of the conscutum. <b>c</b> anal shields. <b>d</b> spiracle. <b>e</b> genital aperture	82
2.9	<i>Hyalomma excavatum</i> : Female: <b>a</b> dorsal view. <b>b</b> spiracle. ( <b>c – n</b> ) morphological variations of genital aperture.	83
2.10	<i>Hyalomma impeltatum</i> : Male: <b>a</b> dorsal view. <b>b</b> posterior region of the conscutum. <b>c</b> anal shields. <b>d</b> spiracle. <b>e</b> genital aperture.	86
2.11	<i>Hyalomma rufipes</i> : Male: <b>a</b> dorsal view. <b>b</b> dense setae. <b>c</b> posterior region of the conscutum. <b>d</b> anal shields. <b>e</b> spiracle. <b>f</b> genital aperture	88
2.12	<i>Hyalomma truncatum</i> : Male: <b>a</b> dorsal view. <b>b</b> posterior region of the conscutum. <b>c</b> anal shields. <b>d</b> spiracle. <b>e</b> genital aperture.	90
2.13	<i>Rhipicephalus annulatus</i> : <b>a</b> dorsal view of male. <b>b</b> anal shields. <b>c</b> spiracle. <b>d</b> dorsal view of female. <b>e</b> female genital aperture. <b>f</b> mouth parts.	92
2.14	<i>Rhipicephalus pulchellus</i> : Male: <b>a</b> dorsal view. <b>b</b> ventral view. <b>c</b> adanal plates. <b>d</b> spiracle.	94
2.15	<i>Rhipicephalus sanguineus</i> : Male: <b>a</b> dorsal view. <b>b</b> posterior region of the conscutum. <b>c</b> adanal plates. <b>d</b> spiracle.	96
2.16	<i>Rhipicephalus sanguineus</i> : Female: <b>a</b> dorsal view. <b>b</b> spiracle. <b>c</b> genital aperture. <b>d</b> engorged female with male in mating position.	97
3.1	Environmental suitability map of Crimean-Congo hemorrhagic fever in the Old World.	122
3.2	Environmental suitability of the primary potential vector of Crimean-Congo hemorrhagic fever virus, <i>Hyalomma marginatum</i> , in the Old World.	123
3.3	Environmental suitability maps of the remaining tick vectors; <i>Amblyomma variegatum</i> , <i>Dermacentor marginatus</i> , <i>Hyalomma rufipes</i> , <i>H. truncatum</i> , <i>Rhipicephalus appendiculatus</i> , and <i>R. evertsi evertsi</i> .	124

4.1	Current potential distribution of <i>Hyalomma rufipes</i> and <i>H. truncatum</i> based on present-day climatic conditions.	143
4.2	Predicted future potential distribution of <i>Hyalomma rufipes</i> under four future representative concentration pathways of climate conditions in 2050.	145
4.3	Predicted future potential distribution of <i>Hyalomma truncatum</i> under four future representative concentration pathways of climate conditions in 2050.	146
4.4	Summary of the binary modeled potential distributions of <i>Hyalomma rufipes</i> under future conditions in 2050.	148
4.5	Summary of the binary modeled potential distributions of <i>Hyalomma truncatum</i> under future conditions in 2050.	149