

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

بسم الله الرحمن الرحيم





HANAA ALY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكرونيله



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



HANAA ALY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HANAA ALY



Screening of Oral Tamoxifen Ocular Toxicity in Breast Cancer Patients

Thesis

Submitted to Faculty of Medicine, A in shams University for partial fulfillment of Master Degree in Ophthalmology

Presented by

Mai Mohamed Ahmed Ahmed
M.B., B.Ch
Faculty of Medicine, Misr University for Science and Technology
(MUST)

Under Supervision of

Prof. Dr. Abdelrahman Gaber Salman

Professor of Ophthalmology

Faculty of Medicine, Ain Shams University

Dr. Randa Hesham Ali Abdelgawad

Lecturer of Ophthalmology

Faculty of Medicine, Ain Shams University

Dr. Ahmed Said Ibrahim Abd El Motaal Lecturer of Clinical Oncology Faculty of Medicine, Ain Shams University

Dr. Ahmed Tharwat Abdelfattah

Lecturer of Ophthalmology Misr University for Science and Technology (MUST)

> F aculty of Medicine, Ain Shams University Cairo-Egypt-2021

Acknowledgments

First and foremost, I feel always indebted to ALLAH, the Most Kind and Most Merciful.

I'd like to express my respectful thanks and profound gratitude to Prof. Dr. Abdelrahman Gaber Salman, Professor of Ophthalmology, Faculty of Medicine, Ain Shams University, for his keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.

I am also delighted to express my deepest gratitude and thanks to Prof. Dr. Azza Mohamed Ahmed Said, Professor of Ophthalmology, Faculty of Medicine – Ain Shams University, for her kind care, continuous supervision, valuable instructions, constant help and great assistance throughout this work.

I am deeply thankful to Dr. Randa Hesham Ali Abdelgawad, Lecturer of Ophthalmology, Faculty of Medicine – Ain Shams University, for her keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.

I would like to thank Dr. Ahmed Said I brahim Abd El Motaal, Lecturer of Clinical Oncology, Faculty of Medicine – Ain Shams University, for his keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.

I'd like to express my respectful thanks and profound gratitude to Dr. Ahmed Tharwat Abdelfattah, Lecturer of Ophthalmology, Misr University for Science and Technology (MUST), for his keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.

I'd like to express my respectful thanks and profound gratitude to Dr. Samah Mahmoud F awzy, lecturer of Ophthalmology, Faculty of Medicine – Ain Shams University, for her keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.

Finally, would like to express my appreciation and gratitude to all my family, especially my caring and loving parents for their support till this work was completed.

Mai Mohamed Ahmed Ahmed



سورة البقرة الآية: ٣٢

Contents

Title	Page No.
List of Abbreviations	i
List of Tables	
List of F igures	
Introduction	
Aim of the Work	4
Review of Literature	
Breast cancer	5
Ocular Toxicity of Tamoxifen	16
Diagnosis of Tamoxifen Ocular Toxicity	24
Patients and Methods	40
Results	49
Discussion	88
Conclusion	93
Recommendations	95
Summary	96
References	
Arabic Summary	

L ist of A bbreviations

Abb. Full term

ASCO	American Society of Clinical Oncology	
BCVA	Best Corrected Visual Acuity	
BMI	Body mass index	
BRCA	Breast Cancer gene	
CMT	Central macular thickness	
DNA	Deoxyribonucleic acid	
EBCTCG	Early Breast Cancer Trialists' Collaborative Group	
EPIC	European prospective investigation into cancer and nutrition	
ER	Estrogen receptor	
ERG	Electroretinography	
ETDRS	Early Treatment Diabetic Retinopathy Study	
GCC	Ganglion cell complex	
HER-2	Human epidermal growth factor receptor 2	
HRT	Hormone replacement therapy	
IOP	Intraocular pressure	
ISCEV	International Society for Clinical Electrophysiology of Vision	
MAPK- ERK	Mitogen-activated protein kinases extracellular signal-regulated kinases	
MD	Mean deviation	
MFERG	multifocal electroretinogram	
MPOD	Macular pigment optical density	
NCI	Egypt National Cancer Institute	
PI3K/AKt	Phosphoinositide-3-Kinase, protein Kinase B	
PERG	Pattern electroretinogram	
PR	Progesterone receptor	
PROS	Preserved photoreceptor outer segment	

RNFL	Retinal nerve fiber layer	
RPE	Retinal pigment epithelium	
SD	Standard deviation	
SD-OCT	Spectral domain optical coherence tomography	
SERM	Selective estrogen receptor modulator	
SWS	Short wavelength sensitive	
TNM	Tumor-Node-Metastasis staging	
VEP	Visual evoked potential	

∟ ist of ⊤ables

Table	No.	Title	Page No.
1	Mean	age of studied patients	49
2		duration of tamoxifen therapy	
3		on of tamoxifen therapy divided	
4		(logMAR) and its correlation ifen therapy	
5		ary of BCVA (logMAR) differoups regarding duration of tame	
6		value of IOP value and its on of tamoxifen therapy	
7		IOP value difference betwee ing duration of tamoxifen intake	
8	Summ	ary of cup disc values and its ifen intake duration	correlation to
9	differe	ary of cup disc values and dence between two groups regard ifen intake	ing duration of
10		mottling distribution among	
11		mottling affection difference regarding duration of tamoxife	
12	Retina	ptive values of mean average l Nerve Fiber layer (RNFL) th	nickness in our
13	Fiber 1	ary of average peripapillary Layer (RNFL) values and their ifen therapy duration	correlation to
14	Descri	ptive data of average peripartiber layer (RNFL) affection	pillary Retinal
15	Summ Fiber	ary of difference peripapillary layer (RNFL) values between ing duration of tamoxifen intake i	Retinal Nerve n both groups

L ist of ⊤ables (cont..)

Table	No.	Title	Page	No.
16	fiber	nary of difference peripapillary layer (RNFL) affection between ding duration of tamoxifen intake	n both groups	61
17		riptive values of mean macular th in different quadrants		64
18		riptive data of total macular thick		
19	quadı	nary of mean macular thicknes rants and its correlation to tam ion	oxifen intake	66
20		nary of mean macular thickne		
21	betwe	nary of total macular thickn een both groups regarding kifen intake	duration of	68
22		nary of average Ganglion cell coness values		71
23		riptive data of Average Ganglion ') affection		
24		nary of average Ganglion cell cost correlation to duration of tamox	• '	
25	Comp differ	nary of mean of average Golex (GCC) complex thick ence in both groups regarding sifen intake	ness values duration of	74
26	Sumr	nary of average Ganglion Cell Collex affection difference between b	omplex (GCC)	
27		S layer thickness disruption amounts	_	76
28	-	affection among the studied patie		
29	Color	vision affection among the studie	ed patients	81

L ist of ⊤ables (cont..)

Table I	No. Title	Page No.
30	Summary of color vision affection to tamoxifen therapy	
31	Summary of total macular thickness values vision affection	
32	Summary of parameters in our study correlation to duration of tamoxifen	
33	Descriptive data of parameters used in outheir correlation to duration of tamoxifen	•

L ist of Figures

Fig. No.	. Title Page	No.
1	Distribution of estimated female breast cancer new cases, deaths, and 5-year prevalence by world region, 2012	
2	Breast cancer incidence rates/100,000 person-years by age group and year of diagnosis, 1999–2008	
3	Mechanisms of Action and Resistance in Estrogen Receptor (ER)–Targeted Therapy	
4	Fundus photography of both eyes of one patient showing crystalline deposits in perifoveal area	
5	A and B, Slit-lamp external photograph of the right eye shows subepithelial crystallin-like substance deposition in the cornea	
6	Slit-lamp examination (A) and cobalt blue filter after fluorescein instillation (B) of the patient's right eye depicting whorl-like, subepithelial deposits sparing the periphery and central cornea	
7	Ishihara color vision test plates	
8	Ishihara color vision test plates	
9	SD-OCT of the right eye. The black arrow showed the hyperreflective substances between the ganglion cell layer and inner plexiform layer correspond with	
10	SD-OCT of the left eye. A defect in the outer retinal layer with sharp edges can be seen	
11	SD-OCT of the right eye. A disruption of the photoreceptor layer can be seen.	
12	SD-OCT showed bilateral disruption of the ellipsoid zone and interdigitation zone	
13	The biphasic waveform of the ERG of a normal patient	

L ist of Figures (cont..)

Fig. No.	Title	Page	No.
14	Representative fullfield ERG waveforms tamoxifen subject according to The International Society for Clinical Electrophysiology of (ISCEV) standard protocol	national Vision	33
15	A typical standard PERG		35
16	(A) MFERG trace array derived from 103 responses of the first-order kernel. (B) dimensional plot of multifocal ERG redensity.	Three-esponse	27
17	density Visual evoked potential in a case of tantoxicity	noxifen	
18	Fransworth panel D-15 device used in our stu-		
19	Fransworth panel D-15 device used in our stu-	•	
20	Retnia scan RS 3000 advance nidek OCT devi	ce used	
21	RETI port/scan 21 Electrophysiology device our study	used in	
22	OCT optic nerve scan of one of our patients		46
23	OCT-macula map of one of our patients		46
24	OCT-of ganglion cell complex (GCC) of one patients.		47
25	Pie diagram showing groups of tamoxifen intake	therapy	
26	OCT optic nerve scan of one of our patients within normal average RNFL thickness in bot		
27	OCT optic nerve scan of one of our patients thinning of average RNFL thickness in super of right eye.	s shows ior half	
28	OCT optic nerve scan of one of our patients below normal thinning of average RNFL thick nasal quadrant of rt eye and borderline thin average RNFL thickness in nasal quadrant of le	s shows kness in ning of	

L ist of Figures (cont..)

Fig. No.	Title	Page	No.
29	OCT- macula map of one of our patients within normal average total macular thickness shows within average normal thickness in both	s, GCC	69
30	OCT macula map of one of our patients borderline thinning of the average total rethickness in nasal and lower quadrants in our in left eye	nacular ter ring	69
31	OCT-macula map of one of our patients borderline thinning of the central macular th of both eyes and borderline thinning in s quadrant of left eye and temporal quadrant of	ickness uperior of right	
22	eye in the 3 mm zone		70
32	Pie diagram showing IS/OS layer thickness the studied group (N=50)	_	76
33	OCT-macula radial of one of our patients interruption in IS/OS layer thickness in both a		77
34	interruption in IS/OS layer thickness in both e OCT-macula radial of one of our patients	-	/ /
	interruption in IS/OS layer thickness in both e		77
35	PERG of one of our patients shows delayed wave in Right eye.		79
36	MFERG of one of our patients showed reduce amplitude in ring 1		80
37	Pie diagram showing color vision distribution among the studied group (N=50).	defects	
38	Fransworth D-15 Hue test of patient in ou with normal color vision.	r study	
39	Fransworth D-15 Hue test of patient in our student	dy with	
	tritanomaly		
40	Fransworth D-15 Hue test of patient in our stupprotanomaly.		
41	Fransworth D-15 Hue test of patient in our with deutanomaly.		85