

بسم الله الرحمن الرحيم



HOSSAM MAGHRABY



شبكة المعلومات الجامعية التوثيق الالكتروني والميكرو فيلم



HOSSAM MAGHRABY

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
على هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن


تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HOSSAM MAGHRABY



بعض الوثائق الأصلية تالفة



HOSSAM MAGHRABY



بالرسالة صفحات

لم ترد بالأصل



HOSSAM MAGHRABY

**Possible teratogenic potential
of the interaction between caffeine and
certain antiasthmatic
drugs in rats**

THESIS

Presented by

MONA GEORGE YOUNAN

**Master Degree in Pharmacy
(Cairo University)**

**Assistant Lecturer of Pharmacology
Faculty of Pharmacy,
Suez Canal University**

**Submitted For
The Degree of Ph. D.
in**

**Pharmaceutical Sciences
(PHARMACOLOGY
& TOXICOLOGY)**

**FACULTY OF PHARMACY
CAIRO UNIVERSITY**

1998

UNDER THE
SUPERVISION OF

Prof. Dr.

Samir F. SAAD

Professor of Pharmacology
Faculty of Pharmacy,
Cairo University.

Prof. Dr.

M. Raouf. HAMED

Professor of Pharmacology
National Organization for Drug
Control and Research, Egypt.

ACKNOWLEDGEMENT

I dedicate this work to all those who are interested science and scientific research works , especially to my two outstanding professors: Dr. Samir F. Saad, Professor of Pharmacology, Faculty of Pharmacy, Cairo University and Dr. M. Raouf Hamed, Professor of Pharmacology, National Organization for Drug Control and Research, Ministry of Health, who gave me their kind assistance without limits.

All the gratitude, appreciation and thanks I can express to them will not be sufficient for their kind guidance, precious advice, priceless support and paternal auspices extended to me throughout the whole work.

I extend my sincere thanks to members of Pharmacology Department in NODCAR especially Professor Dr. Salwa Methwaly and Dr. Sawssan Karass. My thanks are also due to members of histopathological Department in NODCAR, especially assistant Professor A. El-Zawahry.

Thanks and indebtedness are also due to my father, mother, husband and to my daughter "Sara" for their continuous support and assistance and for giving me every convenience.

Thanks to our Almighty God who helped me to finish this work and beseech His Divinity for more fruitful and useful works.

APPROVAL SHEET

Approved

Committee in charge

(1) Prof. Dr. Samir F. Saad

.....Saad, S.F.....

(2) Prof. Dr. M. Raouf Hamed

.....M. Raouf Hamed.....


(3) Prof. Dr. Zarif Isaac

.....Zarif Isaac.....

(4) Prof. Dr. M. Adel Nasra

.....M. Adel Nasra.....

DEVELOPMENTAL PERIODS



The embryonic period ends at the end of the eighth week; by this time, the beginnings of all essential structures are present. The fetal period, extending from the ninth week until birth, is characterized by growth and elaboration of structures. Sex is clearly distinguishable by 12 weeks. The above 11- to 38-week fetuses are about half actual size.

(Reproduced from Tuchmann-Duplessis, 1975a)

CONTENTS

	Page
- List of tables	i
- List of figures	ix
- List of plates	xviii
- INTRODUCTION	
1. Congenital malformations	1
1.1 Aetiology of congenital malformations.	3
1.2 Dysmorphogenic mechanisms.	11
1.3 Methods for evaluation of drug toxicity.	15
1.4 Physico-chemical techniques.	21
2. Caffeine	
2.1 Chemical structure.	22
2.2 Physical properties.	22
2.3 Pharmacodynamic properties.	22
2.4 Pharmacokinetic properties.	25
2.5 Toxicology of caffeine.	27
2.6 Postnatal effects.	30
3. Theophylline	
3.1 Chemical structure.	32
3.2 Physical properties.	32
3.3 Pharmacodynamic properties.	32
3.4 Pharmacokinetic properties.	34
3.5 Toxicology of theophylline.	36
3.6 Postnatal effects.	38
4. Salbutamol	
4.1 Chemical structure.	39
4.2 Physical properties.	39
4.3 Pharmacodynamic properties.	39
4.4 Pharmacokinetic properties.	41
4.5 Toxicology of salbutamol.	41

- AIM OF THE WORK	43
- MATERIAL AND METHODS	44
1. Animals	44
2. Drugs and dosage preparations.	44
3. Time of administration.	44
4. Types of the performed investigations.	44
5. Experimental design.	45
- Experimental procedures:	
6. Teratology.	49
7. Behavioural testing.	51
8. Histopathological examination.	56
- RESULTS	58
- Tables and Figures	107
- Plates	258
- DISCUSSION	284
- SUMMARY	301
- REFERENCES	305
- ARABIC SUMMARY	

LIST OF TABLES

- (1) : Effect of daily maternal administration of caffeine from the 7th to the 14th day of gestation on the outcome of pregnancy in rats.
- (2) : Effect of daily maternal administration of salbutamol from the 7th to the 14th day of gestation on the outcome of pregnancy in rats.
- (3) : Effect of daily maternal administration of theophylline from the 7th to the 14th day of gestation on the outcome of pregnancy in rats.
- (4): Effect of daily maternal coadministration of caffeine (25mgKg⁻¹) and salbutamol (1 or 2 mg Kg⁻¹) from the 7th to the 14th day of gestation on the outcome of pregnancy in rats.
- (5): Effect of daily maternal coadministration of caffeine (50mgKg⁻¹) and salbutamol (1 or 2 mg Kg⁻¹) from the 7th to the 14th day of gestation on the outcome of pregnancy in rats.
- (6) : Effect of daily maternal coadministration of caffeine (25mgKg⁻¹) and theophylline (60 or 120 mg Kg⁻¹) from the 7th to the 14th day of gestation on the outcome of pregnancy in rats.
- (7) : Effect of daily maternal coadministration of caffeine (50mgKg⁻¹) and theophylline (60 or 120 mg Kg⁻¹) from the 7th to the 14th day of gestation on the outcome of pregnancy in rats.
- (8) : Effect of daily maternal administration of caffeine from the 7th to the 14th day of gestation on fetal growth parameters.
- (9) : Effect of daily maternal administration of salbutamol, from the 7th to the 14th day of gestation on fetal growth parameters.
- (10) : Effect of daily maternal administration of theophylline from, the 7th to the 14th day of gestation on fetal growth parameters.
- (11): Effect of daily maternal coadministration of caffeine (25mgKg⁻¹) and salbutamol (1 or 2 mg Kg⁻¹) from the 7th to the 14th day of gestation on fetal growth parameters.

- (12): Effect of daily maternal coadministration of caffeine (50mgKg^{-1}) and salbutamol (1 or 2 mg Kg^{-1}) from the 7th to the 14th day of gestation on fetal growth parameters.
- (13): Effect of daily maternal coadministration of caffeine (25mgKg^{-1}) and theophylline (60 or 120 mg Kg^{-1}) from the 7th to the 14th day of gestation on fetal growth parameters.
- (14): Effect of daily maternal coadministration of caffeine (50mgKg^{-1}) and theophylline (60 or 120 mg Kg^{-1}) from the 7th to the 14th day of gestation on fetal growth parameters.
- (15): Effect of daily maternal administration of caffeine from the 7th to the 14th day of gestation on the incidence of internal anomalies in rats fetuses.
- (16): Effect of daily maternal administration of salbutamol from the 7th to the 14th day of gestation on the incidence of internal anomalies in rats fetuses.
- (17): Effect of daily maternal administration of theophylline from the 7th to the 14th day of gestation on the incidence of internal anomalies in rats fetuses.
- (18): Effect of daily maternal coadministration of caffeine (25mgKg^{-1}) and salbutamol (1 or 2 mg kg^{-1}) from the 7th to the 14th day of gestation on the incidence of internal anomalies in rats fetuses.
- (19): Effect of daily maternal coadministration of caffeine (50mgKg^{-1}) and salbutamol (1 or 2 mg kg^{-1}) from the 7th to the 14th day of gestation on the incidence of internal anomalies in rats fetuses.
- (20): Effect of daily maternal coadministration of caffeine (25mgKg^{-1}) and theophylline (60 or 120 mgkg^{-1}) from the 7th to the 14th day of gestation on the incidence of internal anomalies in rats
- (21): Effect of daily maternal coadministration of caffeine (50 mgkg^{-1}) and theophylline (60 or 120 mgkg^{-1}) from the 7th to the 14th day of gestation on the incidence of internal anomalies in rats fetuses.

- (22): Effect of daily maternal administration of caffeine from the 7th to the 14th day of gestation on the incidence of skeletal anomalies in rats fetuses.
- (23): Types of induced skeletal abnormalities, following a daily maternal administration of caffeine from the 7th to the 14th day of gestation.
- (24): Effect of daily maternal administration of salbutamol from the 7th to the 14th day of gestation on the incidence of skeletal anomalies in rat fetuses.
- (25): Types of induced skeletal abnormalities, following a daily maternal administration of salbutamol from the 7th to the 14th day of gestation.
- (26): Effect of daily maternal administration of theophylline from the 7th to the 14th day of gestation on the incidence of skeletal anomalies in rat fetuses.
- (27): Types of induced skeletal abnormalities, following a daily maternal administration of theophylline from the 7th to the 14th day of gestation.
- (28): Effect of daily maternal coadministration of caffeine (25mgKg⁻¹) and salbutamol (1 or 2 mg Kg⁻¹) from the 7th to the 14th day of gestation on the incidence of skeletal anomalies in rat fetuses.
- (29): Types of induced skeletal abnormalities, following a daily maternal coadministration of caffeine (25 mg Kg⁻¹) and salbutamol (1 or 2 mg Kg⁻¹) from the 7th to the 14th day of gestation.
- (30): Effect of daily maternal coadministration of caffeine (50mgKg⁻¹) and salbutamol (1 or 2 mg Kg⁻¹) from the 7th to the 14th day of gestation on the incidence of skeletal anomalies in rat fetuses.
- (31): Types of induced skeletal abnormalities, following a daily maternal coadministration of caffeine (50 mg Kg⁻¹) and salbutamol (1 or 2 mg Kg⁻¹) from the 7th to the 14th day of gestation.