

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY



Randomized Clinical Trial to Evaluate the Efficacy of Tranexamic Acid in Irrigant Solution on Blood Loss in Transurethral Resection of the Prostate

Thesis

Submitted in partial fulfillment of the Master Degree in Urology

By

Ahmed Fawaz El-Zhary Mohamed

Bachelor of Medicine and Surgery

Under supervision of

Prof. Dr. Ahmed Mohamed Saafan

Assistant Professor of Urology Faculty of Medicine, Ain Shams University

Dr. Ahmed Tawfick Hassan

Lecturer of Urology Faculty of Medicine, Ain Shams University

Faculty of Medicine
Ain Shams University
2021



سورة البقرة الآية: ٣٢

Acknowledgment

First and foremost, I feel always indebted to AUAH, the Most Kind and Most Merciful.

I'd like to express my respectful thanks and profound gratitude to **Prof. Dr. Ahmed**Mohamed Saafan, Assistant Professor of Urology Faculty of Medicine- Ain Shams University for his keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.

I am also delighted to express my deepest gratitude and thanks to **Dr. Ahmed Tawfick****Thassan, Lecturer of Urology, Faculty of Medicine, Ain Shams University, for his kind care, continuous supervision, valuable instructions, constant help and great assistance throughout this work.

I would like to express my hearty thanks to all my family for their support till this work was completed.

Last but not least my sincere thanks and appreciation to all patients participated in this study.

Ahmed Fawaz El-Zhary Mohamed

List of Contents

Title	Page No.
List of Tables	i
List of Figures	iii
List of Abbreviations	v
Introduction	1
Aim of the Work	3
Review of Literature	
Benign Prostatic Hyperplasia (B.P.H)	4
Tranexamic Acid	48
Patients and Methods	56
Results	62
Discussion	87
Summary	98
Conclusion	101
References	102
Arabic Summary	١

List of Tables

Table No	. Title	Page No.
Table (1):	Evaluation of age, medical history history in all 50 patients	
Table (2):	Comparison between the two study groubasic demographic and clinical characte	
Table (3):	Comparison between the two study groups operative labs (CBC, Coagulation p functions, IPSS):	rofile, Renal
Table (4):	Comparison in non tranexamic acid gro the CBC values (Hemoglobin c Hematocrite, total leucocytic count, plat	oncentration,
Table (5):	Analysis of the changes in (Coagula Renal functions, IPSS) in the non tran- group at pre operative and post operative	nexamic acid
Table (6):	Comparison in tranexamic acid group of CBC values (Hemoglobin concentration, total leucocytic count, platelets), in pre of 4hrs post operative time, 24hrs p	Hematocrite, perative time,
Table (7):	Analysis of the changes in (Coagula Renal functions, IPSS) in the tranexam at pre operative and post operative	ic acid group
Table (8):	Comparison between the two study grou CBC at 4hrs post operative (HGB, HCT, T	
Table (9):	Comparison between the two study grounds CBC at 24hrs post operative, post (Coagulation profile, Renal functions, IF	st operative
Table (10):	Comparison between the two study groupre operative (Urine analysis, PSA (pelvi-abd U/S (pre and post), also DRE):	Total, Free),
Table (11):	Comparison between the two study grouthe Irrigation Volume (dL) in three immediately after surgery, T2: 4hrs po T3: 24 hrs post operative)	e times (T1: ost operative,

List of Tables (Cont...)

Table No	. Title	Page No.
Table (12):	Comparison between the two study groups Hemoglobin concentration in fluid (g/dL) times (T1: immediately after surgey, T2: operative, T3: 24 hrs post operative)	in three 4hrs post
Table (13):	Comparison between the two study groups Blood loss (ml) in three times (T1: immediatory, T2: 4hrs post operative, T3: 24 operative)	ately after hrs post
Table (14):	Comparison in the non tranexamic group (Irrigation Volume (dL), Hemoglobin con in fluid (g/dL), and Blood loss (ml) in the tl (T1: immediately after surgery, T2: operative, T3: 24 hrs post operative)	centration aree times 4hrs post
Table (15):	Comparison in the tranexamic group (Irrigation Volume (dL), Hemoglobin con (g/dL), and Blood loss (ml) in the three t immediately after surgery, T2: 4hrs post T3: 24 hrs post operative)	c in fluid imes. (T1: operative,
Table (16):	Collection of all difference values bet groups	ween two
Table (17):	Comparison between two groups (Complications, Surgery duration, Hosp (days), Quality of endoscopic view)	regarding oital Stay

List of Figures

Fig. No.	Title	Page No.
Figure (1):	Prostate Zones	7
Figure (2):	1 =Peripheral Zone, 2= Central Transitional Zone, 4= Anterior Zone	•
Figure (3):	(a) Internal anatomy of normal pro- Internal anatomy of benign prostatic hyp	
Figure (4):	Bipolar resection	25
Figure (5):	Gyrus PlasmaKinetic system Med vaporization Loop bipolar resection	
Figure (6):	Extension of the incisions at 5 and 7 o'c prostatic urethra on both sides verumontanum by Collings loop	of the
Figure (7):	Laser Vaporization of the prostate	30
Figure (8):	Treatment algorithm for BPH	33
Figure (9):	Resection of the median lobe from the bl until above the verumontanum	
Figure (10):	Final aspect of the prostatic lodge after	ΓURP39
Figure (11):	Antifibrinolytic action of tranexamic acid	d49
Figure (12):	Summarizing (pre- and 4hrs post oper 24hrs post operative time) hemoglobin con (gm/dL) in non tranexamic group	ncentration
Figure (13):	Summarizing (pre- and 4hrs post oper 24hrs post operative time) hematocrit in the non tranexamic group	percentage
Figure (14):	Summarizing (pre- and 4hrs post operative time) has concentration (gm/dL) in tranexamic gro	nemoglobin
Figure (15):	Summarizing (pre- and 4hrs post oper 24hrs post operative time) hematocrit in the tranexamic group	percentage
Figure (16):	Summarizing the statistical significant between two groups regarding (HGB) a operative	difference t 4hrs post

List of Figures (Cont...)

Fig. No.	Title	Page No.
Figure (17):	Summarizing the statistical significant between two groups regarding (HCT) at operative.	4hrs post72
Figure (18):	Summarizing the statistical significant between two groups regarding (HGB) at operative	24hrs post
Figure (19):	Summarizing the statistical significant between two groups regarding (HCT) at operative	24hrs post
Figure (20):	Summarizing the statistical significant between two groups regarding (Irrigation (dL) in three times (T1 and T2 and T3)	on Volume
Figure (21):	Summarizing the significant difference two groups regarding total irrigation volu	
Figure (22):	Summarizing the statistical significant between two groups regarding (F concentration in fluid (g/dL) in (T1: time T2: 4hrs post operative), and non difference in T3	Hemoglobin of surgery, significant
Figure (23):	Summarizing the significant difference two groups regarding total has concentration in irrigation fluid	nemoglobin
Figure (24):	Summarizing the statistical significant between two groups regarding (Blood lo three times (T1: immediately after su 4hrs post operative, T3: 24 hrs post operative)	ess (ml)) in ergery, T2:
Figure (25):	Summarizing the statistical significant between two groups regarding the total I	
Figure (26):	Summarizing the statistical significant between two groups regarding (HGB g/dl	
Figure (27):	Summarizing the statistical significant between two groups regarding (HCT %) of	

List of Abbreviations

Abb.	Full term
BNE:	.Bladder Neck Elevation
	.Bladder outlet obstruction
<i>BPH</i> :	. Benign prostatic Hyperplasia
<i>CABG</i> :	. Coronary Artery Bypass Graft
<i>CBC</i> :	. Complete Blood Picture
<i>CIC</i> :	. Coagulation Intermittent Cutting
<i>CR</i> :	. Creatinine
CRASH2:	. Clinical Randomization of antifibrinolytic insignificant head injury 2
<i>DHT</i> :	.Di Hydro Testosterone
<i>DJ</i> :	$. Double\ J\ stent$
<i>DM</i> :	.Diabetes mellitus
<i>DO</i> :	. Detrusor over activity
DRE:	Digital Rectal Examination
<i>DVT</i> :	. Deep Venous Thrombosis
	.Erectile dysfunction
<i>ER:YAG:</i>	. Erbium Yttrium-Aluminum-Garnet crystal
	laser
HCT:	
	. Hepatitis C virus
<i>HGB</i> :	
	.High intensity focal ultra sound
	.Heavy Menstrual Bleeding
	. Holmium Yttrium-Aluminum-Garnet Laser
	. Holmium laser ablation of prostate
	. Holmium laser enucleation of prostate
	. Holmium laser resection of prostate
HPS laser:	.High Performance system laser
<i>HTN</i> :	. Hypertension

List of Abbreviations (Cont...)

Abb.	Full term
<i>INR</i> :	.International normalized ratio
<i>IPSS</i> :	.International prostate symptoms score
<i>K</i> :	. Potassium
<i>KTP</i> :	. Potassium Titanyl Phosphate, Green light laser
<i>LBO</i> :	.Lithium tri borate crystals laser
<i>LUTS</i> :	.Lower Urinary Tract Symptoms
<i>NA</i> :	. Sodium
<i>ND:YAG:</i>	.Neodymium-doped Yttrium Aluminum Garnet
	crystal laser
<i>NSAID</i> :	.Non Steroidal Anti Inflammatory Drug
<i>PDE5</i> :	.Phosphodiesterase TYPE 5
<i>PDE5I</i> :	.Phosphor di esterase type 5 inhibitor
<i>PE</i> :	.Pulmonary Embolism
<i>PKI</i> :	.Predictive drug kinase inhibitor
<i>PLT</i> :	.Platelet
<i>PSA</i> :	. Prostatic specific antigen
Pt:	. Prothrombine time
Ptt:	. Partial thrombin time
<i>PVP</i> :	. Photo selective laser vaporization of prostate
<i>PVR</i> :	.Post voiding residual volume
<i>Q max:</i>	.Maximum flow rate
<i>QOL</i> :	. Quality of life
<i>RBCS</i> :	.Red Blood Cells
<i>RF</i> :	.Radio frequency current
<i>SD</i> :	. Standard Deviation
SIG:	. Significant
<i>T1:</i>	. Time immediately after surgery
<i>T</i> 2:	. Time 4 Hrs after surgery
<i>T3</i> :	. Time 24 hrs after surgery

List of Abbreviations (Cont...)

Abb.	Full term	
THA:	Total Hip Arthroplasty	
<i>TLC</i> :	Total Leukocyte Count	
<i>TPV</i> :	Total prostatic volume	
TRUS:	Trans urethral ultra sound	
<i>TUIP</i> :	Trans urethral incision of prostate	
<i>TUMT</i> :	Trans urethral microwave thermotherapy	
<i>TUNA</i> :	Trans urethral needle ablation	
TUR:	Trans urethral Resection	
<i>TURP</i> :	Trans urethral resction of prostate	
<i>TXA</i> :	Tranexamic Acid	
<i>US</i> :	United States	
<i>UTI</i> :	Urinary Tract Infections	
<i>VLAP</i> :	Visual laser aplation of prostate	

Introduction

PH is a nonmalignant enlargement of the prostate gland caused by cellular hyperplasia of the glandular and stromal elements. BPH leads to troublesome lower urinary tract symptoms in some men. Transurethral resection of prostate (TURP) is one of the most common and well-developed technologies used to treat BPH. It is recognized as the gold standard for surgical BPH treatments. (1)

Prostate tissues contain large venous sinuses which may cause some complications such as bleeding, TURP syndrome. Hemorrhage is the most common complication of this type of surgery. (1)

Tranexamic acid is a synthetic derivative of amino acid lysine and has strong affinity for lysine binding site of plasminogen. It inhibits the conversion of plasminogen to plasmin which is required for fibrinolysis. (2)

The intravenous route and the local administration of tranexamic acid, both have reduced blood loss effectively in some surgeries such as cardiac surgery, oral surgery, dental surgery, knee arthroplasty and endoscopic sinus surgery. (2)

The topical use of Antifibrinolytic agents such as tranexamic acid (TXA) demonstrates benefits without any significant risk, systemic absorption and thromboembolic disorders.