

# بسم الله الرحمن الرحيم



-Caron-





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





# جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

# قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغيار





### Cognitive Dysfunction in PTSD: Cross Sectional Descriptive Study in Recent Trauma Survivors in Egypt

#### Thesis

Submitted for Partial Fulfillment of Medical Doctorate Degree in **Psychiatry** 

#### By

Mohamed El Tohamy Atia Mohamed

M.B.B.Ch, M.SC

Under Supervision of

#### **Prof. Dr. Ahmed Saad Mohamed**

Professor of Psychiatry
Faculty of Medicine, Ain Shams University

#### Prof. Dr. Ghada Mohamed Abdel Razik

Professor of Psychiatry
Faculty of Medicine, Ain Shams University

#### Prof. Dr. Doha Mostafa El Serafi

Professor of Psychiatry
Faculty of Medicine, Ain Shams University

#### Prof. Dr. Maged Bahi El Den Mohamed

Head of Psychiatric Department Military Medical Academy

Faculty of Medicine, Ain Shams University
2021



سورة البقرة الآية: ٣٢

### Acknowledgments

First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.

I wish to express my deepest thanks, gratitude and appreciation to **Prof. Dr. Ahmed Saad**Mohamed, Professor of Psychiatry, Faculty of Medicine, Ain Shams University, for his meticulous supervision, kind guidance, valuable instructions and generous help.

Special thanks are due to **Prof. Dr. Ghada**Mohamed Abdel Razik, Professor of Psychiatry,
Faculty of Medicine, Ain Shams University, for her sincere efforts, fruitful encouragement.

I am deeply thankful to **Prof. Dr. Doha**Mostafa El Serafi, Professor of Psychiatry, Faculty
of Medicine, Ain Shams University, for her great help,
outstanding support, active participation and
guidance.

Really I can hardly find the words to express my gratitude to **Prof. Dr. Maged Bahi & Den Mohamed,** Head of Psychiatric Department, Military Medical Academy, for his supervision, continuous help, encouragement throughout this work.

I would like to express my hearty thanks to all my family & my lovely wife for their support till this work was completed.

Mohamed El Johanny Atia Mohamed

#### **ABSTRACT**

Posttraumatic stress disorder (PTSD) is a psychiatric syndrome that develops after exposure to terrifying and life-threatening events including warfare, motor-vehicle accidents, physical and sexual assault. It is characterized by intense reliving of the traumatic event through disruptive memories and nightmares, avoidance of reminders of the event, and hyper-vigilance toward potential threats in the environment. For most, the trauma is acute and transient and results in minimal functional impairment.

PTSD is characterized by affective and cognitive dysfunction. In later a long time, considerable proof from functional neuro-imaging has indicated that disruptions in prefrontal, limbic and interactive brain function may underlie the disorder.

PTSD has been connected to wide neuropsychological deficits over numerous domains. Many studies of PTSD have found that lower pre-morbid cognitive capacity may be a risk issue for the consequent advancement of the disorder.

Preservative behaviors are a common clinical marker of Executive functioning dysfunction. Perseveration concerning aspects of the trauma may be an important and basic contributing factor within the advancement and support of PTSD.

In spite of the fact that presentation to stressful and traumatic occasions is very common solely a little proportion of people in the long run, develop vital posttraumatic adjustment difficulties or PTSD.

The cognition affection is characterized by a variety of cognitive deficits that can start in infancy through youth, counting developmental delays in language and intellectual functioning, executive functioning, and attention.

Maltreated patients have been found to encounter incredible shortfalls in executive functioning skills as compared to non-maltreated controls.

Moreover, cognitions associated with the traumatic encounters may be thoughtabout inside this domain, counting attributions of fault, that are found to be related with more regrettable results of depression and PTSD.

The study outcomes for post-traumatic stress disorder in sample of Egyptian patients recently survived from trauma and there effect on cognitive functions.

We hypothesized that the presence of traumatic events may precipitate post traumatic disorder which may aggravate cognitive dysfunction in sample of Egyptian patients recently survived from trauma.

Keywords: Cognitive Dysfunction in PTSD: Cross Sectional Descriptive Study in Recent Trauma Survivors in Egypt

### Tist of Contents

Title	Page No.
List of Tables	i
List of Figures	iv
List of Abbreviations	v
Introduction	1
Rationale of the Study	3
Hypothesis	4
Aim of the Work	5
Review of Literature	
■ Trauma & Stress Related Disorders	6
Risks & Comorbidity of PTSD	28
■ PTSD in Relation to Cognitive Functions	43
Subjects and Methods	58
Results	70
Discussion	103
Strengths and Limitations	120
Conclusion	122
Summary	124
Recommendations	129
References	131
Arabic Summary	

## Tist of Tables

Table No.	Title	Page No.
Table 1:	Socio-demographic data among case history of exposure to trauma	
Table 2:	Posttraumatic stress diagnostic sca	le(PDS)
m 11 o	among cases with history of exposure to t	
Table 3:	History of psychiatric illness among exposed to trauma:	
Table 4:	Comparison between patients with P without PTSD as regard Socio-demograpusing Student t-test, Chi-Square test &	TSD & hic data
m 11 =	Exact test.	
Table 5:	Comparison between patients with P without PTSD as regard History of psy	
	illness using Chi-Square test	
Table 6:	Comparison between patients with P	
	without PTSD as regard PDS using Exact test & Chi-Square test	
Table 7:	Wechsler Memory Scale (WMS) amon	
	with history of exposure to trauma	-
Table 8:	Wechsler Adult Intelligence Scale	
T 11 0	among cases with history of exposure to t	
Table 9:	Wisconsin Sorting Card test (WSCT) among cases with history of exposure to t	
Table 10:	Comparison between patients with P	
	without PTSD as regard WMS using	
	Exact test	
Table 11:	Comparison between patients with P	
	without PTSD as regard WMS score Student t-test	<del>-</del>
Table 12:	Comparison between patients with P	
	without PTSD as regard WAIS using	
	Exact test & Chi-Square test	

## Tist of Tables cont...

Table No.	Title	Page No.	
Table 13:	Comparison between patients with I without PTSD as regard WAIS score Student t-test	es using	. 89
Table 14:	Comparison between patients with I without PTSD as regard WSCT score Student t-test	es using	90
Table 15:	Comparison between patients with PTS: without PTSD group & control group a WMS using Fisher's Exact test & Ch test	s regard i-Square	. 91
Table 16:	Comparison between patients with PTS2 without PTSD group & control group a WMS scores using One Way ANOVA Post-hoc Bonferroni test	D group, s regard test &	
Table 17:	Comparison between patients with PTS: without PTSD group & control group a WAIS using Fisher's Exact test & Ch	D group, s regard i-Square	
Table 18:	Comparison between patients with PTS without PTSD group & control group a WAIS scores using One Way ANOVA Post-hoc Bonferroni test	D group, s regard test &	
Table 19:	Comparison between patients with PTS2 without PTSD group & control group a WSCT scores using One Way ANOVA Post-hoc Bonferroni test	D group, s regard A test &	. 95
Table 20:	Cognitive dysfunction and Number of trevents using Fisher's Exact test & Chest	raumatic i-Square	.96
Table 21:	Cognitive dysfunction Mode and nutraumatic events between cognitive dysgroups within cases group using Fisher test & Chi-Square test	mber of sfunction r's Exact	

## Tist of Tables cont...

Table No.	Title Page 1	No.
Table 22:	Age & education between cognitive dysfunction groups using Chi-Square test	98
Table 23:	Physical injury between cognitive dysfunction groups using Chi-Square test	
Table 24:	Correlations between symptoms severity score and WSCT64 among 200 patients using pearson correlation test	100
Table 25:	Correlations between symptoms severity score and WMS among 200 patients using pearson correlation test	
Table 26:	Correlations between symptoms severity score and WAIS among 200 patients using pearson correlation test	

# List of Figures

Fig. No.	Title	Page No.
Figure 1:	HPA axis in PTSD	29
Figure 2:	Noradrenergic feed forward circuit	32

## Tist of Abbreviations

Abb.	Full term
<del>-</del>	sSerotonin receptor
5-HTTLPR	Hydroxytryptamine-transporter-linked
	$polymorphic\ Region$
	Acute Stress Disorder
	Adjustment Disorder
	Arcuate fasciculus
<i>ANOVA</i>	Analysis of variance tests
<i>AO</i>	Adult-Onset
<i>APA</i>	American psychiatry association
<i>ARMS</i>	A risk mental state
<i>ATR</i>	Anterior thalamic radiation
<i>ANKK1</i>	Ankyrin repeat and Kinase domain
	containing 1
<i>BDNF</i>	Brain-derived neurotropic factor
	British National Formulary
CNR1	A gene on chromosome 6q14-
	q15 that encodes one of two cannabinoid rec
	eptors,
<i>CBT</i>	Cognitive behavioral therapy
	Cortico-cortical evoked potentials
<i>CD</i>	Cluster of differentiation
	Cingulum hippocampus
<i>CGI</i>	Clinical Global Impression
<i>CNS</i>	Central nervous system
	Catechol-O-methyltransferase
_	Dopamine Receptor 2
	DopamineAgonist Receptor
	Dorsolateral prefrontal cortex
	Diagnostic and Statistical Manual of
	Mental Disorders, first edition
<b>DSM II</b>	Diagnostic and Statistical Manual of
	Mental Disorders, second edition

### **DSM III** ......Diagnostic and Statistical Manual of Mental Disorders, third edition

### Tist of Abbreviations cont...

Abb.	Full term
DOMESTIC	
DSM VI	Diagnostic and Statistical Manual of
DOM F	Mental Disorders, fourth edition
DSM-5	Diagnostic and Statistical Manual of
DDII	Mental Disorders, fifth edition
	Dopamine Beta-Hydroxylase
	Emotion Expression
	Electroence phalogram
	Echo Planar Imaging
	Emotion Recognition Task
	Fluoro-dopamine (fluoro-phenylalanine)
<i>FH.</i>	Family history
<i>FMRI</i>	Functional MRI
FKBP5	FK506 binding protein of 12 kDa
$G^X E$	Gene verses Environment Eposure
<i>GABA</i>	$ Gamma \hbox{-} amino \hbox{-} buteric \hbox{-} acid$
<i>GAD</i>	Generalized anxiety disorder
<i>GAD</i>	Glutamic acid decarboxylase
<i>GAF</i>	Global Assessment Function
GABRA2	GABA transporter
GCCR	Gene Encoding Glucocorticoid
<i>GRD</i>	Genetic risk and deterioration syndrome
GWAS	Genome-wide association study
	. Hypothalamic-Pituitary-Adrenal
	Hippocampal sclerosis
	Inferior fronto-occipital fasciculus
<i>IL</i>	
<i>IQ</i>	Intelligence quotient
KYNA	~ <u>-</u>
<i>MD</i>	
	Major Histocompatibility Complex

MNI ......Standard Montreal Neurological Institute
MRI ......Magnetic resonance imaging

### Tist of Abbreviations cont...

Abb.	Full term
MRS	.Magnetic resonance spectroscopy
<i>MS</i>	
	National Alliance on Mental Illness
	North American Prodrome Longitudinal
	Study
<i>NART</i>	National reading test
<i>NCCMH</i>	.National Collaborating Centre for Mental
	Health
<i>NCCT</i>	.Sodium-chloride symporter ( <u>N</u> a+- <u>C</u> l- co-
	transporter)
NEa1	Noradrenergic alpha 1
<i>NHS</i>	National Health Service
<i>NICE</i>	National Institute for Health and Clinical
	Excellence
<i>NPY</i>	.Neuropeptide Y
<i>NLLS</i>	Non-linear Least Squares
<i>NMDAr</i>	.N-methyl-Daspartate receptor
	.N-methyl-d-aspartic acid receptor
<i>NO</i>	
	.NADPH oxidase pathway
NRG-1	•
	Outreach and support in South London
	.Obsessive compulsive disorder
	Ordinary Least Squares
OR	
<i>PCR</i>	.Polymerase chain reaction
	.Positron emission tomography
PROPELLER	.Periodically rotated overlapping parallel
	lines with enhanced reconstruction
	.Post-traumatic stress disorder
<i>PUFAs</i>	.Polyunsaturated fatty acids