

بسم الله الرحمن الرحيم



-Call 4000





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة يعبدا عن الغبار













بالرسالة صفحات لم ترد بالأصل





Cardiovascular Manifestations of Behçet Disease among Egyptian Patients

Thesis

Submitted for Partial Fulfillment of Master Degree in **Internal Medicine**

By

Bassant Megahed Ibrahim Hassanin M.B.B.Ch

Under Supervision of

Prof. Dr. Dalia Fayez Mohammed

Professor of Internal Medicine & Rheumatology Faculty of Medicine, Ain Shams University

Ass. Prof. Dr. Nashwa Aly El-morshedy

Assistant Professor of Internal Medicine & Rheumatology Faculty of Medicine, Ain Shams University

Dr. Mohamed Abdelmoneam Teama

Lecturer of Internal Medicine & Rheumatology Faculty of Medicine, Ain Shams University

Faculty of Medicine - Ain Shams University 2020



سورة البقرة الآية: ٣٢

Acknowledgments

First and foremost, I feel always indebted to Allah the Most Beneficent and Merciful.

I wish to express my deepest thanks, gratitude and appreciation to Prof. Dr. Dalia Fayez Mohammed, Professor of Internal Medicine & Rheumatology, Faculty of Medicine, Ain Shams University, for her meticulous supervision, kind guidance, valuable instructions and generous help.

Special thanks are due to Prof. Dr. Nashwa Aly Elmorshedy, Assistant Professor of Internal Medicine & Rheumatology, Faculty of Medicine, Ain Shams University, for her sincere efforts, fruitful encouragement.

I am deeply thankful to Dr. Mohamed Abdelmoneam Teama, Lecturer of Internal Medicine & Rheumatology, Faculty of Medicine, Ain Shams University, for his great help, outstanding support, active participation and guidance.

I would like to express my hearty thanks to all my family and friends for their support till this work was completed.

Bassant Megahed Ibrahim Hassanin

List of Contents

| Title | Page No. |
|--|----------|
| List of Tables. | 5 |
| List of Figures | 8 |
| List of Abbreviations | 12 |
| Introduction | 1 - |
| Aim of the Work | 3 |
| Review of Literature | |
| ■ Behçet's Disease | 4 |
| Cardiovascular Manifestation of Behcet's Disease | 61 |
| Patients and Methods | 95 |
| Results | 102 |
| Discussion | 138 |
| Summary and Conclusion | 149 |
| References | 153 |
| Arabic Summary | |

List of Tables

| Table No. | Title Pag | e No. |
|-------------|--|-----------|
| Table (1): | The distribution of clinical findings amo different countries | _ |
| Table (2): | The international criteria for Behce disease | |
| Table (3): | Updated European League again Rheumatism recommendations for t management of Behçet's syndrome | he |
| Table (4): | Treatment of major vessel disease in Behce disease | |
| Table (5): | The international criteria for behce disease | |
| Table (6): | Behçet's Disease Current Activity Form | 97 |
| Table (7): | BDCAF score | 98 |
| Table (8): | Distribution of demographic data amo studied behcet's patients | _ |
| Table (9): | Distribution of cumulative clinical data among the studied behoet's patier according to international study group (IS criteria | nts G) |
| Table (10): | Distribution of Behcet Disease Curre Activity Form (BDCAF) score among studi behcet patient | ed |
| Table (11): | Distribution of cardiovascul manifestations in studied behcet's patients | |
| Table (12): | Distribution of Laboratory investigations our studied behcet's patients | |
| Table (13): | Distribution of lipid profile among studi behcet's patients | |

List of Tables cont...

| Table No. | Title | Page No. |
|-------------|---|--------------------|
| Table (14): | Distribution of ECG findings in our behcet's patients | |
| Table (15): | Distribution of Echo findings among behcet's patients | |
| Table (16): | Distribution of carotid artery duplex famong studied behcet's patients | _ |
| Table (17): | Distribution of most common immunosuppressive drugs among behcet's patients. | studied |
| Table (18): | Correlation between Disease act BDCAF score) and laboratory fine TLC,Hb,PLT,ESR,CRP) | dings (|
| Table (19): | Relation of Disease activity (BDCAF and cardiovascular findings in ECG, and Carotid duplex and lipid profile studied patients | ECHO in our |
| Table (20): | Correlation between BDCAF (activity) and lipid profile, carotid dup echo findings | lex and |
| Table (21): | Relation between lipid profile and echo, carotid duplex findings in our patients | studied |
| Table (22): | Correlation between lipid profile para and carotid intimal thickening and echo parameters | dother |
| Table (23): | Relation between anterior uveitis an ECHO, Carotid duplex findings, vendarterial vascular affection in our behoet's patients | ous and studied |

List of Tables cont...

| Table No. | Title | Page No. |
|-------------|--|---------------------|
| Table (24): | Relation between posterior uveitis ar ECHO, carotid duplex, vascular arte venous affection in our study group | rial and |
| Table (25): | Relation between genital ulcers an ECHO and carotid duplex findings, arterial and venous affection in our behcet's patients | vascular studied |
| Table (26): | Relation between skin lesions and Echo, carotid duplex findings, arterial and venous affection in our behcet's patients | vascular studied |
| Table (27): | Relation between neumanifestations and ECG, ECHO, duplex findings, vascular arterial and affection in our studied behaet's patie | carotid l venous |
| Table (28): | Relation between pathergy test an ECHO, carotid duplex findings, arterial and venous affection in our behoet's patients | vascular studied |
| Table (29): | Relation between lipid profile and c clinical data of ISG criteria | |

List of Figures

| Fig. No. | Title | Page No. |
|--------------|---|---------------------------|
| Figure (1): | The role of autoimmune T cells cytokines in the pathogenesis of Bedisease | hcet's |
| Figure (2): | Interaction between multiple facto Behcet's disease | |
| Figure (3): | Oral ulcers Behcet's disease | 13 |
| Figure (4): | Genital ulcers Behcet's disease | 14 |
| Figure (5): | Mucocutaneous lesions of Behcet's di | sease17 |
| Figure (6): | A person with hypopyon which caseen in anterior uveitis in a person Behçet's disease | with |
| Figure (7): | A. Vitreous involvement with s diffuse vitritis. B. Inferior prerestring of pearls" vitreous exu tending to accumulate along the posedge of the vitreous base | etinal dates terior |
| Figure (8): | A. Fundus photograph. B. Fluore angiography of the same patie month after initial examination | ntone |
| Figure (9): | Pulmonary angiography shows a pulmonary artery aneurysm | _ |
| Figure (10): | Coronal T2 images of a patient shother pontine—midbrain lesion externation both sides | nding |
| Figure (11): | Comparison between intestinal B Disease and Crohn's Disease | - |
| Figure (12): | Pathergy Test | 42 |
| Figure (13): | Pathergy reaction | 42 |

List of Figures cont...

| Fig. No. | Title | Page No. |
|--------------|---|-------------------|
| Figure (14): | Thrombosis is a result of multifaction derangements in systemic inflamm fibrinolysis, and the endothelium | ation, |
| Figure (15): | Neck, thoracic, and abdominal colla in a BD patient with long-standing cava superior thrombosis | vena |
| Figure (16): | Post-thrombotic syndrome (chevenous insufficiency) | |
| Figure (17): | Aortography shows a huge aneury ascending aorta | |
| Figure (18): | Multislice CT scan shows a pseudoaneurysm of aortic arch | |
| Figure (19): | MRI scan discloses multiple pulm artery aneurysms (arrow) on descending pulmonary arteries of right lower lobe in a patient with BD | the f the |
| Figure (20): | MRI scan shows pulmonary a thrombosis on the descending branch the right pulmonary artery in a pawith BD (arrow) | ach of atient |
| Figure (21): | Transthoracic echocardiography in parasternal short-axis view show thrombus (arrow) in the right ven attached to the interventricular sept | ws a ntricle |
| Figure (22): | Contrast-enhanced pulmonary angiography shows acute embolus and mildly enlarging the right middle artery (arrow). | filling e lobe |

List of Figures cont...

| Fig. No. | Title | Page | No. |
|--------------|---|-----------------|-----|
| Figure (23): | Thorax CT scan shows multiple grass lesions compatible with hemore in a BD patient with pulmonary a involvement | rhage artery | 89 |
| Figure (24): | Sex distribution among studied beh | | 103 |
| Figure (25): | Distribution of cumulative clinical among studied behoet patients according to international study group (ISG) | rding | 105 |
| Figure (26): | Distribution of lipid profile in our st behcet's patients. | | 110 |
| Figure (27): | Distribution of different parameter lipid profile among studied be patients. | hcet's | 110 |
| Figure (28): | ECG changes among patients | | 111 |
| Figure (29): | Distribution of Valvular affection a our studied patients | | 114 |
| Figure (30): | Distribution of Diastolic Dysfunction intracardiac masses among our st behcet patients. | udied | 114 |
| Figure (31): | Distribution of carotid artery d findings in our studied patients | | 115 |
| Figure (32): | Distribution of most common immunosuppressive drugs among studied behcet's patients | our | 117 |
| Figure (33): | Increased incidence of Mitral Re with disease activity | | 120 |