

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

## بسم الله الرحمن الرحيم





MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



MONA MAGHRABY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

## جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



MONA MAGHRABY



# Role of Diffusion Weighted MRI in Diagnosis of Breast Masses (BIRADS III and BIRADS IV)

#### Thesis

Submitted for Partial Fulfillment of Master Degree in **Diagnostic and Interventional Radiology** 

By

Shimaa Maghawry El Sayed
M.B.B.Ch., Ain Shams University

Under Supervision of

#### **Prof. Dr. Mounir Sobhy Guirguis**

Professor of Radio diagnosis Faculty of Medicine, Ain Shams University

#### Dr. Ali Hagag Ali Nour El Deen

Lecturer of Radio diagnosis Faculty of Medicine, Ain Shams University

Faculty of Medicine, Ain Shams University
2021



سورة البقرة الآية: ٣٢

#### Acknowledgments

First and foremost, I feel always indebted to **Allah** the Most Beneficent and Merciful.

I wish to express my deepest thanks, gratitude and appreciation to **Prof. Dr. Mounir Sobhy**Guirguis, Professor of Radio diagnosis, Faculty of Medicine, Ain Shams University, for his meticulous supervision, kind guidance, valuable instructions and generous help.

Special thanks are due to **Dr. Ali Haqaq**Ali Mour El Deen, Lecturer of Radio diagnosis,

Faculty of Medicine, Ain Shams University, for his sincere efforts, fruitful encouragement.

I would like to express my hearty thanks to all my family for their support till this work was completed.

Shimaa Maghawry El Sayed

#### Tist of Contents

Title	Page No.
List of Tables	i
List of Figures	iv
List of Abbreviations	xii
Introduction	1
Aim of the Work	3
Review of Literature	
Pathological Considerations	4
MRI Imaging of the Breast	25
Patients and Methods	58
Results	63
Case Presentation	93
Discussion	120
Summary and Conclusion	130
References	133
Arabic Summary	

#### Tist of Tables

Table No.	Title	Page No.
Table (2-1):	Enhancement patterns of the brea	ast lesions 35
<b>Table (4-1):</b>	Shows the range, mean and deviation of the patient's age	
<b>Table (4-2):</b>	Shows the site and size of breast l	esions 63
<b>Table (4-3):</b>	Shows the echogenicity, shape, m denisty of breast lesions	-
<b>Table (4-4):</b>	Shows the breast lesions which a with secondary sign (edema retraction)	or nipple
<b>Table (4-5):</b>	Shows the breast lesions which a with microcalcification	
Table (4-6):	Shows the mammography and u BIRADS classification of the breast lesions	examined
<b>Table (4-7):</b>	Shows the signal intensity of the lesions in T1WI and T2WI	
<b>Table (4-8):</b>	Shows MRI enhancement charact the breast lesions	
<b>Table (4-9):</b>	Shows the type of time/signal curve pattern of enhancement lesions.	of breast
Table (4-10):	Shows characters of the breast I DW images as restricted or diffusion and the mean, standard and range of the ADC values of the lesions	facilicated deviation the breast
<b>Table (4-11):</b>	Shows the MRI BIRADS classif breast lesions	

#### Tist of Tables cont...

Table No.	Title	Page No.
Table (4-12):	Shows the histopathological anal breast lesions.	•
Table (4-13):	Shows the mean, standard devirange of the patient's age regardenign and malignant breast lesion	arding the
<b>Table (4-14):</b>	Shows the site and size of t lesions in correlation whistopathology results	rith the
Table (4-15):	Shows the echogenicity, shape, medical density of the breast lesions in density with the histopathology results	correlation
Table (4-16):	Shows the breast lesions whas associated with secondary a microcalcification or not	sign and
<b>Table (4-17):</b>	Shows the ultrasound and mam BIRADS classification of the brea in comparison to the histopatholo	ast lesions
<b>Table (4-18):</b>	Table shows the signal intensity enhancement and time/signal curve	intensity
Table (4-19):	Shows the characteristics of the lesions on DW images as well as range and standard deviation of values in benign and malignatesions	the mean, f the ADC int breast
Table (4-20):	Shows the MRI BIRADS classic the breast lesions in correlations that the breast lesions in correlations are the breast lesions.	tion with

#### Tist of Tables cont...

Table No.	Title	Page No.
Table (4-21):	Correlation between age and s breast lesions to the ADC values.	
Table (4-22):	Shows the echogenicity of the lesions in relation to the ADC value.	
Table (4-23):	Shows the shape and margin lesions in relation to the ADC value	
Table (4-24):	Shows the breast lesions which with secondary sign and microca not with relation to the ADC value.	lfication or
Table (4-25):	Shows the density of breast relation to the ADC values	
Table (4-26):	Shows the ultrasound and mam BIRADS classification of the bre in relation to the ADC va histopathology	ast lesions alues and
Table (4-27):	Shows classification of the detections according to their counenhanced breast MRI in rahistopathology.	riteria on elation to
Table (4-28):	Shows the characters of the m lesions on DW images in r histopathology	elation to
Table (4-29):	Shows the type of time/signal cur of enhancement in relation to values	the ADC
Table (4-30):	Shows the ADC values of the brein relation to the MRI classification and histopathology	BIRADS

## List of Figures

Fig. No.	Title	Page No.
Fig. (1.1):	a & b-Microscopic and gross picture fibroadenoma, respectively	
Fig. (1.2):	Intermediate grade DCIS	
Fig. (1.3):	a & b-Microscopic and gross pictures o	
<b>g</b> ; (_1,0);	respectively	
Fig. (1.4):	Microscopic picture of Paget's disease.	
Fig. (2.1):	Double breast MRI coil	
Fig. (2.2):	Typical set up for breast MRI	
Fig. (2.3):	Pre-contrast T2 of a left breast simple	
<b>g</b> : (=:0);	(arrow)	-
Fig. (2.4):	Pre-contrast T2 of a left breast spec mass proved to be invasive ductal	ulated
	cancer (arrow)	28
Fig. (2.5):	Fat suppression makes enhancing leasier to appreciate	
Fig. (2.6):	An example of the 3D FLASH sequence	e29
Fig. (2.7):	MIP of subtracted images coronal	view
	points out a nodular enhancement i	in the
	right breast (arrow)	31
Fig. (2.8):	53-year-old woman with palpable nod	lule of
	right breast	32
Fig. (2.9):	Axial T2-weighted image of the breast	33
Fig. (2.10):	Sagittal T2 -fat suppression MR im-	age of
	the breast	33
Fig. (2.11):	Axillary LNs in sagittal T2 fat supp	$\operatorname{ressed}$
	MRI	34
Fig. (2.12):	Contrast enhanced MRI subtraction	image
	showing a non-mass like enhancemen	
	distribution in the right breast, proved	
	lobular carcinoma in situ	
Fig. (2.13):	Internal enhancement of focal mass by	MRI 38
Fig. (2.14):	Malignant features	39

Title	Page No.
· =	
Type III enhancement curve in	filtrating
Diffusion of water molecules	43
2 symmetric motion-probing gradies into a single-shot SE T2-weighted s	nt pulses sequence,
which is hyperintense in (a), the ab-value (1000 s/mm²) DW-MR im dark on (b), ADC maps represent ADC values (restricted dark of the control o	xial high age, and ting low iffusion),
A 53-year-old woman with ductal card situ. The axial T1-weighted image segmentally distributed, asymmetric, intensity, non-mass lesion in the left. The maximal intensity projection subtraction image shows heterogeneous enhancement in the left breast. diffusion-weighted image shows the lesion to be a high-signal intensity. The axial plane apparent diffusion of map shows a mixed green and yellow the apparent diffusion coefficient value.	cinoma in ge shows iso-signal breast. <b>B</b> . n of a us clumpy <b>C.</b> The he main lesion. <b>D.</b> coefficient area and ue of this
	Three types of enhancement kinetic with breast MRI

Fig.	No.	Title	Page No.
Fie	(2.24):	Fibroadenoma	50
_	(2.24): (2.25):	False-negative result in a patier	
rıg.	(2.20).	invasive ductal carcinoma	
Fig.	(2.26):	Intraductal papilloma	
_	(2.27):	Mucinous carcinoma	
_	(2.28):	Fibroadenoma in the left breast	
_	(4.1):	The mean ages of the patients in ben	
		malignant breast lesions	-
Fig.	<b>(4.2):</b>	The echogenicity of breast lesi	ons in
		correlation to the histopathology resu	ılts 73
Fig.	<b>(4.3):</b>	The shape and margin of the breast	lesions
		in correlation to the histopathology r	
Fig.	<b>(4.4)</b> :	The ultrasound and mamme	0 1 0
		BIRADS classification of the breast	
	>	in comparison to the histopathology.	
Fig.	<b>(4.5)</b> :	The time/signal intensity curve par	
		enhancement of the breast lesi	
TC:	(4.0).	correlation to the histopathology	
rıg.	<b>(4.6)</b> :	The characters of the breast lesions	
Tie.	(4.7).	images in correlation to the histopath ADC value of breast lesions in correlations.	
rıg.	<b>(4.7):</b>	with histopathology	
Fig	(4.8):	MRI BIRADS classification of the	
1 15.	(4.0).	lesions in correlation with	
		histopathology	
Fig.	<b>(4.9)</b> :	Correlation between age and size	
- <b>o</b> *	/ -	ADC values	
Fig.	(4.10):	Relation between the ADC values a	
J		echogenicity of the breast lesions	82
Fig.	<b>(4.11)</b> :	Relation between the ADC valu	es and
_		shape, margin of the breast lesions	84

Fig. No.	Title	Page No.
Fig. (4.12):	The ADC values of the breast less correlation to ultrasound	and
Fig. (4.13):	mammography BIRADS classification Relation of the ADC values time/signal intensity curve of the lesions	to the breast
Fig. (4.14):	Relation of the ADC values to the BIRADS classification	ne MRI
Fig. (4.15):	Relation of the ADC values histopathology	to the 91
Fig. (4.16):	Receiver operating characteristic (ROC) for ADC value to differ between BIRADS III and BIRADS IV	curve rentiate
Fig. (5.1):	Ultrasound showing hypoechoic well lesion in the left breast	
Fig. (5.2):	MRI axial images showing lower quadrant oval well defined mass displaying lowT1 (A) and hetero	lesion ogenous
Fig. (5.9).	(bright and intermediate) T2 (B)	
Fig. (5.3):	MRI Axial A) Dynamic THRIVE sequ	
Fig. (5.4):	Diffusion sequence and post processismap	
Fig. (5.5):	Mammography showing well circum mass with lucent halo in the right	nscribed t upper
Fig. (5.6):	outer quadrant  Complementary ultrasound showin	g right
Fig. (5.7):	breast well defined hypoechoic mass.  MRI axial images well defined righ outer mass lesion displaying low T1	t upper (A) and
F! (F.0)	bright T2 with fat saturation (B)	
Fig. (5.8):	MRI axial A) Dynamic THRIVE sequ	
Fig. (5.9):	Diffusion sequence and post procession	_

Fig. No.	Title Page No.
Fig. (5.10):	Mammography showing edematous breast changes as thickened skin, coarsening trabeculations and increased breast density99
Fig. (5.11):	Complementary ultrasound showing right breast ill defined hypoechoic irregular lump 99
Fig. (5.12):	MRI axial images show right outer central ill defined lump of low T1 (A) and intermediate T2 with fat saturation (B)., associated with edematous changes
Fig. (5.13):	MRI sagittal A) Dynamic THRIVE sequence. B) Subtraction image Right outer central heterogenous ill defined lump with edematous breast changes. C) Time/signal intensity analysis of ROI
Fig. (5.14):	Diffusion sequence and post processing ADC map101
Fig. (5.15):	Mammography showing left breast well-defined mixed density lesion, predominantly hyperdense
Fig. (5.16):	Complementary ultrasound shows left breast well defined solid mass of mixed echogenicity
Fig. (5.17):	MRI axial images left breast well defined solid mass shows heterogenous hyperintense signal in STIR (A) and low
Fig. (5.18):	signal in T2 (B)