

بسم الله الرحمن الرحيم





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بعض الوثائق الأصلية تالفة





بالرسالة صفحات
لم ترد بالأصل





Addition of intrathecal atropine for prevention of postoperative nausea and vomiting in patients received intrathecal bupivacaine / morphine in lower limb surgeries

Thesis

Submitted for partial fulfillment of the Master
Degree in Anesthesia

By

Ramez Mohamed Kamel Amin Elkady

M.B.B. Ch, Faculty of Medicine, Assiut University

Under supervision of

Prof. Ibrahim Abd Elghani Ramadan

Professor of Anesthesiology, Intensive Care & Pain Management
Faculty of Medicine – Ain Shams University

Dr. Ashraf Elsayed Elagamy

Assistant Professor of Anesthesiology, Intensive Care & Pain
Management
Faculty of Medicine – Ain Shams University

**Dr. Wael Sayed Ahmed Abd Elghaffar
Elgharabawy**

Lecturer of Anesthesiology, Intensive Care & Pain Management
Faculty of Medicine – Ain Shams University

**Faculty of Medicine
Ain Shams University
2020**



First of all, thanks to **Allah** whose magnificent help was the main factor in completing this work.

No words could express my deepest thanks and appreciation to **Prof. Ibrahim Abd Elghani Ramadan**, Professor of Anesthesiology, ICU and Pain Management, Faculty of Medicine, Ain Shams University, for inspiring me with the idea of this work. his patience, precious advice and guidance enlightened my way throughout this work.

I want also to express my profound gratitude to **Dr. Ashraf Elsayed Elagamy**, Assistant Professor of Anesthesiology, ICU and Pain Management, Faculty of Medicine, Ain Shams University, for his patience, valuable advice and continuous help in completing this work.

I am also deeply indebted to **Dr. Wael Sayed Ahmed Abd Elghaffar Elgharabawy**, Lecturer of Anesthesiology, ICU and Pain Management, Faculty of Medicine, Ain Shams University, for his kind help, guidance, useful advices, continuous encouragement and support all through my entire work.

Finally, my deepest thanks to all professors, colleagues and my family especially my wife **Marwa Elkady** who helped me in the production of this work.

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

سببناك لا تعلم لنا
إلا ما علمتنا إنك أنت
العليم العظيم

صدقة الله العظيم

سورة البقرة الآية: ٣٢

Contents

Subjects	Page
List of abbreviations.....	II
List of figures.....	IV
List of tables.....	VI
• Introduction	1
• Aim of the Work.....	3
• Review of Literature	
♦ Chapter (1): Anatomy of spinal cord and vertebral column	4
♦ Chapter (2): Technique of intrathecal anesthesia	10
♦ Chapter (3): Pharmacology of local anesthetics and additives of intrathecal anesthesia	15
♦ Chapter (4): Postoperative nausea and vomiting.....	30
• Patients and Methods.....	37
• Results.....	46
• Discussion.....	71
• Conclusion.....	74
• Summary.....	77
• References.....	80
• Arabic Summary.....	\

List of Abbreviations

5 HT	5-HydroxyTryptamine/serotonin
$\alpha 2$	Alpha 2
ACL	Anterior Cruciate Ligament.
ASA	The American Society of Anesthesiologists
CBC	Complete Blood Count
CL	Clearance
Cm	Centimeter
CNS	Central Nervous System
CSF	Cerebro Spinal Fluid
CTZ	Chemoreceptor Trigger Zone
δ	Delta
DHS	Dynamic Hip Screw
ECG	Electrocardiography
GI	Gastro Intestinal
H	Hour
IM	Intramuscular
IT	Intrathecal
IV	Intravenous
κ	Kappa opioid receptor
Kg	Kilogram

L	Liter
Mcg	Microgram
Mg	Milligram
Min	Minute
ml	Milliliter
mmHg	Millimeter mercury
MW	Molecular Weight
Na+	Sodium
NK-1	Neurokinin
NMDA	N-Methyl-D-aspartic Acid
μ	Mu opioid receptor
O2	Oxygen
PCO 2	Arterial Partial Pressure of Carbon Dioxide
PFN	Proximal Femur Nail
pH	Hydrogen ion measure
PONV	Post-Operative Nausea and Vomiting
SAMBA	The Society of Ambulatory Anesthesiology
T1/2	Half life
TDS	Transdermal Scopolamine
VAS	Visual Analogue Scale
Vdss	Volume of distribution at steady state.

List of Figures

<u>No.</u>	<u>Figure</u>	<u>Page</u>
<u>1</u>	Spinal cord anatomy. Notice the termination of the spinal cord	5
<u>2</u>	Spinal cord within its meninges.	7
<u>3</u>	The spinal cord in transverse section from a thoracic segment.	8
<u>4</u>	Sitting position for spinal anesthesia.	11
<u>5</u>	Lateral position for spinal anesthesia.	11
<u>6</u>	Cross section of the spinal cord and adjacent ligaments.	13
<u>7</u>	Technique of spinal anesthesia.	14
<u>8</u>	Structure of Bupivacaine.	18
<u>9</u>	Structure of Morphine.	22
<u>10</u>	Effect of Morphine	25
<u>11</u>	Structure of Atropine.	28
<u>12</u>	The inputs and receptors involved in causing PONV.	31
<u>13</u>	Age among BMA group and BM group.	46
<u>14</u>	Heart rate among the studied cases.	51
<u>15</u>	Systolic blood pressure among the studied cases	54
<u>16</u>	Diastolic blood pressure among the studied cases.	54
<u>17</u>	Respiratory rate among the studied cases.	56

<u>No.</u>	<u>Figure</u>	<u>Page</u>
<u>18</u>	O2 saturation (%) among the studied cases.	58
<u>19</u>	Incidence of postoperative nausea in both groups.	60
<u>20</u>	Incidence of postoperative vomiting in both groups.	62
<u>21</u>	Total number of postoperative nausea and vomiting episodes in both groups.	64
<u>22</u>	Time of first anti-emetic (min) and frequency of subsequent doses for the BM group and the BMA group.	66
<u>23</u>	Xerostomia and pruritis among the BM group and the BMA group.	69

List of Tables

<u>No.</u>	<u>Table</u>	<u>Page</u>
<u>1</u>	Pharmacokinetic properties of several amide local anesthetics.	17
<u>2</u>	Physicochemical and pharmacokinetic properties of bupivacaine.	19
<u>3</u>	Additives used for receptor-mediated analgesic effect.	20
<u>4</u>	Opioid receptor subtypes, their functions, and their endogenous peptide affinities.	23
<u>5</u>	Simplified Apfel scoring system.	32
<u>6</u>	Degree of motor block assessed by modified bromage scale.	42
<u>7</u>	Intensity of pain assessed by VAS (visual analog scale).	42
<u>8</u>	Comparison between BMA group and BM group according to demographic data.	45
<u>9</u>	Comparison between BMA group and BM group according to ASA classification.	46
<u>10</u>	Comparison between BMA group and BM group according to type and duration of operation.	47
<u>11</u>	Comparison between BMA group and BM group according to sensory level and block degree at 15 min.	48

<u>No.</u>	<u>Table</u>	<u>Page</u>
<u>12</u>	Comparison between BMA group and BM group according to time of regression and total duration of motor.	49
<u>13</u>	Comparison between BMA group and BM group according to heart rate (beat/minute) among the studied cases.	50
<u>14</u>	Comparison between BMA group and BM group according to systolic blood pressure (mmHg).	52
<u>15</u>	Comparison between BMA group and BM group according to diastolic blood pressure (mmHg).	53
<u>16</u>	Comparison between BMA group and BM group according to Respiratory rate (breath/minute).	55
<u>17</u>	Comparison between BMA group and BM group according to O2 saturation (%).	57
<u>18</u>	Comparison between BMA group and BM group according to postoperative nausea.	59
<u>19</u>	Comparison between BMA group and BM group according to postoperative vomiting.	61
<u>20</u>	Comparison between BMA group and BM group according to total number of postoperative nausea and vomiting episodes.	63
<u>21</u>	Comparison between BMA group and BM group according to time of first anti-emetic	65