

شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلو

# بسم الله الرحمن الرحيم





HANAA ALY



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكرونيله



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



HANAA ALY



شبكة المعلومات الجامعية التوثيق الإلكترونى والميكروفيلم

# جامعة عين شمس التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HANAA ALY





# Relation between postmenopausal osteoporosis and IL-17 serum level

#### Thesis

Submitted in Partial Fulfillment of the Requirement of the Master Degree in Physical Medicine, Rheumatology and Rehabilitation

### 

M.B., B.Ch. Faculty of Medicine- Ain Shams University

### Supervisors

### Prof. Dr. Nagwa Mohammed Mahmoud Nassar

Professor of Physical Medicine, Rheumatology & Rehabilitation Faculty of Medicine - Ain Shams University

### Prof. Dr. Ola Abd El-Nasser Abd El-Aziz

Professor of Physical Medicine, Rheumatology & Rehabilitation Faculty of Medicine - Ain Shams University

#### Dr. Reem Mohammed Ezz El-Deen Elmallah

Assistant Professor of Physical Medicine, Rheumatology & Rehabilitation Faculty of Medicine - Ain Shams University

> Faculty of Medicine Ain Shams University 2021



سورة البقرة الآية: ٣٢

## Acknowledgment

First and foremost, my deep gratefulness and indebtedness is to **Allah**, "The Most Gracious and The Most Merciful".

I would like to express my sincere gratitude to **Prof. Dr.**Nagwa M. M. Nassar Prof. of Physical Medicine,
Rheumatology & Rehabilitation, Faculty of Medicine, Ain

Shams University who offered me advice during the present work. I will appreciate her continuous guidance, valuable directions, extensive revision, objective criticism, and she was helping me to overcome many difficulties during the study.

In addition, my deep appreciation and sincere thanks to **Prof. Dr. Ola Abd El-Nasser Abd El-Aziz** Prof. of Physical Medicine, Rheumatology & Rehabilitation, Faculty of Medicine, Ain Shams University for her valuable and expert supervision. She offered me advice during the present work. Her kind insight and encouragement helped me to bring out the best work.

Furthermore, I would like to express my sincere gratitude to **Dr. Reem Mohammed Ezz El-Deen Elmallah,** Assistant professor of Physical Medicine, Rheumatology & Rehabilitation, Faculty of Medicine, Ain Shams University, for her continuous guidance, sincere help during the stages of this work, and for her continuous advice and encouragement during the preparation of the thesis.

At last, but not least all thankfulness, deep respects to members of Physical Medicine, Rheumatology & Rehabilitation department, Faculty of Medicine, Ain Shams University, central Tanta University laboratory where this study was done.

Samar Samy El-Hoseiny

# List of Contents

Title	Page No.
List of Abbreviations	i
List of Tables	iv
List of Figures	vi
Abstract	viii
Introduction	1
Aim of the Work	3
Review of Literature	
Osteoporosis in Postmenopausal Women	4
The IL-17 Family of Cytokines	41
Patients and Methods	54
Results	68
Discussion	84
Summary	100
Conclusion	104
Recommendations	105
References	106
Appendices	135
Arabic Summary	

### List of Abbreviations

Abb.	Full term
AP1	Activator protein 1
	Adenomatous polyposis coli
	Ankylosing spondylitis
	Area under curve
	B cell activating factor
	Bone mineral density
	Bone mineral densities
bone ALP	Bone alkaline phosphatase
	Calmodulin-dependent protein kinase
<b>CBP</b>	CREB binding protein
	Cluster of differentiation 4+
<b>CRD</b>	Cysteine rich domain
<b>CREB</b>	cAMP-response element-binding protein
<b>CTX</b>	C-telopeptides of type 1 collagen
<b>DC</b>	Dendritic cells
Dkk	Dickkopf
	Dual photon absorptiometry
	Deoxypyridinoline
	Dual energy X-ray absorptiometry
ELISA	Enzyme-linked immunosorbent assay
	Extracellular signal regulated kinase
<b>ESCEO</b>	European society for clinical and economic
	aspect of osteoporosis and osteoarthritis
	Enterotoxigenic Bacteroides fragilis
	Food and drug administration
	Fracture risk assessment tool
FZD	
	Glycogen synthase kinase 3β
	Human mesenchymal stem cells
IBD	Inflammatory bowel disease

### List of Abbreviations Cont...

Abb.	Full term
ΙΕΝ-γ	Interferon-v
IKK	
IL-17	
IL17Rs	
	3 innate lymphoid cells
	.International osteoporosis foundation
	Interquartile Range
JAK	
LEF1	Lymphocyte enhancer factor 1
	Lipopolysaccharide
LRP5	lipoprotein receptor related protein 5
LRP5/6	Low-density lipoprotein receptor-related
	protein 5/6
MAPKs	Mitogen activated protein kinase
M-CSF	Macrophage colony stimulating factor
	Major histocompatibility complex
	Matrix metalloproteinases
	.Microsatellite instability
	Microsatellite stable
	Nuclear factor kB
NKT	
	N-telopeptides of type 1 collagen
OC	
OP	
	Osteoprotegerin
	Phosphoinositide 3-kinase
	Procollagen type 1 N-terminal propeptide
	Planar cell polarity
PKC	Protein kinase C

# List of Abbreviations Cont...

Abb.	Full term
PQCT	Peripheral Quantitative Computed
-	Tomography
PTH	Parathyroid hormone
	parathyroid hormone-related protein
PYR	- ·
	Quantitative Computed Tomography
-	Quantitative Ultrasound
RA	Rheumatoid arthritis
<b>RANK</b>	Receptor activator of NF-κB
	Receptor activator of NF-κB ligand
<b>RBPs</b>	RNA-binding proteins
Ror	Receptor tyrosine kinase like orphan receptor
	Related orphan receptor gamma
SEFIR	Similar expression of fibroblast growth factor
	genes and IL-17Rs
SERMs	Selective estrogen receptor modulators
SOST	Sclerostin
<b>SPA</b>	Single photon absorptiometry
<b>SXA</b>	Single-energy X-ray absorptiometry
TCF	T cell factor
TCR	T cell receptor
TGF-β	Transforming growth factor-β
Th17	
<b>TNF-α</b>	Tumor necrosis factor-α
TRAF6	Tumor necrosis factor receptor associated
	factor 6
	Ubiquitin-specific peptidase 25
	Women's Health Initiative
	World health organization
WNT	Wingless

## List of Tables

Table No.	Title	Page No.
Table (1):	Common causes of secondary osteopo	prosis 17
<b>Table (2):</b>	World Health Organization criter classification of patients with bone medensity measured by dual-energy absorptiometry	nineral X-ray
<b>Table (3):</b>	Methods of diagnosis of osteoporosis	21
<b>Table (4):</b>	Clinical Risk Factors Considered in F	FRAX 25
<b>Table (5):</b>	Biochemical Markers of Bone Formand Resorption in Urine and Serum.	
<b>Table (6):</b>	Comparison between osteoporotic and non-osteoporotic group as region demographic data	arding
<b>Table (7):</b>	Comparison between osteoporotic an osteoporotic groups regarding vertel hip Examination	bral &
<b>Table (8):</b>	Comparison between Non-osteoporot osteoporotic groups regarding radio data	logical
<b>Table (9):</b>	Comparison between osteoporotic an osteoporotic groups regarding IL-17 level	serum
<b>Table (10):</b>	Correlation between each of IL-17 level, DEXA score, FRAX, clinical laboratory data of participant cases	al and
<b>Table</b> (11):	Correlation between IL-17 serum lev history among studied cases	
<b>Table (12):</b>	Correlation between IL-17 serum lev vertebral & hip examination	el and

# List of Tables Cont...

Table No.	Title	Page No.
Table (13):	Correlation between DEXA score history of smoking & history of fra among studied cases	acture
<b>Table (14):</b>	Correlation between DEXA score vertebral & hip examination	
<b>Table (15):</b>	Correlation between FRAX and hist fracture	v
<b>Table (16):</b>	Correlation between FRAX and context examination among studied cases	
Table (17):	Correlation Study between IL-17 slevel, and DEXA score and FRA participant cases	AX of
Table (18):	Sensitivity, Specificity for each of Score, FRAX, and IL-17 serum lediagnose patients with osteoporosis	vel to
	without osteoporosis	83

## List of Figures

Fig. No.	Title	Page No.
Figure (1):	The bone remodeling process	
Figure (2):	The effect of estrogen on bone menopause and post menopause	-
Figure (3):	Bone remodelling	13
Figure (4):	Wnt signal transduction	16
Figure (5):	Dual energy x-ray absorptiometry	20
Figure (6):	Fracture Risk Assessment Tool	23
Figure (7):	Treatment Pathways According to Characterisation of Fracture Risk	Based
	on FRAX® Probability	
Figure (8):	Treatment Strategies for Osteoporo	
Figure (9):	Interleukin 17 (IL-17) family cyto and their receptors	
<b>Figure (10):</b>	Effects of IL-17 on osteoblast, osteo and chondrocyte differentiation	•
Figure (11):	Summary of interleukin-17A function	
Figure (12):	IL-17 Protective and Path Functions	ologic
Figure (13):	FRAX calculation tool-web- algorithm program input of variables	based FRAX
Figure (14):	Interpretation of fracture risk	
Figure (15):	Box plot showing distribution of carregard DEXA score	ses as
<b>Figure (16):</b>	Box plot showing distribution of carregard FRAX	ses as
<b>Figure (17):</b>	Box plot showing distribution of carregard IL-17 serum level	

# List of Figures Cont...

Fig. No.	Title	Page No.
Figure (18):	Scatter figure showing between age and FRAX	
Figure (19):	Scatter figure showing between Duration of meno	pause and
<b>Figure (20):</b>	Correlation study of serum DEXA score (a) and FRAX inc	
Figure (21):	ROC curve for each of Di FRAX, and IL-17 serum level patients with osteoporosis fr osteoporosis	to diagnose com without

### **Abstract**

**Background:** Osteoporosis is a bone condition that makes bones thinner and more fragile because of reduced bone density and it puts people at risk of fractures, especially of the hip, spinal vertebrae and wrist.

**Aim and objectives:** The aim of work is to study possible relation between serum level of IL-17 and postmenopausal osteoporosis.

**Subjects and methods:** A case control study that was conducted at Physical Medicine, Rheumatology and Rehabilitation department at Ain Shams University hospital including thirty postmenopausal females' study will be diagnosed as having primary post-menopausal" type I". The duration of the study ranged from 6-12 months.

**Results**: results revealed that IL-17 level were 120 (60 - 280) with range (45-480), the mean DEXA Score level was  $1.03 \pm 0.27$  with range (0.62-1.7) and the median FRAX level was 0.55 (0.3 - 1.2) with range (0.1-1.9). There was high statistically significant difference between with and without osteoporosis groups regarding IL-17, DEXA Score and FRAX and there was no statistically significant difference as regard ESR, CRP and PTH.

**Conclusion**: Serum IL-17 was significantly elevated in osteoporotic postmenopausal women when compared to healthy postmenopausal women.

**Keywords:** Postmenopausal, Osteoporosis, IL-17, DEXA Score, Rehabilitation.