



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكرو فيلم

بسم الله الرحمن الرحيم



HANAA ALY



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكروفيلم



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلم



HANAA ALY



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكروفيلم

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HANAA ALY



**“THE EFFECT OF NANO MINERAL TRIOXIDE
AGGREGATE ON THE HEALING PROCESS OF
PERIAPICAL TISSUES AFTER ENDODONTIC SURGERY
AND ITS SEALING ABILITY AND ADAPTABILITY
AS A ROOT-END FILLING MATERIAL”**

Thesis

*Submitted to the Faculty of Dentistry - Ain Shams University
for Partial Fulfillment of the Requirements for Doctorate Degree in
Endodontics*

By

Marwa Wagih Saad ElBoraey

Assistant Lecturer of Endodontics, Future University

(B.D.S, Cairo University-2009, M.D.S, Cairo University-2017)

**Faculty of Dentistry
Ain Shams University
2022**


SUPERVISORS

Prof. Dr. Ihab Hassanien

Professor of Endodontics, Endodontic department,
Faculty of Dentistry
Ain Shams University
Dean of Faculty of Dentistry, Galala University

Dr. Mohamed Mokhtar Nagy

Assistant professor of Endodontics, Endodontic department
Faculty of Dentistry
Ain Shams University



قَالُوا سُبْحَانَكَ
لَا عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا
إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ

صَدَقَ اللَّهُ الْعَظِيمُ

سورة البقرة - آية (٣٢)

Acknowledgement

First, All the gratitude and thanks to Allah, the Most Gracious and the Most Merciful who helped me to accomplish this study.

I would like to express all my heart full of very special thanks and my deepest gratitude and appreciation to my mentor Prof. Dr. Ihab Hassanien, Professor of Endodontics, Faculty of Dentistry, Ain Shams University and Dean of Faculty of Dentistry, Galala University, for his time, effort, advice, valuable comments, and the priceless support he gave to me throughout my journey.

I would like to express my sincere gratitude to Assis. Prof. Dr. Mohamed Mokhtar Nagy, Assistant professor of Endodontics, Faculty of Dentistry, Ain shams university, for his invaluable help, for his time and effort, his support and guidance and his continuous cooperation throughout the details of every part of this study to be completed.

I would like to express my deepest gratitude and thankfulness to each and every patient who took part in this study who gave me chance to complete this whole work,

Last but not least, I would like to thank all professors and staff members in the Endodontics Department in Ain Shams University for their help and support.

Dedication

All grateful words are not enough to express my feelings to:

My Father & Mother,

*who are always supporting me in my life, for their lifelong
inspiration for me to be the best and for their endless love
and support in my hard times and who taught me
perseverance and supported me in all stages of this study*

My Sister, My Brother, My Lovely Friends

For Their endless support, supreme love, and absolute care

My Professors, Colleagues,

*at the Faculty of Dentistry, Future University in Egypt
for standing by my side all through the course of this study*

This work is especially dedicated to

***My Grandfather Saad El-Boraey,**
who is truly missed*

LIST OF CONTENTS

	<i>Page</i>
LIST OF TABLES.....	II
LIST OF FIGURES.....	III
LIST OF ABBREVIATIONS	V
INTRODUCTION	1
REVIEW OF LITERATURE	4
AIM OF THE STUDY	32
MATERIALS & METHODS	34
RESULTS.....	68
DISCUSSION.....	89
SUMMARY	104
CONCLUSION	107
RECOMMENDATIONS	108
REFERENCES	109
APPENDIX	127
ARABIC SUMMARY	-

LIST OF TABLES

Table No.	Title	Page
1	The recorded means, standard deviation (SD) values of the gap thickness between dentin-material interface measurements in (μm) in the longitudinal section of Nano-MTA and MTA groups	72
2	The recorded means, standard deviation (SD) values of the gap thickness between dentin-material interface measurements (μm) in the transverse section of Nano-MTA and MTA groups	75
3	The mean and standard deviation (SD) of age and frequency & percentage of gender for tested groups	77
4	The mean, standard deviation (SD) values of volumetric changes in bone lesion size in (mm^3) of different groups	83
5	The mean, standard deviation (SD) values of percentage of volumetric change in bone lesion size in (mm^3) of different groups	85

LIST OF FIGURES

Figure no.	Title	Page
1	Study design for comparison between MTA and Nano-MTA	35
2	Photograph showing (A) DTE ultrasonic unit, (B) Satelec surgical ultrasonic tip (AS3D) for root-end cavity preparation.....	41
3	A photograph showing a tooth with root-end resection; and a radiograph showing the tooth after ultrasonic root-end preparation	41
4	A diagram showing the grouping of the experimental samples in the leakage and the adaptation study	42
5	A photograph showing ProRoot MTA root-end filling material	43
6	A photograph showing Nano-MTA root-end filling material	43
7	TEM images of the prepared MTA	45
8	XRD pattern of prepared MTA	45
9	A longitudinal section in the apical 3 mm into two halves using a slow-speed diamond saw	49
10	Scanning Electron Microscopic evaluation of sealing ability with 1000x magnification in a longitudinal section	49
11	A transverse section to obtain 1 mm thick section using a slow-speed diamond saw	50
12	Scanning Electron Microscopic evaluation of adaptability with 1000x magnification in a transverse section	50
13	Consort flow diagram of the study	54

Figure no.	Title	Page
14	A photograph showing a final image obtained by CBCT.....	64
15	The sequential steps of measuring the volume of periapical bone lesion from the CBCT by the aid of 3D slicer software	66
16	SEM micrograph for longitudinal section of retro-cavity sealed with Nano-MTA showing good adaptation. Homogenous filling mass is clearly detectable (X 1000)...	69
17	SEM micrograph for longitudinal section of retro-cavity sealed with Nano-MTA showing homogenous material with almost no cracks or voids (X 1000)	69
18	SEM micrograph for longitudinal section of retro-cavity sealed with MTA. The interface between MTA and retro-cavity margins appeared to be relatively good but with thin areas of gaps (X 1000).....	71
19	SEM micrograph for longitudinal section of retro-cavity sealed with MTA showing few cracks and voids within the bulk of the material. (X 1000)	71
20	A bar chart showing the recorded mean values for the gap thickness between dentin-material interface measurements in the longitudinal section of Nano-MTA and MTA groups	72
21	SEM micrograph for transverse section of retro-cavity sealed with Nano-MTA showing good adaptation. Homogenous filling mass with no voids or cracks is clearly detectable (X 1000)	73
22	SEM micrograph for transverse section of retro-cavity sealed with MTA. The interface between MTA and retro-cavity margins appeared to be relatively good but with thin areas of gaps. Figure showing few cracks and voids within the bulk of the material (X 1000)	74

Figure no.	Title	Page
23	A bar chart showing the recorded mean values for the gap thickness between dentin-material interface measurements in the transverse section of Nano-MTA and MTA groups ..	75
24	Bar chart showing percentage of gender distribution in tested groups.....	78
25	Bar chart showing the mean age in tested groups	78
26	Bar chart representing mean values of volumetric changes in bone lesion size within each group	84
27	Bar chart representing mean values of volumetric changes in bone lesion size for different groups	84
28	Bar chart representing percentage of volumetric change in bone lesion size for different groups	86
29	A photograph showing Cone Beam CT scan using Nano-MTA as root-end filling material, (A): The preoperative CBCT image, (B): The 12 months follow-up CBCT image showing decrease in the size of periapical radiolucency....	87
30	A photograph showing Cone Beam CT scan using MTA as root-end filling material, (A): The preoperative CBCT image, (B): The 12 months follow-up CBCT image showing decrease in the size of periapical radiolucency....	88

LIST OF ABBREVIATIONS

Abbreviation	Full term
?????	?????

INTRODUCTION

Surgical endodontic treatment is an option for teeth with apical periodontitis and may be indicated for teeth with periapical pathology when nonsurgical retreatment is impractical or unlikely to improve the previous results or when a biopsy is needed^(1,2). Endodontic surgery procedure consists of the exposure of the involved root end, root end resection and root-end cavity preparation to receive a biocompatible root-end filling material. This root-end resection results in an exposure of the root dentin and the root canal filling material (gutta-percha). The placing of a root-end filling after root-end resection is mandatory to support bone regeneration. Various materials have been advocated for use as retrograde filling. An ideal endodontic retro-filling material should possess numerous characteristics such as biocompatibility to be well tolerated by periapical tissue, adhere, and adapt to the dentinal wall to prevent leakage of microorganisms and their by-products, and allow healing of the periapical tissues. Teeth treated with a root-end filling after apicectomy showed significantly better healing than teeth treated by smoothing of the orthograde gutta-percha root filling only. ⁽³⁾

The main aim of root-end filling material is to be able to prevent the infiltration of bacteria into the root canals⁽⁴⁾, so root-end filling materials should have appropriate sealing ability and adaptability as the amount of microorganisms in the infected area is a key factor determining the healing or persistence of the apical lesion. ⁽⁵⁾