



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكروفيلم

بسم الله الرحمن الرحيم



HANAA ALY



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكروفيلم



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلم



HANAA ALY



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكروفيلم

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HANAA ALY



**Horizontal Ridge Expansion Using Alveolar Ridge
Splitting and Platelet Rich Fibrin (PRF) or
Xenogeneic Graft with Simultaneous Implant
Placement (Randomized Controlled Clinical and
CBCT Study)**

Thesis

**Submitted in Partial Fulfillment of the Requirements of PhD Degree
in Periodontology**

By

Omar Ahmed Abdel Salam Sitten

B.D.S. 2007, M.Sc. 2016

Faculty of Dentistry, Ain Shams University

Supervisors

Prof. Dr. Ahmed Youssef Gamal

Professor of Oral Medicine, Periodontology, Oral Diagnosis and
Radiology, Faculty of Dentistry, Ain Shams University

Vice Dean for Postgraduate Studies and research, Faculty of
Dentistry, 6 October University

Ass. Prof. Dr. Mohamed Sherif El Mofty

Associate Professor of Oral Medicine, Periodontology, Oral
Diagnosis and Radiology Faculty of Dentistry, Ain Shams
University

Vice Dean for Community Service and Environmental
Development, Faculty of Dentistry, Nahda University

Acknowledgment

First thanks to ***Almighty Allah***, for giving me strength to carry on this study and for all his blessings.

I'd like to express a lot of thanks and appreciation to ***Prof. Dr. Ahmed Youssef Gamal***, Professor of Oral Medicine, Periodontology, Oral Diagnosis and Radiology, Faculty of Dentistry, Ain-Shams University for his great valuable supervision and kind guidance.

Very special thanks to ***Ass. Prof. Dr. Mohamed Sherif El Mofty*** Associate Professor of Oral Medicine, Periodontology, Oral Diagnosis and Radiology, Faculty of Dentistry, Ain-Shams University for his great efforts and support throughout this work.

Many thanks and appreciations are also extended to all staff members and colleagues at the oral medicine and periodontology department, Faculty of Dentistry, Ain-Shams University for their support and help.

Finally I would like to express special thanks to Dean and my colleagues at Faculty of Dentistry, MTI University.

Omar Ahmed AbdAlsalam Sitten

Dedication

THIS THESIS IS DEDICATED TO MY FATHER FOR BEING MY
ROLE MODEL AND HIS GREAT HELP AND SUPPORT
THROUGHOUT THIS WORK, MY MOTHER FOR HER ENDLESS
LOVE, I REALLY HONORED TO BE YOUR SON, MY WIFE AND LIFE
PARTNER FOR HER SUPPORT AND LOVE AND TO MY LOVELY
SONS



**التكبير الأفقي لحرف السخية عن طريق التقسيم وباستعمال
البلازما الغنية بالألياف أو الطعم الأجنبي مع وضع غرسات
الأسنان في وقت متلازم (دراسة اكلينيكية تحكيمية و باستخدام
التصوير الاشعاعي ثلاثي الأبعاد)**

رسالة

**تمهيدية مقدمة لكلية طب الفم والأسنان جامعة عين شمس توطئة
للحصول على درجة الدكتوراة في علاج اللثة**

مقدمة من

الطبيب/ عمر أحمد عبد السلام ستين

**بكالوريوس طب وجراحة الفم والأسنان
ماجستير طب الفم وعلاج اللثة**

تحت إشراف

أ.د/ أحمد يوسف جمال

أستاذ طب الفم وأمراض اللثة والتشخيص والأشعة

كلية طب الأسنان – جامعة عين شمس

وكيل الكلية لشئون الدراسات العليا كلية طب الأسنان – جامعة 6 أكتوبر

أ.م.د/ محمد شريف المفتي

أستاذ مساعد طب الفم وأمراض اللثة و التشخيص و الأشعة

كلية طب الأسنان – جامعة عين شمس

وكيل الكلية لشئون خدمة المجتمع وتنمية البيئة كلية طب الأسنان – جامعة النهضة

كلية طب الأسنان

جامعة عين شمس

2022

List of Contents

Title	Page No.
List of Tables	II
List of Figures	IV
List of Abbreviations	IX
INTRODUCTION AND REVIEW OF LITERATURE	1
AIM OF THE STUDY	38
MATERIALS, SUBJECTS AND METHODS	39
RESULTS	74
DISCUSSION	111
CONCLUSIONS	123
RECOMMENDATIONS	124
SUMMARY	125
REFERENCES	130
ARABIC SUMMARY	-

List of Tables

Table No.	Title	Page No.
1	Descriptive data of all studied cases (16 cases) with horizontal bone defect managed by ridge splitting procedure with simultaneous implant placement.	74
2	Plaque index of both groups preoperative as well as 6 and 9 months postoperatively.	77
3	Gingival index of both groups preoperative as well as 6 and 9 months postoperatively	80
4	Follow up results of soft tissue thickness in group I preoperatively as well as 6 months and 9 months postoperatively and their statistical significance.	83
5	Follow up results of soft tissue thickness in group II preoperatively as well as 6 months and 9 months postoperatively and their statistical significance.	86
6	Follow up results of bucco-lingual bone width in group I preoperatively as well as immediately, 6 and 9 months postoperatively and their statistical significance.	88
7	Follow up results of bucco-lingual bone width in group II preoperatively as well as immediately, 6 and 9 months postoperatively and their statistical significance.	91
8	Follow up results of marginal bone level in group I immediately as well as 6 months and 9 months postoperatively and their statistical significance.	93

<i>Table No.</i>	<i>Title</i>	<i>Page No.</i>
9	Follow up results of marginal bone level in group II immediately as well as 6 months and 9 months postoperatively and their statistical significance.	95
10	Comparison of mean value of studied parameters in both groups with horizontal bone defect managed by ridge splitting with simultaneous implant placement	97
11	Marginal bone level in relation to implant level in both studied groups.	102
12	Comparison between both groups regarding value of change of studied parameters preoperatively, immediately, 6 and 9 months postoperatively	104
13	Clinical and radiographic parameters of cases of Group 1 (PRF membrane) preoperatively, at 6 and 9 months postoperatively.	109
14	Clinical and radiographic parameters of cases of Group 2 (Xenogenic bone particulates) preoperatively, at 6 and 9 months postoperatively.	110

List of Figures

<i>Fig. No.</i>	Title	Page No.
1	Edentulous space of missing upper right canine and first premolar.	56
2	Full thickness flap elevation.	56
3	Ridge splitting with one horizontal cut and two vertical cuts at premolar area (Buccal view).	57
4	Ridge splitting with one horizontal cut and two vertical cuts at premolar area (Occlusal view).	57
5	Implant placement in splitted ridge at premolar area.	58
6	PRF membrane prepared from patient blood.	58
7	Placement of PRF membrane in intercortical gap.	59
8	Suturing of flap edges by interrupted sutures.	59
9	Implant exposure after 6 months and placement of healing abutment for 2 weeks.	60
10	Porcelain fused to metal crown with occlusal hole to restore upper right first premolar (Occlusal view).	60
11	Porcelain fused to metal crown with occlusal hole to restore upper right first premolar (Buccal view).	61

Fig. No.	Title	Page No.
12	Preoperative CBCT to measure bucco-palatal bone width.	61
13	Immediately postoperative CBCT to measure bucco-palatal bone width & marginal bone height.	62
14	Six months postoperative CBCT to measure bucco-palatal bone width & marginal bone height.	62
15	Nine months postoperative CBCT to measure bucco-palatal bone width & marginal bone height.	63
16	Panoramic view of implant at 6 and 9 months postoperatively.	63
17	Edentulous space of missing upper right canine and first premolar.	64
18	Full thickness flap elevation.	64
19	Ridge splitting using bone chisels and mallet to separate ridge.	65
20	Parallel pins to check implants positions after initial drilling.	65
21	Bone expander for expansion of bone at osteotomy site.	66
22	Implants placement in splitted ridge at canine and 1st premolar area.	66
23	Placement of xenogenic bone particulate in intercortical gap then covered by collagen membrane.	67

Fig. No.	Title	Page No.
24	Membrane stabilization by horizontal mattress suture.	67
25	Suturing of flap edges by interrupted sutures.	68
26	Implants exposure after 6 months and placement of healing abutments for 2 weeks.	68
27	Implants stock abutments were screwed before crowns cementation.	69
28	Porcelain fused to metal crowns to restore upper right canine and 1st premolar (Occlusal view).	69
29	Porcelain fused to metal crowns to restore upper right canine and 1st premolar (Buccal view).	70
30	Preoperative CBCT to measure bucco-palatal bone width.	70
31	Immediately postoperative CBCT to measure bucco-palatal bone width & marginal bone height.	71
32	Six months postoperative CBCT to measure bucco-palatal bone width & marginal bone height.	71
33	Nine months postoperative CBCT to measure bucco-palatal bone width & marginal bone height.	72
34	Panoramic view of implants at 6 and 9 months postoperatively.	72
35	Mean age in years in both groups.	75
36	Sex distribution of cases in both groups.	75

Fig. No.	Title	Page No.
37	Plaque index preoperative and 9 months postoperatively.	78
38	Gingival index preoperatively and postoperatively 9 months.	81
39	Soft tissue thickness of cases in group I preoperatively as well as 6 months and 9 months postoperatively.	84
40	Soft tissue thickness of cases in group II preoperatively as well as 6 months and 9 months postoperatively.	86
41	Bone width of cases in group I preoperatively as well as immediately, 6 months and 9 months postoperatively.	89
42	Bone width of cases in group II preoperatively as well as immediately, 6 months and 9 months postoperatively.	91
43	Marginal bone level of cases in group I immediately as well as 6 months and 9 months postoperatively.	93
44	Marginal bone level of cases in group II preoperatively as well as immediately, 6 months and 9 months postoperatively.	95
45	Buccolingual bone mean width in both groups preoperatively, 6 month and 9 months postoperatively.	98
46	Soft tissue thickness mean in both groups preoperatively and 9 months postoperatively.	99

<i>Fig. No.</i>	Title	Page No.
47	Marginal bone level in relation to implant level immediate and 9 months postoperative in both groups.	100
48	Variations in buccolingual bone width among cases of group I.	105
49	Variations in buccolingual bone width among cases of group II.	105
50	Variations in marginal bone level in cases of group I.	106
51	Variation of marginal bone level in cases of group II.	106
52	Comparison between both groups regarding value of change of studied parameters preoperatively or immediately and 9 months postoperatively.	107