

بسم الله الرحمن الرحيم





شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بعض الوثائق الأصلية تالفة





بالرسالة صفحات
لم ترد بالأصل



B1✓10V

The Impact of Environmental and Parental Conditions on Mass Production and Rearing Process of *Oreochromis Species* and Their Hybrid

A THESIS

***Submitted to the Faculty of Science
Zoology Department
Alexandria University
In Fulfillment of the Requirements for
The Degree of
Ph.D.***

By

Samir Guindy Ghabrial

National Institute of Oceanography & Fisheries

*B.Sc. (Alexandria University)
Dip. EST (Delft – The Netherlands)
M.Sc. (Alexandria University)*

Supervised by

Prof. Dr. Milad I. Michael

*Emeritus Professor of Experimental
Embryology Faculty of Science
Alexandria University*

Prof. Dr. Magda I. Zaki

*Professor of Aquaculture
National Institute of Oceanography
and Fisheries, Alexandria*

*To whom I owe a great debt of love
and thanks,
my late mother
Sophia*

Acknowledgement

The author is most grateful to Prof. Dr. Magda F. Zaki, Professor of Aquaculture, National Institute of Oceanography and Fisheries, Alexandria, for her constant supervision and extensive help. Her valuable advice, generous guidance, assistance, helpful criticism and follow-up, made this work a reality.

I'd like to express my deepest and sincere gratitude to Professor Dr. Milad F. Michael, Professor of Experimental Embryology for his great care, help and concern throughout the course of this work. His stimulating influence was a motive to push me forward.

Thanks for both of you.

Table of Contents

	<i>Page</i>
List of Tables	iii
List of Figures	vi
Introduction and Literature Review	1
Binomial System of Tilapia	2
Identification and Biochemical Analysis	3
Spawning and Environmental Aspects	5
Cage Culture	6
Hybridization	7
Objectives	8
Materials and Methods	10
I. Sampling and Identification	10
1- Morphometric Identification	10
2- Biochemical Identification	11
II. Fish Spawning Techniques and Seed Production	11
1- Induced Spawning Techniques	12
2- Spawning in Concrete Tanks at El-Max Fish Culture Station	13
3- Spawning in Fiber Glass Tanks at the NIOF	14
4- Spawning in Concrete Basins at Kafr El-Hataba	16
III. Net-Enclosures and Cage Culture	19
Design and Construction	21
Food Composition and Processing of Diet for Fingerlings	23
Stocking Rates	23
IV. Biochemical Analysis	24
1- Fast-System Iso-Electric Focusing	24
2- Phast-System Electrophoresis	27
3- Total Amino Acid Extraction	29

Results and Observations	Page 31
I. Species Identification	31
1- Morphometric Identification	31
2- Biochemical Identification	36
II. Spawning, Seed Production and Cage Culture	44
1- Spawning in Concrete and Fiber Glass Tanks at El-Max and NIOF	44
2- Spawning in Concrete Basins at Kafr El-Hataba	47
3- Culture in Net-Enclosures	49
4- Culture in Cages	57
III. Biochemical Analysis	63
1- Fast-System Isoelectric Focusing (IEF)	63
* IEF of Muscle Protein	63
* IEF of Plasma Protein	70
* IEF of Hepatic Protein	76
* IEF of Flesh of Fingerlings	82
* IEF of Gonadal Protein	88
2- Phast-System Electrophoresis	
* Muscle Protein	94
* Hepatic Protein	99
* Gonadal Protein	103
* Flesh of fingerlings	107
3- Total Muscle Protein and Amino Acid Content	112
Discussion	120
Summary and conclusion	150
References	159

List of Tables

<i>Table No.</i>	<i>Description</i>	<i>Page</i>
1	Summary of the external characteristics distinguishing both pure parent species, i.e. <i>O. niloticus</i> and <i>O. aureus</i> found naturally	35
2	The isoelectric points (PI's) values and the percentages of quantitative values for scanning flesh and gill protein fractions of two <i>Oreochromis niloticus</i> and <i>Oreochromis aureus</i> and their natural hybrid	42
3	Spawning in Concrete and Fiberglass Tanks at El-Max Fish Culture Station of NIOF (March – Oct. 1993)	46
4	Spawning and Environmental Conditions in Concrete Basins at Kafr El Hataba (Dec. 1992 – Sep. 1993)	48
5	Spawning and Environmental Conditions in Fiberglass Tanks Indoors at the Semi-commercial Hatchery NIOF (March – Oct. 1993)	48
6	Average physico-chemical parameters and nutrient status in the net-enclosures at El-Max Research Station (monthly measured during the 180 days rearing period)	52
7	Percentage composition and chemical analysis of the fingerlings diet (25% protein content pelleted diet) used in both net enclosures at El Max fish farm and cage culture at Kafr El Hataba.	53
8	Percentage composition and chemical analysis of the fingerlings diet (31% protein content pelleted diet) used in both net enclosures at El Max fish farm and cage culture at Kafr El Hataba.	53
9	Vitamin mixture concentration used in both protein diets expressed in IU or gm / ton of the whole ingredient	54
10	Mineral mixture concentrations used in both protein diets expressed in gm / ton of the whole ingredient	54
11	Mean of body weight, total length and condition factor (K) of <i>Oreochromis spp.</i> & their hybrid fingerlings receiving different feeds in 1.6 m ³ net-enclosures for 180 days (April – September).	55

Table No.	Description	Page
12	Growth parameters, food utilization and survival rate of <i>Oreochromis spp.</i> & their hybrid reared in net enclosures using various feeds for 180 days (April – September).	56
13	Average monthly parameters and nutritional status in the cage's activities at Kafr El Hataba (during six month rearing period).	60
14	Mean of body weight, total length and condition factor (K) of <i>Oreochromis spp.</i> and their hybrid fingerlings receiving different feed in cages for 180 days	61
15	Growth performance, feed utilization and survival rates of <i>Oreochromis spp.</i> & their hybrid reared in cages using various feed for 180 days	62
16	The isoelectric points (PI's) values and the percentage of quantitative values for scanning of white muscle protein fractions of <i>Oreochromis spp.</i> and their hybrid.	68
17	The isoelectric points (PI's) values and the percentage of quantitative values for scanning of plasma protein fractions of <i>Oreochromis spp.</i> and their hybrid.	74
18	The isoelectric points (PI's) values and the percentage of quantitative values for scanning of liver (hepatic) protein fractions of <i>Oreochromis spp.</i> and their hybrid.	80
19	The isoelectric points (PI's) values and the percentage of quantitative values for scanning of the flesh protein fractions of fingerlings of <i>Oreochromis spp.</i> and their hybrid.	86
20	The isoelectric points (PI's) values and the percentage of quantitative values for scanning of gonad's protein fractions of <i>Oreochromis spp.</i> and their hybrid.	92
21	Relative mobility (Rf) with their corresponding molecular weight (Mwt.) and the percentage of quantitative values for scanning protein fraction in the white muscle of <i>Oreochromis spp.</i> and their hybrid	98
22	Relative mobility (Rf) with their corresponding molecular weight values (Mwt.) and the percentage of quantitative values for scanning protein fraction in the hepatic protein of <i>Oreochromis spp.</i> and their hybrid	102

Table No.	Description	Page
23	Relative mobility (Rf) with their corresponding molecular weight values (Mwt.) and the percentage of quantitative values for scanning protein fractions in testis and ovaries of <i>Oreochromis spp.</i> and their hybrid	106
24	Relative mobility (Rf) with their corresponding molecular weight values (Mwt.) and the percentage of quantitative values for scanning protein fraction in the flesh of the fingerlings of <i>Oreochromis spp.</i> and their hybrid	111
25	Percentages of muscle composition on wet weight basis of <i>Oreochromis spp.</i> and their hybrid	114
26	Total amino acids content in the flesh of both pure strains of <i>Oreochromis spp.</i> and their resulting hybrids expressed in gram / 100 g.f.w.	115
27	Total and mean values of amino acids contents of <i>Oreochromis spp.</i> and the hybrids and the mean values of each amino acid in the three groups of females only, males only and hybrids expressed in gram / 100 g.f.w.	118

List of Figures

Figure no.	Description	Page
1	Five brick concrete tanks for spawning experiments (at El-Max). Each tank has its own gate – the five tanks with a main reservoir.	15
2	Fiberglass tanks (1.5 ton water capacity) for hybrid production at El-Max.	15
3	The three identical concrete basins at Kafr El-Hataba covered with semi-transparent plastic sheets	17
4	The semi-transparent plastic sheets stretched on flexible metal tubes covering the unit completely	17
5	Water pump as the reservoir supplier at the top of the basins	18
6	Clay porcelain tubes used as nests during fish spawning and incubation	18
7	Net enclosures used for supplemental feeding of fish fry (experimental)	20
8	Another type of net enclosures used for supplemental feeding of fish fry (experimental)	20
9, 10, 11	Floating or walkway type cage – in Nile estuary at Kafr El-Hataba (commercial)	22
12	<i>Oreochromis niloticus</i>	34
13	<i>Oreochromis aureus</i>	34
14(a)	IEF of phast gel (3-9) of protein pattern in muscle of <i>Oreochromis species</i> . a (<i>O. niloticus</i> adult), b(<i>O. aureus</i> adult), c & d (<i>O. niloticus</i> fry), e (hybrid fingerlings) and St (standard protein)	37
14(b)	IEF of phast gel (3-9) of flesh protein in <i>Oreochromis species</i> a, c and e (flesh of hybrid, <i>O. niloticus</i> and <i>O. aureus</i>) b, d and f (gills of hybrid, <i>O. niloticus</i> and <i>O. aureus</i> respectively) St (standard protein from 3-10)	37

Figure no.	Description	Page
15	The electrophoretograms and densitograms of flesh and gills of <i>Oreochromis species</i> a, b, c, d, e and f (muscle of <i>O. niloticus</i> , <i>O. aureus</i> , hybrid <i>O. niloticus</i> fingerlings, <i>O. aureus</i> fingerling and hybrid fingerling) g, h and i (gills of <i>Oreochromis niloticus</i> , <i>Oreochromis aureus</i> and hybrid)	38
16-a	IEF of phast-gel (3-9) of white muscle protein pattern of <i>Oreochromis species</i> and their hybrid	66
16-b	The electrophortogram and densitogram of phast gel isoelectric focusing for white muscle protein of <i>Oreochromis species</i> and their hybrid	67
17-a	IEF of phast-gel (3-9) of plasma protein pattern of <i>Oreochromis species</i> and their hybrid	72
17-b	The electrophortogram and densitogram of phast gel isoelectric focusing for plasma protein of <i>Oreochromis species</i> and their hybrid	73
18-a	IEF of phast-gel (3-9) of hepatic protein pattern of <i>Oreochromis species</i> and their hybrid	78
18-b	The electrophortogram and densitogram of phast gel isoelectric focusing for hepatic protein of <i>Oreochromis species</i> and their hybrid	79
19-a	IEF of phast-gel (3-9) of flesh protein pattern of fingerlings of <i>Oreochromis species</i> and their hybrid	84
19-b	The electrophortogram and densitogram of phast gel isoelectric focusing of fingerling's protein of <i>Oreochromis species</i> and their hybrid	85
20-a	IEF of phast-gel (3-9) of gonadal protein pattern of <i>Oreochromis species</i> and their hybrid	90
20-b	The electrophortogram and densitogram of phast gel isoelectric focusing for gonadal protein of <i>Oreochromis species</i> and their hybrid	91
21-a	Polyacrlamide gel electrophoresis (PAGE) phast-gel gradient (8-25) of white muscle protein pattern of <i>Oreochromis species</i> and their resulting hybrid.	96